

# Dietary Folate Intake and Breast Cancer Risk: Results from the Shanghai Breast Cancer Study<sup>1</sup>

Martha J. Shrubsole, Fan Jin, Qi Dai, Xiao-Ou Shu, John D. Potter, James R. Hebert, Yu-Tang Gao, and Wei Zheng<sup>2</sup>

Division of General Internal Medicine and Vanderbilt-Ingram Cancer Center, Vanderbilt University, Nashville, Tennessee 37232 [M. J. S., Q. D., X-O. S., W. Z.]; Department of Epidemiology and Biostatistics, Norman J. Arnold School of Public Health, University of South Carolina, Columbia, South Carolina 29208 [M. J. S., J. R. H.]; Cancer Prevention Research Program, Fred Hutchinson Cancer Research Center, Seattle, Washington 98109 [J. D. P.]; and Shanghai Cancer Institute, Shanghai, China 200032 [F. J., Y-T. G.]

## ABSTRACT

Folate is involved in DNA synthesis, repair, and methylation. It has been hypothesized that high intake of folate may reduce the risk of human cancers, including cancer of the breast. Using data from a population-based case-control study of breast cancer conducted in urban Shanghai during 1996–1998, we evaluated the association of dietary folate intake and breast cancer risk among 1321 cases and 1382 controls, 25–64 years of age, who never drank alcohol regularly or used vitamin supplements. Usual dietary habits were assessed with an in-person, interviewer-administered food frequency questionnaire developed and tested for use in this population. Unconditional logistic regression models were used to calculate odds ratios (ORs) and their 95% confidence intervals (95% CIs) after adjusting for potential confounding factors. Dietary folate intake was inversely associated with breast cancer risk (*P* for trend, 0.05) with an adjusted OR of 0.71 (95% CI, 0.56–0.92) observed among women who were in the highest quintile of intake. The inverse association was stronger after further adjusting for total fruit and vegetable and animal food intakes (OR, 0.62; 95% CI, 0.46–0.82; *P* for trend, 0.01). A more pronounced inverse association between folate intake and breast cancer risk (OR, 0.47; 95% CI, 0.25–0.88; *P* for trend, 0.01) was observed among women who consumed high levels of folate cofactors (methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub>) than those whose intake levels of these nutrients were low. Dietary intake of methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub> were not independently related to risk of breast cancer after adjusting for confounding factors. Thus, our study adds additional support to the protective role of dietary folate in breast carcinogenesis and suggests further that the effect of folate may be modified by dietary intake of methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub>.

## INTRODUCTION

Folate, a B vitamin found naturally in many food sources, particularly in dark green leafy vegetables, is essential for regenerating methionine, the methyl donor for DNA methylation, and for producing the purines and pyrimidine thymidylate required for DNA synthesis and repair. The role of folate is well documented in the prevention of neural tube defects and is increasingly studied in relation to cardiovascular disease risk (1, 2). Evidence for its potential role in carcinogenesis is encouraging, although the effect is less well studied (3). Several studies have examined the association of dietary folate with colorectal neoplasms (4–21), and most of them reported a statistically significant inverse association. However, only a few studies have examined the association of dietary folate intake with breast cancer risk, and the results from these studies have been inconsistent (22–28). With the exception of only one study (26), the observed inverse association was restricted in subgroups of women, and many of the reported associations were not statistically significant. One

study found that folate intake was related to a reduced risk of breast cancer only among premenopausal women (24), whereas others found that the inverse association was restricted to postmenopausal women (23, 27), particularly those who drank alcohol regularly. One nested case-control study, using stored serum samples, found no association between serum folate and breast cancer risk (29). Almost all of the previous studies were conducted in predominantly Caucasian North American populations.

Although two previous studies have evaluated folate according to methionine intake (27, 28), no previous study has evaluated the potential modifying effects of vitamin B<sub>12</sub> and vitamin B<sub>6</sub>. All of these nutrients have important roles in folate metabolism. The methyl group for homocysteine remethylation to methionine is donated by folate. Vitamins B<sub>6</sub> and B<sub>12</sub> are obligate cofactors; vitamin B<sub>6</sub> is a coenzyme for the conversion to the form of folate used for DNA synthesis or methylation, and vitamin B<sub>12</sub> facilitates the methyl transfer to homocysteine. Therefore, it is conceivable that a high intake of these folate cofactors and/or their interaction with folate may reduce breast cancer risk. The only published study examining these cofactors found an inverse association between serum B<sub>12</sub> and homocysteine and breast cancer risk (29).

Since 1998, folic acid fortification of cereal grain foods has been mandated in the United States (30); prior to this time, fortification was optional and not uniformly implemented. Therefore, studies of folate intake conducted in North American populations during the past decade may have been limited by the lack of an appropriate food composition database and, thus, by an inability to assess usual folate intake. Some previous studies were also not able to account for intake of folic acid, a more readily bioavailable form of folate than folate in food and present in vitamin supplements; this may have resulted in misclassifying the intake for some subjects.

To evaluate the association between dietary folate intake and breast cancer risk, we analyzed data from a large population-based case-control study conducted in Shanghai, China. Unlike their Western counterparts, most Chinese women have diets composed mainly of unprocessed and unfortified foods, and they rarely take vitamin supplements; this facilitates assessment of folate intake and reduces potential misclassification. Also few women drink regularly. These unique features provide us a better opportunity to evaluate the hypotheses relating dietary folate intake to breast cancer risk than would be available in the United States and other Western countries.

## SUBJECTS AND METHODS

**Subject Recruitment.** The Shanghai Breast Cancer Study is a population-based case-control study conducted in urban Shanghai, China during 1996–1998. Detailed study methods have been published elsewhere (31). Briefly, cases were identified using a rapid-case ascertainment system, supplemented by the Shanghai Cancer Registry. All incident breast cancer cases, newly diagnosed during the study period, and meeting the following criteria were eligible for this study: 25–64 years of age, resident of urban Shanghai, no previous history of any cancer, and alive at the time of interview. A total of 1602 eligible breast cancer cases were identified during the study period, of

Received 5/10/01; accepted 8/10/01.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup>This research was supported by USPHS Grant RO1CA6427 from the National Cancer Institute.

<sup>2</sup>To whom requests for reprints should be addressed, at Center for Health Services Research, Vanderbilt University, Nashville, TN 37232-8300. Phone: (615) 936-0682; Fax: (615) 936-1269; E-mail: wei.zheng@mcmail.vanderbilt.edu.

which 1459 cases (91.1%) completed in-person interviews. The major reasons for nonparticipation were refusal (109 cases, 6.8%), death prior to interview (17 cases, 1.1%), and inability to locate (17 cases, 1.1%). Written informed consent was obtained from all subjects, and relevant committees for the use of human subjects in research approved the study protocol. Two senior pathologists confirmed all diagnoses.

Controls had inclusion criteria identical to those of the cases with the exception of a breast cancer diagnosis. Controls were randomly selected and were frequency matched on age (5-year intervals) to the expected age distribution of the cases in a 1:1 ratio. Controls were selected using the Shanghai Resident Registry, a population registry containing address and demographic information for all residents of urban Shanghai. Of the 1724 eligible controls, 1556 (90.3%) completed in-person interviews. The remaining women were not included in the study either because of refusal (166 controls, 9.6%) or death prior to interview (2 controls, 0.1%).

**Data Collection and Nutrient Estimation.** All subjects completed an in-person interview that used a structured questionnaire and included anthropometric measurements. Dietary intakes were assessed using a 76-item food frequency questionnaire that ascertained mostly single food items as raw ingredients. Only 6 of the food items were processed foods (tofu, fried tofu, breads, noodles, candy, and desserts). The food frequency questionnaire was designed to capture >85% of foods consumed by Shanghai residents and to assess usual diet over the past 5 years, ignoring any recent change. Typically studies conducted in the United States only estimate intake over the past 12 months, which could be problematic, particularly when cases are interviewed several months after cancer diagnosis. Each subject was asked about the frequency with which she ate a specific food (daily, weekly, monthly, yearly, or never), followed by a question on the raw amount typically eaten; *i.e.*, how many lians (1 lian is 50 g) or jins (1 jin is 500 g). For all seasonal foods, subjects were asked to describe their consumption when the food was available on the market. Only 306 cases and controls reported drinking alcoholic beverages regularly ( $n = 122$ ; 4.1%) or regularly using a vitamin B or multivitamin supplement ( $n = 188$ ; 6.2%). Alcohol use was missing for 6 (0.2%) subjects. Because alcohol consumption may increase folate requirements and because the data on the folate content of vitamins were not available, all analyses were limited to the 1321 (90.5%) cases and 1382 (88.8%) controls who were known not to consume alcohol and not to take vitamin supplements. Total dietary intakes of folate, methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub> were calculated by summing the product of the micronutrient content of each food item, usual portion consumed, and frequency of consumption. For each food item listed in the food frequency questionnaire, an identical (82%) or equivalent (17%) item from the food composition database of the USDA was identified, and the values for folate, methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub> were used in the estimate of intake levels of these nutrients. One minor food item (conch) was excluded because of no comparable item in the USDA food database.

**Data Analysis.** Quintile distributions among controls were used to categorize all dietary intake variables. Pearson correlation coefficients were used to measure the correlation between all dietary intake variables and daily fruit and vegetable and animal intakes. ORs<sup>3</sup> were used to measure the association of breast cancer risk with dietary intake. Unconditional logistic regression models were used to obtain maximum likelihood estimates of the ORs and their 95% CIs, after adjusting for potential confounding variables. Risk factors identified previously as having an independent association with breast cancer in this population were controlled in all models (31). These included breast cancer in a first-degree relative, history of fibroadenoma, age at menarche, age at first live birth, age at menopause, BMI, physical activity, and menopausal status. Models also controlled for age, education, and household income. Age was included as a continuous variable throughout, and categorical variables were treated as indicator variables in the model. Energy adjustment was performed using the residual method (32). Other dietary factors were also controlled in some analyses. Tests for trend were performed by entering categorical variables as continuous. The folate cofactor score was calculated by summing the ranks in the tertiles for each of methionine, vitamin B<sub>6</sub>, and vitamin B<sub>12</sub>. Analyses stratified by each folate cofactor and the cofactor score were used to

evaluate their potential modifying effects. Tests for multiplicative interaction were done by including a multiplicative variable between two variables in the logistic model. All statistical tests were based on two-sided probabilities using SAS, version 8.0 (SAS Institute, Inc., Cary, NC).

## RESULTS

Comparisons between cases and controls on selected demographic factors and established breast cancer risk factors are presented in Table 1. Results were very similar between the total sample and among the subset used in the folate analysis (data not shown). Compared with controls, cases had higher education, an earlier age at menarche, a later age at first live birth, a later age at menopause, and a higher daily intake of animal foods. Cases were more likely to have a higher BMI or waist:hip ratio, a family history of breast cancer among first-degree relatives, a history of fibroadenoma, and were less likely to exercise regularly than controls.

Mean and median dietary intake values for total energy, folate, and folate cofactors (methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub>) among controls are presented in Table 2. Average intake according to plant and animal source is also presented. Plant foods were the major source for folate (88%) and vitamin B<sub>6</sub> (72%). Animal foods were the major source of vitamin B<sub>12</sub> (99%). Methionine was obtained approximately equally from both plant and animal foods. Daily intake of fruits and vegetables was moderately correlated with total folate ( $r = 0.56$ ) and methionine ( $r = 0.40$ ) and closely correlated with total B<sub>6</sub> intake ( $r = 0.75$ ). Daily intake of animal foods was closely correlated with methionine ( $r = 0.81$ ) and vitamin B<sub>6</sub> ( $r = 0.62$ ) and moderately associated with folate ( $r = 0.36$ ) and vitamin B<sub>12</sub> (0.40). The correlations between folate and its cofactors were 0.62, 0.24, and 0.73 for methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub>, respectively (data not shown).

As shown in Table 3, breast cancer risk was significantly inversely associated with dietary folate intake (OR for highest *versus* lowest quintile of intake, 0.71; 95% CI, 0.56–0.92; *P* for trend, 0.05). This

Table 1 Distribution of cases and controls by descriptive characteristics for a subset of women in the Shanghai Breast Cancer Study who were known not to drink alcohol or take vitamin supplements, 1996–1998

Subject characteristic	Cases ( <i>n</i> = 1321)	Controls ( <i>n</i> = 1382)	<i>P</i> <sup>a</sup>
Age (yr), mean ± SD	48.1 ± 7.9	46.9 ± 8.7	0.02
Education, %			
No formal education	3.8	5.3	
Elementary school	8.2	8.4	
Middle or high school	75.6	76.7	
College or above	12.5	9.6	0.04
Household income (Yuan), %			
<4000	20.1	19.0	
4000–5999	32.8	32.9	
6000–7999	12.5	13.2	
8000–8999	20.1	23.3	
9000+	14.6	11.5	0.08
Breast cancer in first-degree relative, %	3.6	2.2	0.02
Ever had breast fibroadenoma, %	9.9	4.9	<0.01
Age at menarche (yr)	14.5 ± 1.6	14.7 ± 1.7	<0.01
Ever had a live birth, %	95.2	96.6	0.06
Number of live births, mean ± SD	1.5 ± 0.8	1.5 ± 0.8	0.32
Age at first live birth (yr), mean ± SD	26.8 ± 4.0	26.3 ± 3.8	<0.01
Postmenopausal, %	33.6	34.7	0.54
Age at menopause (yr), mean ± SD	48.0 ± 4.5	47.5 ± 4.6	0.07
Physically active past 10 years, %	17.6	23.5	<0.01
BMI (kg/m <sup>2</sup> ), mean ± SD	23.5 ± 3.4	23.2 ± 3.3	0.01
Waist:hip ratio, mean ± SD	0.81 ± 0.06	0.80 ± 0.05	<0.01
Daily animal food intake (g), mean ± SD	97.9 ± 50.9	90.1 ± 45.6	<0.01
Daily plant food intake (g), mean ± SD	287 ± 158	286 ± 163	0.92
Daily energy intake (kcal), mean ± SD	1863 ± 458	1840 ± 462	0.22

<sup>a</sup> For  $\chi^2$  test (categorical variables) or *t* test (continuous variables).

<sup>3</sup> The abbreviations used are: OR, odds ratio; CI, confidence interval; BMI, body mass index; THF, tetrahydrofolate; SAM, S-adenosyl-methionine; NHS, Nurses' Health Study; USDA, United States Department of Agriculture.

Table 2 Dietary intake of folate and selected nutrients among controls, Shanghai Breast Cancer Study, 1996–1998

Dietary factor	Mean ± SD	Median (25th, 75th percentiles)	Correlation coefficients with intake of	
			Fruits and vegetables	Animal foods
Folate (μg/day)				
Total	294 ± 172	256 (194, 345)	0.56	0.36
Plant source	259 ± 166	220 (163, 302)	0.55	0.28
Animal source	34 ± 22	31 (19, 44)	0.22	0.70
From plants, %	87 ± 7	88 (83, 92)		
Vitamin B <sub>12</sub> (μg/day)				
Total	4.59 ± 4.14	2.85 (1.32, 8.47)	0.19	0.40
Plant source	0.20 ± 0.47	0.04 (0.01, 0.19)	0.28	0.08
Animal source	4.39 ± 4.09	2.62 (1.16, 8.36)	0.16	0.39
From animals, %	93 ± 12	99 (93, 100)		
Methionine (g/day)				
Total	1.64 ± 0.56	1.55 (1.27, 1.89)	0.40	0.81
Plant source	0.81 ± 0.27	0.77 (0.65, 0.93)	0.38	0.26
Animal source	0.83 ± 0.42	0.76 (0.52, 1.03)	0.29	0.90
From animals, %	49 ± 13	50 (41, 57)		
Vitamin B <sub>6</sub> (mg/day)				
Total	1.73 ± 0.58	1.64 (1.35, 1.99)	0.75	0.62
Plant source	1.24 ± 0.45	1.16 (0.94, 1.42)	0.83	0.30
Animal source	0.50 ± 0.26	0.45 (0.32, 0.62)	0.23	0.86
From plants, %	72 ± 10	72 (65, 78)		

association was further strengthened after adjusting for fruit and vegetable intake and animal food intake (OR, 0.62; 95% CI, 0.46–0.82; *P* for trend, 0.01). Analyses by food source of folate revealed similar associations for both plant and animal sources. The ORs were 1.00, 0.77, 0.89, 0.80, and 0.67 (*P* for trend, 0.02) with increasing plant source folate intake and 1.00, 0.93, 0.85, 0.70, and 0.66 (*P* for trend, <0.01) with increasing animal source folate intake after adjusting for fruit, vegetable, and animal food intakes. A high intake of methionine (*P* for trend, 0.01) was associated with a 30% increased risk of breast cancer; the positive association disappeared after adjusting for animal food intake. A 40% elevated risk was observed with a high intake of vitamin B<sub>6</sub>. Analyses by source of vitamin B<sub>6</sub> showed no association with plant source B<sub>6</sub> (*P* for trend, 0.99) and a positive association with animal source B<sub>6</sub> (*P* for trend, <0.01). Additionally adjusting for animal food intake attenuated the positive association (*P* for trend, 0.89). The ORs for increasing animal source vitamin B<sub>6</sub> were 1.00, 0.98, 1.04, 0.92, and 1.09. No association between vitamin B<sub>12</sub> and breast cancer risk was observed. Stratified analyses by menopausal status showed that similar patterns of associations existed in both pre- and postmenopausal women, although the inverse association with folate intake was more evident in postmenopausal (*P* for trend, 0.03) than in premenopausal women (*P* for trend, 0.57; data not

shown). The adjusted ORs for the highest versus the lowest quintile of folate intake were 0.78 (95% CI, 0.56–1.07) in premenopausal women and 0.66 (95% CI, 0.44–0.99) in postmenopausal women.

Stratified analyses were performed to evaluate potential modifying effects of folate cofactors (methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub>) on the association between dietary folate intake and breast cancer risk (Table 4). Among women who consumed a low level of these nutrients, dietary folate was weakly associated with a reduced risk of breast cancer. A striking inverse association was observed for dietary folate intake among those whose intake levels of methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub> were high. These data suggest an interaction, although tests for multiplicative interaction were not statistically significant. Additional analyses were conducted to evaluate folate intake and breast cancer risk, stratified by dietary sources of methionine and vitamin B<sub>6</sub>. The same pattern of associations was observed (data not shown).

## DISCUSSION

In this large population-based case-control study, we observed that dietary folate intake was inversely associated with the risk of breast cancer. This inverse association was particularly evident after adjust-

Table 3 Risk of breast cancer associated with dietary intake of folate, methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub>, Shanghai Breast Cancer Study 1996–1998

Dietary intake	Adjusted OR (95% CI) by quintile of intake <sup>a</sup>					<i>P</i> for trend
	Q1 (low)	Q2	Q3	Q4	Q5	
Folate						
Cases/controls	320/277	256/276	256/277	275/276	214/276	
OR (95% CI)	1.00 (reference)	0.83 (0.65–1.06)	0.87 (0.67–1.12)	0.93 (0.72–1.19)	0.71 (0.56–0.92)	0.05
OR (95% CI) <sup>b</sup>	1.00 (reference)	0.78 (0.61–1.01)	0.79 (0.61–1.03)	0.85 (0.65–1.11)	0.62 (0.46–0.82)	0.01
Vitamin B <sub>12</sub>						
Cases/controls	247/277	267/276	282/277	259/276	266/276	
OR (95% CI)	1.00 (reference)	1.21 (0.93–1.56)	1.29 (0.99–1.67)	1.18 (0.92–1.53)	1.18 (0.91–1.52)	0.33
OR (95% CI) <sup>b</sup>	1.00 (reference)	1.11 (0.85–1.44)	1.14 (0.87–1.49)	1.05 (0.80–1.36)	1.01 (0.77–1.32)	0.79
Methionine						
Cases/controls	234/277	245/276	234/277	304/276	298/276	
OR (95% CI)	1.00 (reference)	1.08 (0.84–1.40)	1.03 (0.80–1.34)	1.34 (1.04–1.73)	1.31 (1.02–1.68)	0.01
OR (95% CI) <sup>b</sup>	1.00 (reference)	0.95 (0.73–1.24)	0.84 (0.63–1.12)	1.00 (0.74–1.36)	0.79 (0.54–1.16)	0.47
Vitamin B <sub>6</sub>						
Cases/controls	219/277	268/276	268/277	270/276	296/276	
OR (95% CI)	1.00 (reference)	1.27 (0.98–1.64)	1.26 (0.97–1.62)	1.36 (1.05–1.76)	1.44 (1.12–1.85)	0.01
OR (95% CI) <sup>b</sup>	1.00 (reference)	1.26 (0.97–1.65)	1.22 (0.92–1.63)	1.35 (0.99–1.84)	1.46 (1.01–2.13)	0.07

<sup>a</sup> All ORs and 95% CI were calculated in a logistic regression model adjusted for total energy, age, education, family history of breast cancer, personal history of fibroadenoma, age at menarche, parity, age at first live birth, menopausal status, age at menopause, physical activity, and waist:hip ratio. Quintile cutpoints were based on residual energy-adjusted intake among controls.

<sup>b</sup> Additionally adjusted for total fruit and vegetable intake and total animal food intake.

Table 4 Risk of breast cancer associated with dietary folate intake by intake of methionine, vitamin B<sub>12</sub>, and vitamin B<sub>6</sub>, Shanghai Breast Cancer Study, 1996–1998

Dietary intake	Adjusted OR (95% CI) by quintile of folate intake <sup>a</sup>					P for trend
	Q1 (low)	Q2	Q3	Q4	Q5	
Methionine tertile						
T1 (low)	165/162 1.00 (ref)	80/105 0.75 (0.51–1.10)	55/75 0.75 (0.48–1.17)	69/64 1.10 (0.72–1.70)	42/55 0.83 (0.51–1.32)	0.81
T2	76/63 1.00 (ref)	86/107 0.66 (0.41–1.06)	103/115 0.73 (0.46–1.17)	90/110 0.71 (0.44–1.14)	46/65 0.57 (0.34–0.98)	
T3	79/52 1.00 (ref)	90/64 0.95 (0.58–1.55)	98/87 0.82 (0.50–1.32)	116/102 0.81 (0.51–1.30)	116/156 0.57 (0.37–0.89)	
P for interaction = 0.15						
Vitamin B <sub>6</sub> tertile						
T1 (low)	204/205 1.00 (ref)	98/129 0.77 (0.54–1.11)	55/71 0.81 (0.52–1.27)	27/37 0.71 (0.41–1.24)	15/19 0.81 (0.39–1.68)	0.19
T2	69/51 1.00 (ref)	113/108 0.77 (0.48–1.24)	125/123 0.77 (0.48–1.25)	113/117 0.73 (0.45–1.18)	34/61 0.43 (0.24–0.77)	
T3	47/21 1.00 (ref)	45/39 0.51 (0.25–1.04)	76/83 0.47 (0.25–0.89)	135/122 0.57 (0.31–1.85)	165/196 0.41 (0.23–0.73)	
P for interaction = 0.26						
Vitamin B <sub>12</sub> tertile						
T1 (low)	149/139 1.00 (ref)	75/104 0.69 (0.46–1.02)	64/73 0.85 (0.55–1.31)	78/72 1.04 (0.68–1.59)	59/73 0.78 (0.50–1.20)	0.69
T2	65/61 1.00 (ref)	94/94 0.97 (0.60–1.56)	113/98 1.18 (0.73–1.90)	106/119 0.88 (0.55–1.41)	77/88 0.83 (0.51–1.35)	
T3	106/77 1.00 (ref)	87/78 0.90 (0.57–1.42)	79/106 0.63 (0.41–0.99)	91/85 0.91 (0.58–1.44)	78/115 0.53 (0.35–0.82)	
P for interaction = 0.14						
Folate cofactor score						
3–4 (low)	135/141 1.00 (ref)	58/85 0.69 (0.44–1.07)	32/48 0.66 (0.38–1.15)	39/34 1.10 (0.63–1.91)	13/20 0.75 (0.35–1.62)	0.62
5–7	148/114 1.00 (ref)	145/156 0.74 (0.52–1.06)	158/151 0.93 (0.65–1.33)	154/163 0.83 (0.58–1.19)	92/118 0.62 (0.42–0.91)	
8–9	37/21 1.00 (ref)	50/34 0.85 (0.41–1.77)	64/77 0.50 (0.26–0.98)	84/78 0.65 (0.33–1.26)	108/137 0.47 (0.25–0.88)	
P for interaction = 0.49						

<sup>a</sup> All ORs are calculated from a logistic regression model and adjusted for total energy, age, education, family history of breast cancer, personal history of fibroadenoma, age at menarche, parity, age at first live birth, menopausal status, age at menopause, physical activity, and waist:hip ratio. Quintile cutpoints were based on residual energy-adjusted intake among controls.

ment for fruits and vegetables and among women whose usual diets contained a high level of methionine, vitamin B<sub>12</sub>, or vitamin B<sub>6</sub>. These findings are consistent with data from animal and *in vitro* studies implicating a potential role of folate in cancer prevention (33–38).

There are several possible mechanisms by which folate deficiency may contribute to the carcinogenic process:

(a) Folate deficiency may lead to decreased 5-methyl THF, which would result in decreased levels of SAM and global hypomethylation of DNA (39–41). Methylation is important for modulating gene expression, and hypomethylation has been observed to increase mutation rates via genomic instability (42–46).

(b) A second proposed mechanism is an increase in DNA replication errors. When folate levels are low, SAM formation may take precedence over thymidylate synthesis, resulting in a low thymidylate level and increasing misincorporation of uracil into DNA with resulting higher levels of chromosomal breaks (33, 39, 40, 47).

(c) Although somewhat contrary, folate deficiency also appears to be associated with hypermethylation of the CpG islands in the promoter region of several tumor suppressor genes or DNA repair genes, reducing their expression (40, 48). Hypermethylation of these genes is associated with the development of cancer (48–51).

Although folate intake has been investigated extensively in relation to colon cancer risk, only a few epidemiological studies have been conducted to evaluate its association with breast cancer risk. Our findings of an overall inverse association of folate intake with breast cancer risk are supported, in general, by three previous case-control studies (23, 24, 26). Of these, two were conducted in Western New York, United States, one among postmenopausal women and the other among premenopausal women, in which high folate intake was associated with a statistically nonsignificant 30% reduction in risk postmenopausally and a statistically significant 50% reduction in risk

premenopausally (23, 24). The third study, conducted in Uruguay, also found a statistically nonsignificant 30% reduction in risk (26). A 50% reduction in breast cancer risk associated with folate intake was observed among high consumers of alcohol in the NHS, although the relationship was not observed in the total sample (27). On the other hand, two previous studies reported no association between folate intake and breast cancer risk (22, 25). These two studies, however, may have suffered from recall bias or a small sample size.

In our study, the inverse association appeared to be particularly strong among those who consumed a high level of methionine, vitamin B<sub>12</sub>, or vitamin B<sub>6</sub>, even in the presence of no strong overall associations between these cofactors and breast cancer risk. This suggests that the cofactors may be important through their respective roles in folate metabolism; if the inverse association we observed is mostly attributable to folate, then it is expected that an inverse or no association between the cofactors and breast cancer risk would be observed as in this study. Potential modifying effects of vitamin B<sub>6</sub> or B<sub>12</sub> on the association between dietary folate and risk have not been evaluated in previous studies of breast cancer. Only two previous studies have examined the association of folate intake and breast cancer risk according to methionine intake level (27, 28). In a case-cohort study conducted in Canada, no modifying effect of methionine was observed (28). That study did not collect data on the folate content of multivitamins; therefore, intakes may have been underestimated for vitamin users. Results from the NHS cohort indicate no overall association between folate intake and breast cancer risk. An inverse association was observed among those who had a low methionine intake (27). However, the NHS differs from ours in that a large proportion of those participants were vitamin users, and the methionine levels in our population were lower than in the NHS. Specifically, the upper cutpoint for the lowest NHS quintile was close to the median intake level in our study. It is possible that we did not observe

a folate association among those with the lowest intake level because for these subjects folate cannot compensate for their methionine deficiency. It can also be speculated that because animal sources are substantial contributors to methionine intake in Western populations, some factor associated with animal food intake, such as red meat intake, may have masked or surpassed the folate association in the NHS.

Vitamin B<sub>12</sub> and vitamin B<sub>6</sub> are involved in the metabolism of folate, which may explain the potential modifying effect of these nutrients on the association of folate with breast cancer risk. Vitamin B<sub>12</sub> is a cofactor for the methyl transfer from 5-methyl THF to homocysteine to form methionine and THF. In conditions of low vitamin B<sub>12</sub> supply, the transfer of methyl groups may be reduced, thus affecting DNA methylation, synthesis, and repair. Vitamin B<sub>6</sub> is a coenzyme for homocysteine catabolism and for the acquisition of one-carbon units from serine to form 5,10-methylene THF, the folate that is used for DNA synthesis or for 5-methyl THF formation. In the presence of low vitamin B<sub>6</sub> supply, the conversion of THF to 5,10-methylene THF may be reduced which may, in turn, affect DNA synthesis or DNA methylation. Similarly, SAM inhibits methylenetetrahydrofolate reductase so that in the presence of adequate methionine, more folate may be available for use in thymidylate synthesis, reducing the probability of uracil misincorporation.

Because folate data are not available in the Chinese Food Composition Table, we used the USDA food composition data to estimate dietary intake values for Chinese women in Shanghai. Excellent match, however, was achieved between the mostly single foods listed in the food frequency questionnaire used in our study and those from the USDA database. Although it is possible that the absolute level of folate in foods listed in the database and those actually consumed in Shanghai differ because of different growing, shipping, processing, or other practices, this type of misclassification should be nondifferential between cases and controls, leading to attenuation of the true association in most situations. This is also true of any seasonal or other differences of foods within the urban Shanghai food supply. Because food biology and compositions are unlikely to be fundamentally different between foods produced in the United States and China and we are interested primarily in the relative intake of foods and nutrients, the estimates obtained in our study should be a relatively accurate reflection of folate intake in our study participants. Although we could not directly examine the correlation of folate intake using the USDA and Chinese databases, we did assess the correlation of three other water-soluble vitamins: vitamin C, riboflavin, and niacin. We found excellent correlation; all Pearson correlation coefficients were 0.91 or higher. These data provide additional assurance that the data on folate intake may be appropriate. As in any case-control study, the possibility of potential selection and recall biases cannot be dismissed completely. However, both cases and controls in this study had very high participation rates, thus decreasing the potential influence of selection bias on our results. Over 50% of cases were interviewed within 15 days after diagnosis, and over 80% were interviewed within 4 months of diagnosis, thus reducing potential measurement errors attributable to recent dietary changes after diagnosis. Furthermore, Chinese in Shanghai eat most of their food prepared using basic ingredients, and this behavior facilitates the quantification of usual diet over a longer period of time (5 years; Refs. 52–54) than the 12-month period assessed in typical studies conducted in Western countries. Confounding also may be a concern. However, we have collected extensive information on all known breast cancer risk factors and carefully adjusted for them in all data analyses. The strengths of our study include the very high participation rates and the ability to estimate folate intake in a population who were nonusers of

alcohol and vitamin supplements and who mainly consumed unprocessed and unfortified foods.

In summary, our study found evidence of a decreased risk of breast cancer associated with high consumption of folate among women who do not regularly consume alcohol. This relationship was especially apparent among subjects who also consumed higher amounts of methionine, vitamin B<sub>12</sub>, or vitamin B<sub>6</sub>. This study raises the possibility of important nutrient-nutrient interactions in breast carcinogenesis. As the body of literature on the role of polymorphic genes in folate metabolism increases, it is important for future studies to address the relative contributions of these nutrients and genes in breast carcinogenesis.

## ACKNOWLEDGMENTS

We thank Drs. Robert Bostick and Michael Wargovich for helpful comments and Zhi-Xian Ruan for her contribution in coordinating the field work.

## REFERENCES

- Graham, I. M., and O'Callaghan, P. The role of folic acid in the prevention of cardiovascular disease. *Curr. Opin. Lipidol.*, *11*: 577–587, 2000.
- Christensen, B., Arbour, L., Tran, P., Leclerc, D., Sabbaghian, N., Platt, R., Gilfix, B. M., Rosenblatt, D. S., Gravel, R. A., Forbes, P., and Rozen, R. Genetic polymorphisms in methylenetetrahydrofolate reductase and methionine synthase, folate levels in red blood cells, and risk of neural tube defects. *Am. J. Med. Genet.*, *84*: 151–157, 1999.
- Kim, Y. I. Folate and cancer prevention: a new medical application of folate beyond hyperhomocysteinemia and neural tube defects. *Nutr. Rev.*, *57*: 314–321, 1999.
- Lashner, B. A., Heidenreich, P. A., Su, G. L., Kane, S. V., and Hanauer, S. B. Effect of folate supplementation on the incidence of dysplasia and cancer in chronic ulcerative colitis. A case-control study. *Gastroenterology*, *97*: 255–259, 1989.
- Lashner, B. A., Provencher, K. S., Seidner, D. L., Knesebeck, A., and Brzezinski, A. The effect of folic acid supplementation on the risk for cancer or dysplasia in ulcerative colitis. *Gastroenterology*, *112*: 29–32, 1997.
- Benito, E., Stiggelbout, A., Bosch, F. X., Obrador, A., Kaldor, J., Mulet, M., and Munoz, N. Nutritional factors in colorectal cancer risk: a case-control study in Majorca. *Int. J. Cancer*, *49*: 161–167, 1991.
- Freudenheim, J. L., Graham, S., Marshall, J. R., Haughey, B. P., Cholewinski, S., and Wilkinson, G. Folate intake and carcinogenesis of the colon and rectum. *Int. J. Epidemiol.*, *20*: 368–374, 1991.
- Meyer, F., and White, E. Alcohol and nutrients in relation to colon cancer in middle-aged adults. *Am. J. Epidemiol.*, *138*: 225–236, 1993.
- Ferraroni, M., La Vecchia, C., D'Avanzo, B., Negri, E., Franceschi, S., and Decarli, A. Selected micronutrient intake and the risk of colorectal cancer. *Br. J. Cancer*, *70*: 1150–1155, 1994.
- Glynn, S. A., Albanes, D., Pietinen, P., Brown, C. C., Rautalahti, M., Tangrea, J. A., Gunter, E. W., Barrett, M. J., Virtamo, J., and Taylor, P. R. Colorectal cancer and folate status: a nested case-control study among male smokers. *Cancer Epidemiol. Biomark. Prev.*, *5*: 487–494, 1996.
- White, E., Shannon, J. S., and Patterson, R. E. Relationship between vitamin and calcium supplement use and colon cancer. *Cancer Epidemiol. Biomark. Prev.*, *6*: 769–774, 1997.
- Kerber, R. A., and Slattery, M. L. Comparison of self-reported and database-linked family history of cancer data in a case-control study. *Am. J. Epidemiol.*, *146*: 244–248, 1997.
- Giovannucci, E., Stampfer, M. J., Colditz, G. A., Hunter, D. J., Fuchs, C., Rosner, B. A., Speizer, F. E., and Willett, W. C. Multivitamin use, folate, and colon cancer in women in the Nurses' Health Study. *Ann. Intern. Med.*, *129*: 517–524, 1998.
- Giovannucci, E., Rimm, E. B., Ascherio, A., Stampfer, M. J., Colditz, G. A., and Willett, W. C. Alcohol, low-methionine-low-folate diets, and risk of colon cancer in men. *J. Natl. Cancer Inst.*, *87*: 265–273, 1995.
- Bird, C. L., Swendsen, M. E., Witte, J. S., Shikany, J. M., Hunt, I. F., Frank, H. D., Lee, E. R., Longnecker, M. P., and Haile, R. W. Red cell and plasma folate, folate consumption, and the risk of colorectal adenomatous polyps. *Cancer Epidemiol. Biomark. Prev.*, *4*: 709–714, 1995.
- Tseng, M., Murray, S. C., Kupper, L. L., and Sandler, R. S. Micronutrients and the risk of colorectal adenomas. *Am. J. Epidemiol.*, *144*: 1005–1014, 1996.
- Paspatis, G. A., Kalafatis, E., Oros, L., Xourgias, V., Koutsoumpa, P., and Karamanolis, D. G. Folate status and adenomatous colonic polyps. A colonoscopically controlled study. *Dis. Colon Rectum*, *38*: 64–67, 1995.
- Giovannucci, E., Stampfer, M. J., Colditz, G. A., Rimm, E. B., Trichopoulos, D., Rosner, B. A., Speizer, F. E., and Willett, W. C. Folate, methionine, and alcohol intake and risk of colorectal adenoma. *J. Natl. Cancer Inst.*, *85*: 875–884, 1993.
- Slattery, M. L., Potter, J. D., Samowitz, W., Schaffer, D., and Leppert, M. Methylenetetrahydrofolate reductase, diet, and risk of colon cancer. *Cancer Epidemiol. Biomark. Prev.*, *8*: 513–518, 1999.
- Ulrich, C. M., Kampman, E., Bigler, J., Schwartz, S. M., Chen, C., Bostick, R., Fosdick, Beresford, A. A., Yasui, Y., and Potter, J. D. Colorectal adenomas and the

- C677T MTHFR polymorphism: evidence for gene-environment interaction? *Cancer Epidemiol. Biomark. Prev.*, 8: 659–668, 1999.
21. Benito, E., Cabeza, E., Moreno, V., Obrador, A., and Bosch, F. X. Diet and colorectal adenomas: a case-control study in Majorca. *Int. J. Cancer*, 55: 213–219, 1993.
  22. Thorand, B., Kohlmeier, L., Simonsen, N., Croghan, C., and Thamm, M. Intake of fruits, vegetables, folic acid and related nutrients and risk of breast cancer in postmenopausal women. *Public Health Nutr.*, 1: 147–156, 1998.
  23. Graham, S., Hellmann, R., Marshall, J., Freudenheim, J., Vena, J., Swanson, M., Zielezny, M., Nemoto, T., Stubbe, N., and Raimondo, T. Nutritional epidemiology of postmenopausal breast cancer in western New York. *Am. J. Epidemiol.*, 134: 552–566, 1991.
  24. Freudenheim, J. L., Marshall, J. R., Vena, J. E., Laughlin, R., Brasure, J. R., Swanson, M. K., Nemoto, T., and Graham, S. Premenopausal breast cancer risk and intake of vegetables, fruits, and related nutrients. *J. Natl. Cancer Inst.*, 88: 340–348, 1996.
  25. Potischman, N., Swanson, C. A., Coates, R. J., Gammon, M. D., Brogan, D. R., Curtin, J., and Brinton, L. A. Intake of food groups and associated micronutrients in relation to risk of early-stage breast cancer. *Int. J. Cancer*, 82: 315–321, 1999.
  26. Ronco, A., De Stefani, E., Boffetta, P., Deneo-Pellegrini, H., Mendilaharsu, M., and Leborgne, F. Vegetables, fruits, and related nutrients and risk of breast cancer: a case-control study in Uruguay. *Nutr. Cancer*, 35: 111–119, 1999.
  27. Zhang, S., Hunter, D. J., Hankinson, S. E., Giovannucci, E. L., Rosner, B. A., Colditz, G. A., Speizer, F. E., and Willett, W. C. A prospective study of folate intake and the risk of breast cancer. *JAMA*, 281: 1632–1637, 1999.
  28. Rohan, T. E., Jain, M. G., Howe, G. R., and Miller, A. B. Dietary folate consumption and breast cancer risk. *J. Natl. Cancer Inst.*, 92: 266–269, 2000.
  29. Wu, K., Helzlsouer, K. J., Comstock, G. W., Hoffman, S. C., Nadeau, M. R., and Selhub, J. A prospective study on folate, B<sub>12</sub>, and pyridoxal 5'-phosphate (B6) and breast cancer. *Cancer Epidemiol. Biomark. Prev.*, 8: 209–217, 1999.
  30. Food and Drug Administration Food Standards. Amendment of standards of identity for enriched grain products to require addition of folic acid. *Fed. Regist.*, 61: 8781–8789, 1996.
  31. Gao, Y. T., Shu, X. O., Dai, Q., Potter, J. D., Brinton, L. A., Wen, W., Sellers, T. A., Kushi, L. H., Ruan, Z., Bostick, R. M., Jin, F., and Zheng, W. Association of menstrual and reproductive factors with breast cancer risk: results from the Shanghai Breast Cancer Study. *Int. J. Cancer*, 87: 295–300, 2000.
  32. Willett, W. C., and Stampfer, M. J. Total energy intake: implications for epidemiologic analyses. *Am. J. Epidemiol.*, 124: 17–27, 1986.
  33. Blount, B. C., Mack, M. M., Wehr, C. M., MacGregor, J. T., Hiatt, R. A., Wang, G., Wickramasinghe, S. N., Everson, R. B., and Ames, B. N. Folate deficiency causes uracil misincorporation into human DNA and chromosome breakage: implications for cancer and neuronal damage. *Proc. Natl. Acad. Sci. USA*, 94: 3290–3295, 1997.
  34. Duthie, S. J., and Hawdon, A. DNA instability (strand breakage, uracil misincorporation, and defective repair) is increased by folic acid depletion in human lymphocytes *in vitro*. *FASEB J.*, 12: 1491–1497, 1998.
  35. Duthie, S. J., Grant, G., and Narayanan, S. Increased uracil misincorporation in lymphocytes from folate-deficient rats. *Br. J. Cancer*, 83: 1532–1537, 2000.
  36. Kim, Y. I., Shirwadkar, S., Choi, S. W., Puchyr, M., Wang, Y., and Mason, J. B. Effects of dietary folate on DNA strand breaks within mutation-prone exons of the *p53* gene in rat colon. *Gastroenterology*, 119: 151–161, 2000.
  37. Cravo, M. L., Mason, J. B., Dayal, Y., Hutchinson, M., Smith, D., Selhub, J., and Rosenberg, I. Folate deficiency enhances the development of colonic neoplasia in dimethylhydrazine-treated rats. *Cancer Res.*, 52: 5002–5006, 1992.
  38. Kim, Y. I., Salomon, R. N., Graeme-Cook, F., Choi, S. W., Smith, D. E., Dallal, G. E., and Mason, J. B. Dietary folate protects against the development of macroscopic colonic neoplasia in a dose responsive manner in rats. *Gut*, 39: 732–740, 1996.
  39. Wainfan, E., and Poirier, L. A. Methyl groups in carcinogenesis: effects on DNA methylation and gene expression. *Cancer Res.*, 52: 2071s–2077s, 1992.
  40. Kim, Y. I. Folate and carcinogenesis: evidence, mechanisms, and implications. *J. Nutr. Biochem.*, 10: 66–88, 1999.
  41. Rampersaud, G. C., Kauwell, G. P., Hutson, A. D., Cerda, J. J., and Bailey, L. B. Genomic DNA methylation decreases in response to moderate folate depletion in elderly women. *Am. J. Clin. Nutr.*, 72: 998–1003, 2000.
  42. Wachsmann, J. T. DNA methylation and the association between genetic and epigenetic changes: relation to carcinogenesis. *Mutat. Res.*, 375: 1–8, 1997.
  43. Bird, A. P. The relationship of DNA methylation to cancer. *Cancer Surv.*, 28: 87–101, 1996.
  44. Baylin, S. B., Herman, J. G., Graff, J. R., Vertino, P. M., and Issa, J. P. Alterations in DNA methylation: a fundamental aspect of neoplasia. *Adv. Cancer Res.*, 72: 141–196, 1998.
  45. Gonzalgo, M. L., and Jones, P. A. Mutagenic and epigenetic effects of DNA methylation. *Mutat. Res.*, 386: 107–118, 1997.
  46. Cravo, M. L., Pinto, A. G., Chaves, P., Cruz, J. A., Lage, P., Nobre Leitao, C., and Costa, M. F. Effect of folate supplementation on DNA methylation of rectal mucosa in patients with colonic adenomas: correlation with nutrient intake. *Clin. Nutr.*, 17: 45–49, 1998.
  47. Kim, Y. I., Pogribny, I. P., Basnakian, A. G., Miller, J. W., Selhub, J., and Mason, J. B. Folate deficiency in rats induces DNA strand breaks and hypomethylation within the *p53* tumor suppressor gene. *Am. J. Clin. Nutr.*, 65: 46–52, 1997.
  48. Herman, J. G., Umar, A., Polyak, K., Graff, J. R., Ahuja, N., Issa, J. P., Markowitz, S., Willson, J. K., Hamilton, S. R., Kinzler, K. W., Kane, M. F., Kolodner, R. D., Vogelstein, B., Kunkel, T. A., and Baylin, S. B. Incidence and functional consequences of hMLH1 promoter hypermethylation in colorectal carcinoma. *Proc. Natl. Acad. Sci. USA*, 95: 6870–6875, 1998.
  49. Issa, J. P., Ottaviano, Y. L., Celano, P., Hamilton, S. R., Davidson, N. E., and Baylin, S. B. Methylation of the oestrogen receptor CpG island links ageing and neoplasia in human colon. *Nat. Genet.*, 7: 536–540, 1994.
  50. Makos, M., Nelkin, B. D., Lerman, M. I., Latif, F., Zbar, B., and Baylin, S. B. Distinct hypermethylation patterns occur at altered chromosome loci in human lung and colon cancer. *Proc. Natl. Acad. Sci. USA*, 89: 1929–1933, 1992.
  51. Kane, M. F., Loda, M., Gaida, G. M., Lipman, J., Mishra, R., Goldman, H., Jessup, J. M., and Kolodner, R. Methylation of the hMLH1 promoter correlates with lack of expression of hMLH1 in sporadic colon tumors and mismatch repair-defective human tumor cell lines. *Cancer Res.*, 57: 808–811, 1997.
  52. Dwyer, J. T., and Coleman, K. A. Insights into dietary recall from a longitudinal study: accuracy over four decades. *Am. J. Clin. Nutr.*, 65: 1153S–1158S, 1997.
  53. Smith, A. F. Cognitive Processes in Long-Term Dietary Recall. *Vital Health Statistics*, Vol. 6. Washington, DC: United States Government Printing Office, 1991.
  54. Jain, M., Howe, G. R., Harrison, L., and Miller, A. B. A study of repeatability of dietary data over a seven-year period. *Am. J. Epidemiol.*, 129: 422–429, 1989.

# Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

## Dietary Folate Intake and Breast Cancer Risk: Results from the Shanghai Breast Cancer Study

Martha J. Shrubsole, Fan Jin, Qi Dai, et al.

*Cancer Res* 2001;61:7136-7141.

**Updated version** Access the most recent version of this article at:  
<http://cancerres.aacrjournals.org/content/61/19/7136>

**Cited articles** This article cites 53 articles, 16 of which you can access for free at:  
<http://cancerres.aacrjournals.org/content/61/19/7136.full#ref-list-1>

**Citing articles** This article has been cited by 23 HighWire-hosted articles. Access the articles at:  
<http://cancerres.aacrjournals.org/content/61/19/7136.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://cancerres.aacrjournals.org/content/61/19/7136>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.