

Norepinephrine-induced Migration of SW 480 Colon Carcinoma Cells Is Inhibited by β -Blockers¹

Kai Masur,² Bernd Niggemann, Kurt S. Zanker, and Frank Entschladen

Institute for Immunology, Witten/Herdecke University, 58448 Witten, Germany

Abstract

β -Adrenoceptors are highly expressed on SW 480 colon carcinoma cells as was assessed by flow cytometry. We investigated the influence of norepinephrine on the migration of these cells using time-lapse videomicroscopy. Norepinephrine-treatment increased the locomotor activity within the population from 25% spontaneously locomoting cells to 65% locomoting cells. The β 1/2-blocker propranolol but not the β 1-blocker atenolol inhibited this increase. The intracellular signaling solely of norepinephrine-induced locomotion involved protein tyrosine kinase activity, whereas both spontaneous and norepinephrine-induced migration were reduced by inhibiting phospholipase C γ and protein kinase C α activity. In summary, norepinephrine-induced locomotion of SW 480 cells is β 2-adrenoceptor mediated and distinct from spontaneous locomotion concerning the PTK involvement.

Introduction

Chemokines are well described substances that initiate the locomotion of leukocytes (1, 2). These peptides act on the cells via seven helices or serpentine receptors (3). Besides chemokines, other ligands of serpentine receptors have been described that regulate the migration of leukocytes, e.g., formylated peptides (4). Recently, catecholamines have been described as being involved in the regulation of dendritic cell locomotion (5) and neutrophil granulocyte locomotion (6). Catecholamines act on the cells via adrenoceptors, which are members of the serpentine family. Serpentine receptors are on the intracellular side coupled to G protein-mediated signaling pathways (3) and to PTKs³ via β -arrestin (7). Catecholamines have been reported to enhance the carcinogenic effect of tobacco-specific nitrosamine as was measured by the development of pulmonary adenocarcinoma in hamsters (8). Therefore, catecholamines have effects on both tumor cells and leukocytes. In our study we investigated whether catecholamines are involved in the regulation of tumor cell migration.

Materials and Methods

Cell Culture. The colon carcinoma cell line SW 480 (American Type Culture Collection, Rockville, MD) was maintained in antibiotic-free Leibovitz L-15 medium (PAA Laboratories GmbH, Linz, Austria), supplemented with 10% FCS (PAA Laboratories GmbH) in a humidified atmosphere without CO₂ addition.

Cell Migration Assay. Cultured cells were harvested using a trypsin/EDTA solution. Six $\times 10^4$ cells were mixed with 150 μ l of buffered liquid collagen [1.63 mg/ml collagen type I (pH 7.4); Collagen Corporation, Fremont, CA) containing MEM (Sigma, Deisenhofen, Germany) as well as norepineph-

rine and the investigated pharmacological substances. Self-constructed glass chambers (9) were filled with this mixture. After polymerization of the collagen, the chambers were sealed, and cell locomotion within the three-dimensional collagen lattices was recorded by time-lapse videomicroscopy overnight at 37°C. For the analysis of migratory activity, 30 cells of each sample were randomly selected and two-dimensional projections of paths were calculated by computer-assisted cell tracking in 20-min step intervals.

Only the investigated substances were added to the collagen lattices; the cells were not incubated with any of the substances prior to the mixing of the cells with these substances and the buffered collagen. Norepinephrine was used at 1, 10, and 100 μ M; either propranolol or atenolol was added in equimolar concentrations to norepinephrine. The src-specific PTK (3) inhibitor PP2 (10), the PLC γ -specific inhibitor U73122 (11), and the PKC inhibitor Go6976, specific for the α isotype (12), were added alone or in combination with 10 μ M norepinephrine. All of these pharmacological substances were provided by Calbiochem-Novabiochem GmbH, Bad Soden, Germany. None of the substances was used in a concentration that was cytotoxic as was assessed by flow cytometry.

Flow Cytometry. The expression of α - and β -adrenoceptors of the SW 480 cells was determined using a FACS Calibur flow cytometer (Becton Dickinson, Heidelberg, Germany). Generally, 1×10^5 cells were incubated with 10 μ g/ml primary antibody for 10 min at room temperature. The antibodies directed against the β 1-, β 2-, α 2_B-, and α 2_C-adrenoceptors were provided by Santa Cruz Biotechnology, Santa Cruz, CA; the antibodies directed against the α 1- and α 2_A-adrenoceptors were provided by Dianova, Hamburg, Germany. After the incubation, the cells were washed and incubated with a FITC-conjugated secondary antibody (Fab fragment goat antimouse or antirabbit; Jackson ImmunoResearch Laboratories, West Grove, PA) under the same conditions. Nonspecific binding was determined by an unlabeled isotypic control antibody (Coulter-Immunotech, Hamburg, Germany). Additionally, flow cytometry was used to determine the cell viability subsequent to the migration experiments. The collagen matrices were digested using collagenase type I and IV (Worthington Biochemical Corp., Freehold, NJ), the cells were harvested and subjected to flow cytometry after propidium iodide staining. No changes of the cell viability attributable to treatment with the above mentioned inhibitors were observed throughout the experiments.

Results

Expression of Adrenoceptors. Flow cytometric analyses of the expression of α - and β -adrenoceptors subtypes revealed a faint expression of the α 1- and α 2_A-adrenoceptors expression with a mean FITC-fluorescence intensity of 18.8 and 18.4, respectively, as compared with the isotypic control (14.8; Fig. 1). The α 2_B- and α 2_C-adrenoceptors mean FITC-fluorescence intensity was slightly higher (21.3 and 30.9, respectively), and the mean FITC-fluorescence intensity of the isotypic control was 5.8. In contrast to this low expression of α -adrenoceptors, a very high expression of β -adrenoceptors was observed. The mean FITC-fluorescence intensity was 315.7 for the β 1-adrenoceptor and 325.7 for the β 2-adrenoceptor, compared with 14.8 for the isotypic control.

Norepinephrine-induced Cell Migration. After incorporation within a three-dimensional collagen matrix, the SW 480 colon carcinoma cells developed a spontaneous locomotor activity of 25% locomoting cells (Fig. 2). This spontaneous locomotor activity of the SW 480 cells has been described previously by Kubens and Zanker (13).

Received 1/25/01; accepted 2/12/01.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported by the Deutsche Krebshilfe, Bonn, Germany, and the Fritz-Bender-Foundation, Munich, Germany.

² To whom requests for reprints should be addressed, at Institute for Immunology, Witten/Herdecke University, Stockumer Str. 10, 58448 Witten, Germany. E-mail: kaimasur@uni-wh.de.

³ The abbreviations used are: PTK, protein tyrosine kinase; PKC, protein kinase C; PLC, phospholipase C.




Fig. 1. Analyses of the α - and β -adrenoceptor expression of SW 480 colon carcinoma cells. The antibody bound FITC-fluorescence of specific antibodies (filled gray) was compared with an unspecific isotypic control (black line).

Treatment of the SW 480 cells with 10 μ M norepinephrine augmented the locomotor activity to 65%. Treatment of the cells with 1 μ M norepinephrine increased the locomotor activity to a lesser extent, whereas 100 μ M norepinephrine led to results similar to those with 10 μ M norepinephrine (not shown). The addition of the β -adrenoceptor blocker propranolol had no effect on the spontaneous locomotor activity (Fig. 2A), but reduced the norepinephrine-induced locomotion to the activity of the untreated control. In contrast, the β 1-specific adrenoceptor blocker atenolol only marginally influenced the norepinephrine-induced locomotion (Fig. 2B).

Signaling Cascade of Norepinephrine-induced Migration. In search of the signal transduction underlying the spontaneous and norepinephrine-induced locomotion of the SW 480 cells, we inhibited key regulatory molecules using specific inhibitors. The activity of PTKs of the src-family was inhibited using PP2 (Fig. 3A). This inhibitor led to a reduction of the norepinephrine-induced locomotion from 70% locomoting cells to the level of spontaneous locomotor activity (20% locomoting cells) within the course of the experiment. The spontaneous locomotion was not reduced by this inhibitor. In contrast, inhibition of the PLC γ using the specific inhibitor U73122 (Fig. 3B) and inhibition of the PKC α using the specific inhibitor Go6976 (Fig. 3C) led to a reduction of both spontaneous and norepinephrine-induced locomotion as well.

Discussion

We have reported previously that the intracellular signaling that regulates the migration of neutrophil granulocytes and T lymphocytes differs between spontaneous locomotion and the migration induced by chemokines (14). Herein we used norepinephrine to investigate

whether the engagement of serpentine receptors might initiate a type of tumor cell migration that differs from the spontaneous migration. Spontaneous locomotor activity is supposed to be initiated by integrin binding to the extracellular matrix (2). Using MV3 melanoma cells, we showed that the blocking of the collagen-binding $\alpha_2\beta_1$ integrin inhibited spontaneous locomotor activity (15). Our investigation proved that norepinephrine induces migration in addition to the spontaneous migration of the SW 480 colon carcinoma cells. This migratory type is different from the spontaneous locomotion with regard to the involvement of src-PTKs (Fig. 4). The functional link of the adrenoceptors to a PTK signaling is given by β -arrestin (7). Downstream from the signal transduction, the PLC γ is activated by tyrosine phosphorylation caused by PTKs (16). The activity of this phospholipase is an integrating signal for the regulatory signal transduction of both spontaneous (*i.e.*, matrix-induced) and norepinephrine-induced locomotion, because both types of migration are reduced by an inhibitor specific for this enzyme (Fig. 4). The enzymatic activity of the PLC γ generates the second messenger diacylglycerol, which is an activator for the PKC α . We have shown previously that the PKC α is the isozyme needed for colon carcinoma cells migration (17). The PKC α plays an important role even in untransformed, normal colon epithelium. Frey *et al.* (18) have shown, that the PKC α in nontransformed intestinal epithelial cells regulates the growth via modulation of Cip/Kip family cyclin-dependent kinase inhibitors and the retinoblastoma suppressor protein. Furthermore, the PKC α is located in focal adhesions of rat embryo fibroblasts (19). Focal adhesions are




Fig. 2. Inhibition of the promigratory catecholamine effect in SW 480 cells by propranolol but not by atenolol. Migration of the cells was induced by the addition of 10 μ M norepinephrine. β -adrenoceptors were blocked by the addition of either 10 μ M propranolol (A) or 10 μ M atenolol (B). The graphs show mean values of three independent experiments (90 cells were analyzed per sample).

multi-protein complexes that are essential for the migration of tumor cells (2).

Norepinephrine is a neurotransmitter that is also released in stress reactions. The long-lasting elevation of catecholamines attributable to chronic stress is known to be a risk factor for heart as well as for cancer diseases (20). Thus, we provide molecular evidence for a functional link between psychoneurological events namely the increasing release of the neurotransmitter norepinephrine *in vivo* and its promigratory influence on tumor cells *in vitro*. This finding gets even more relevant when taking the immune system into consideration: we have shown that two important effector cells of the immune system,




Fig. 3. Signal transduction of the catecholamine-induced SW 480 cell migration. Migration of the cells was induced by addition of 10 μ M norepinephrine. The activity of src-PTKs was inhibited by 10 nM PP2 (A), the PLC γ was inhibited by 4 μ M U73122 (B), and the PKC α was inhibited by 6 nM Go6976 (C). The graphs show mean values of three independent experiments (90 cells were analyzed per sample).




Fig. 4. Model of the signal transduction pathway regulating spontaneous (matrix-induced) and norepinephrine-induced locomotion of SW 480 colon carcinoma cells.

the T lymphocytes and the neutrophil granulocytes, are inhibited in their migratory activity by the presence of epinephrine or norepinephrine.⁴ The inhibiting effect of catecholamines on the formyl-methionyl-leucyl-phenylalanine-induced neutrophil granulocyte migration is caused by an increase of cellular cAMP as was reviewed by Elferink and VanUffelen (6).

The promigratory effect of norepinephrine on the migration of SW480 colon carcinoma cells was inhibited by the β -adrenoceptor-blocking agent propranolol at pharmacological dosages relevant for human beings. More interestingly, atenolol, a specific β 1-adrenoceptor-blocking agent, did only marginally influence migration. This might suggest the use of β 2-blocking, non-heart active pharmaceuticals for the preventive treatment in a diagnosed colon carcinoma to inhibit metastatogenesis in the progress of the cancer disease by following preventive clinical trials. Furthermore, ligands of serpentine in the immune system (*i.e.*, chemokines) not only initiate migration but also cause directed migration within a gradient (14). If a gradient of catecholamines could similarly cause a directed migration of colon carcinoma cells, this would have considerable consequences for the view on the pattern of metastases occurring in this special type of cancer. Therefore, our findings might open new pharmacological possibilities for the preventive treatment of a colon carcinoma, to delay or to inhibit the progression of the disease with regard to invasion and the development of metastases.

References

- Luster, A. D. Chemokines—chemotactic cytokines that mediate inflammation. *N. Engl. J. Med.*, 338: 436–445, 1998.
- Entschladen, F., and Zanker, K. S. Locomotion of tumor cells: a molecular comparison to migrating pre- and postmitotic leukocytes. *J. Cancer Res. Clin. Oncol.*, 126: 671–681, 2000.

⁴ Unpublished observations.

3. Howard, Z. O. M., Ben-Baruch, A., and Oppenheim, J. J. Chemokines: progress toward identifying molecular targets for therapeutic agents. *Trends Biotechnol.*, *14*: 46–51, 1996.
4. Thivierge, M., Parent, J. L., Stankova, J., and Rola-Pleszczynski, M. Modulation of formyl peptide receptor expression by IL-10 in human monocytes and neutrophils. *J. Immunol.*, *162*: 3590–3595, 1999.
5. Maestroni, G. J. M. Dendritic cell migration controlled by $\alpha 1\beta$ -adrenergic receptors. *J. Immunol.*, *165*: 6743–6747, 2000.
6. Elferink, J. G. R., and VanUffelen, B. E. The role of cyclic nucleotides in neutrophil migration. *Gen. Pharmacol.*, *27*: 387–393, 1996.
7. Luttrell, L. M., Ferguson, S. S. G., Daaka, Y., Miller, W. E., Maudsley, S., Della Rocca, G. J., Lin, F. T., Kawakatsu, H., Owada, K., Luttrell, D. K., Caron, M. G., and Lefkowitz, R. J. β -Arrestin-dependent formation of $\beta 2$ adrenergic receptor-src protein kinase complexes. *Science (Washington DC)*, *283*: 655–661, 1999.
8. Schuller, H. M., Porter, B., and Riechert, B. β -Adrenergic modulation of NNK-induced lung carcinogenesis in hamsters. *J. Cancer Res. Clin. Oncol.*, *126*: 624–630, 2000.
9. Friedl, P., Maaser, K., Klein, E. C., Niggemann, B., Krohme, G., and Zanker, K. S. Migration of highly aggressive MV3 melanoma cells in 3-dimensional collagen lattices results in local matrix reorganization and shedding of $\alpha 2$ and $\beta 1$ integrins and CD44. *Cancer Res.*, *57*: 2061–2070, 1997.
10. Salazar, E. P., and Rozengurt, E. Bombesin and platelet-derived growth factor induce association of endogenous focal adhesion kinase with src in intact Swiss 3T3 cells. *J. Biol. Chem.*, *274*: 28371–28378, 1999.
11. Stam, J. C., Michiels, F., van der Kammen, R. A., Moolenaar, W. H., and Collard, J. G. Invasion of T-lymphoma cells: cooperation between Rho family GTPases and lysophospholipid receptor signaling. *EMBO J.*, *17*: 4066–4074, 1998.
12. Martiny-Baron, G., Kazanietz, M. G., Mischak, H., Blumberg, P. M., Kochs, G., Hug, H., Marme, D., and Schachtele, C. Selective inhibition of protein kinase C isozymes by the indolocarbazole Go 6976. *J. Biol. Chem.*, *268*: 9194–9197, 1993.
13. Kubens, B. S., and Zanker, K. S. Differences in the migration capacity of primary human colon carcinoma cells (SW 480) and their lymph node metastatic derivatives (SW 620). *Cancer Lett.*, *131*: 55–64, 1998.
14. Entschladen, F., Gunzer, M., Scheuffele, C. M., Niggemann, B., and Zanker, K. S. T lymphocytes and neutrophil granulocytes differ in regulatory signaling and migratory dynamics with regard to spontaneous locomotion and chemotaxis. *Cell Immunol.*, *199*: 104–114, 2000.
15. Maaser, K., Wolf, K., Klein, C. E., Niggemann, B., Zanker, K. S., Brocker, E. B., and Friedl, P. Functional hierarchy of simultaneously expressed adhesion receptors: integrin $\alpha 2\beta 1$ but not CD44 mediates MV3 melanoma cell migration and matrix reorganization within three-dimensional hyaluronan-containing collagen matrices. *Mol. Biol. Cell* *10*: 3067–3079, 1999.
16. Cook, S. J., and Wakelam, M. J. O. Phospholipases C and D in mitogenic signal transduction. *Rev. Physiol. Biochem. Pharmacol.*, *119*: 13–45, 1992.
17. Masur, K., Niggemann, B., Zanker, K. S., and Entschladen, F. PKC α is a basic mediator of colon carcinoma cell migration. *Signal Transduction*, *1*: 81, 2001.
18. Frey, M. R., Saxon, M. L., Zhao, X., Rollins, A., Evans, S. S., and Black, J. D. Protein kinase C isozyme-mediated cell cycle arrest involves induction of p21(waf1/cip1) and p27(kip1) and hypophosphorylation of the retinoblastoma protein in intestinal epithelial cells. *J. Biol. Chem.*, *272*: 9424–9435, 1997.
19. Jaken, S., Leach, K., and Klauck, T. Association of type 3 protein kinase C with focal contacts in rat embryo fibroblasts. *J. Cell. Biol.*, *109*: 697–704, 1998.
20. Paredes, A., Galvez, A., Leyton, V., Aravena, G., Fiedler, J. L., Bustamante, D., and Lara, H. E. Stress promotes development of ovarian cysts in rats: the possible role of sympathetic nerve activation. *Endocrine*, *8*: 309–315, 1998.

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

Norepinephrine-induced Migration of SW 480 Colon Carcinoma Cells Is Inhibited by β -Blockers

Kai Masur, Bernd Niggemann, Kurt S. Zanker, et al.

Cancer Res 2001;61:2866-2869.

Updated version Access the most recent version of this article at:
<http://cancerres.aacrjournals.org/content/61/7/2866>

Cited articles This article cites 20 articles, 10 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/61/7/2866.full#ref-list-1>

Citing articles This article has been cited by 21 HighWire-hosted articles. Access the articles at:
<http://cancerres.aacrjournals.org/content/61/7/2866.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerres.aacrjournals.org/content/61/7/2866>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.