

Breast Cancer Cells Can Evade Apoptosis-Mediated Selective Killing by a Novel Small Molecule Inhibitor of Bcl-2

Pedro Jose Real,¹ Yeyu Cao,³ Renxiao Wang,³ Zaneta Nikolovska-Coleska,³ Jaime Sanz-Ortiz,² Shaomeng Wang,³ and Jose Luis Fernandez-Luna¹

¹Unidad de Genética Molecular and ²Servicio de Oncología Médica, Hospital Universitario Marques de Valdecilla, Servicio Cantabro de Salud, Santander, Spain; and ³Departments of Internal Medicine and Medicinal Chemistry, and Comprehensive Cancer Center, University of Michigan, Ann Arbor, Michigan

ABSTRACT

Proteins of the Bcl-2 family are key regulators of caspase activation and apoptosis. Some members of this family, notably Bcl-2 and Bcl-x_L, are overexpressed in cancer cells, which have been associated with chemoresistance. We have designed and synthesized a small molecule inhibitor of Bcl-2, named YC137, and studied its role in cancer cells. *In vitro* studies showed that YC137 inhibits the binding of the Bid BH3 peptide to Bcl-2, thus disrupting an interaction essential for the antiapoptotic activity of Bcl-2. This inhibitor induces apoptosis of hematopoietic progenitors overexpressing Bcl-2 but not Bcl-x_L and breast cancer cells that express high levels of Bcl-2. On the contrary, a variety of normal primary cells, including CD34⁺ progenitors, myoblasts, and peripheral blood mononuclear cells, do not respond to the inhibitor. A breast cancer cell line resistant to YC137 was generated. Analysis of resistant cells revealed a reduced expression of Bcl-2, which correlated with low activation of signal transducer and activator of transcription-3 (Stat3) and reduced expression of the human epidermal growth factor receptor-2 (HER2). Of note, YC137-resistant cells were more sensitive to apoptosis induced by chemotherapy. Because HER2 has not been linked previously to the Stat3-Bcl-2 transcriptional pathway, we additionally confirmed that specific blockade of HER2 in breast cancer cells resulted in down-regulation of Stat3 activity and reduced levels of Bcl-2. Consistently, HER2 blockade led to YC137 resistance. These data provide evidence for the selective killing of tumor cells by YC137 and represent the first example of *in vitro* selection of cancer cells refractory to a Bcl-2 inhibitor.

INTRODUCTION

The expression of genes that regulate apoptotic cell death plays an important role in determining the sensitivity of tumor cells to chemotherapy. High expression of the antiapoptotic protein Bcl-2 is found in a number of human hematologic malignancies and solid tumors (1, 2). The functional blockade of Bcl-2 or other antiapoptotic proteins, such as Bcl-x_L, could either induce apoptosis in cancer cells or sensitize these cells for chemotherapy. To this end, it has been reported that down-regulation of Bcl-2 by antisense oligonucleotides induces apoptosis in myeloid leukemia cells, breast and colorectal carcinomas, and lung cancer cells among others, even in the presence of other antiapoptotic genes (3–6). In other cell systems (*i.e.*, MCF7 breast cancer cells, HL-60 myeloid leukemia cell line, and myeloma cells), the use of a Bcl-2 antisense oligonucleotide sensitizes tumor cells to chemotherapeutic drugs (3, 7, 8). Consequently, antiapoptotic members of the Bcl-2 family have attracted intensive interest in drug

discovery to develop a new class of anticancer agents (9). BH3-mediated homodimerizations and heterodimerizations play a key role in regulating the apoptotic activity of Bcl-2 family members (10). Structural analysis of the Bcl-x_L-Bak BH3 peptide complex (11) and more recently the Bcl-2 protein (12) has made possible the identification of small molecules that inhibit the interaction between the BH3 domain of proapoptotic proteins and the hydrophobic cleft of Bcl-2 or Bcl-x_L (13–16). Some of these compounds have been shown to reduce cell viability in cancer cell lines, which suggests that they could lead to the development of new therapeutic agents. However, it has not elucidated the activity of these small molecules on normal cells, such as hematopoietic progenitors and epithelial cells, which are commonly affected after chemotherapy.

The clinical success of small molecule kinase inhibitors has led to the notion that other selective inhibitors may have significant activity against a range of human malignancies (17). However, clinical resistance to these agents has been reported, and it is believed that preexisting mutant cells are selected by the agent to which the mutation confers resistance (18). By analogy, Bcl-2 inhibitors could select for resistant cells. Consequently, research should focus on defining the mechanisms of resistance, so that the emergence of resistant clones might be prevented.

In this study, we show that a novel small molecule inhibitor of Bcl-2, YC137, induces apoptosis in breast cancer cells that express high levels of Bcl-2 but does not have any effect on normal hematopoietic progenitors, peripheral blood mononuclear cells (PBMCs), small intestine epithelial cells, and myoblasts. Furthermore, tumor cells that evade the apoptotic effects of YC137 down-regulate Bcl-2 and become more sensitive to conventional chemotherapy.

MATERIALS AND METHODS

Cell Lines and Primary Cells. Breast cancer cell lines MDA-MB435L (lung metastatic variant of MDA-MB435), MDA-MB435B (brain metastatic variant of MDA-MB435), MDA-MB231, MCF7, and SKBR3 were cultured as described previously (19). SUM185, SUM159, and SUM229 were incubated in DMEM/Ham's F-12 (Biochrom KG, Berlin, Germany), supplemented with 5% fetal calf serum, 5 μg/mL insulin, and 1 μg/mL hydrocortisone (both from Sigma, St. Louis, MO). Normal small intestine FHs cells were grown in DMEM/Ham's F-12 with 10% fetal calf serum, 10 ng/mL cholera toxin, 5 μg/mL transferrin, 5 μg/mL insulin, and 100 ng/mL hydrocortisone (Sigma). HCD-57 and FL5.12 hematopoietic cells and their derivative cell lines transfected with Bcl-2 or Bcl-x_L were cultured as described previously (20).

PBMCs were obtained from normal donors. CD34⁺ cells were selected from the PBMC population of donors undergoing mobilization for allogeneic progenitor cell transplantation as described previously (21). All of the donors signed informed consent according to Guidelines from the Committee for the Protection of Human Subjects at the University Hospital Marques de Valdecilla. Myoblasts were obtained from the Cell Therapy Unit, at the University Clinic of Navarra following a protocol described previously (22). In some experiments, MB435B cells were treated with the recombinant humanized antihuman epidermal growth factor receptor (HER)2 monoclonal antibody trastuzumab (F. Hoffmann-La Roche, Basel, Switzerland) and then additionally analyzed for the expression of Bcl-2.

Chemical Synthesis of YC137. Briefly, 1.35 g (5.95 mmol) of 2,3-dichloro-1,4-naphthoquinone were dissolved in 20 mL of 1,4-dioxane, and

Received 3/17/04; revised 7/9/04; accepted 8/25/04.

Grant support: Fondo de Investigaciones Sanitarias Grants No. PI020091 and C03/10 (Red Temática de Investigación Cooperativa de Centros de Cáncer), Fundación Marcelino Botín (J. L. Fernandez-Luna), and the United States Department of Defense Breast Cancer Program (DAMD17-01-1-0256; S. Wang). P. J. Real is a recipient of a predoctoral fellowship from the Fundación Ramon Areces.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Requests for reprints: Jose Luis Fernandez-Luna, Unidad de Genética Molecular, Hospital Universitario Marques de Valdecilla, Servicio Cantabro de Salud, Edificio Escuela Universitaria de Enfermería, Av. Valdecilla s/n, 39008 Santander, Spain. Phone: 34-942-200952; Fax: 34-942-200952; E-mail: fluna@humv.es.

©2004 American Association for Cancer Research.

then 1 g (5.95 mmol) of 4-hydroxyphenylthiourea was added. The mixture was refluxed for 20 minutes. When the color of the reaction mixture turned to deep purple, 20 mL of EtOH and 1.344 g (12 mmol) of 1,4-diazabicyclo(2,2,2)octane (Dabco) were added and refluxed for 5 hours. The reaction mixture was then cooled down to room temperature, and the solvent was removed under vacuum. Then, the important intermediate (deep violet solid) was purified by flash column chromatography. To a solution of 10 mL of dichloromethane and 110 mg (0.53 mmol) of *N*-methyl-formate-*L*-methionine, 110 mg (0.53 mmol) of dicyclohexylcarbodiimide were added in ice-cold bath, followed by the addition of 142 mg (0.44 mmol) of the intermediate. Then, the reaction mixture was allowed to warm up to room temperature and stirred for 16 hours. The solid of the reaction mixture was filtered out and washed with brine, dried over sodium sulfate, and purified via flash chromatography.

YC137 was solubilized at 10 mmol/L in dimethylsulfoxide and kept at -20°C in the dark until use. This stock solution was diluted $>10^4$ -fold for both binding assays and cell culture experiments.

Competitive Fluorescence Polarization Binding Assay. A sensitive and quantitative *in vitro* binding assay using a fluorescence polarization-based method was developed and used to determine the binding affinity of YC137 to Bcl-2 protein. For this assay, 6-carboxy fluorescein succinimidyl ester was coupled to a 21-residue Bid peptide (Flu-QEDIIRNIARHLAQVGDSDMR), which binds with high affinity to Bcl-2 protein ($K_d = 47$ nmol/L).

Fluorescence polarization experiments were performed in Dynex 96-well, black, and round-bottomed plates (Fisher Scientific, Hampton, NH) using the Ultra Plate Reader (Tecan U.S., Inc., Research Triangle Park, NC). YC137, diluted in assay buffer, was incubated with recombinant His-fused soluble Bcl-2 protein (120 nmol/L) and Flu-Bid peptide (5 nmol/L). It is worthy of note that the assay buffer conditions have a great influence on the binding affinity of YC137. Thus, we tried a number of assay conditions until consistent results were obtained in three independent experiments with a buffer containing 50 mmol/L Bis-Tris (pH 7.4) and 0.01% bovine γ globulin. The plates were mixed and incubated at room temperature for 4 hours to reach equilibrium. Curve fitting was performed using GraphPad Prism software. A nonlabeled Bid BH3 peptide was used as the positive control. The K_i values were calculated using a recently developed equation for fluorescence polarization-based binding assays (23). For determination of the binding affinity to Bcl- x_L protein, we used human Bcl- x_L recombinant His-tagged protein without the COOH-terminal hydrophobic tail (120 nmol/L) and Flu-Bid peptide (5 nmol/L), and the competitive binding assay was performed in the same way as for the Bcl-2 protein.

Transfection and Flow Cytometry Analysis. CD34⁺ progenitors (1×10^6 cells) were cotransfected with 1 μg of a vector encoding enhanced green fluorescence protein (pEGFP-C1, BD Biosciences, Palo Alto, CA) and 4 μg of pSFFV-Bcl-2 (24) by using the human CD34 cell nucleofactor kit (Amaxa, Cologne, Germany). After nucleofection, cells were cultured for 16 h with or without stem cell factor and interleukin (IL)-3 at a final concentration of 100 ng/mL. Then, 0.5 $\mu\text{mol/L}$ YC137 was added to the culture, and after 24 hours of incubation, apoptotic cells were detected, within the EGFP⁺ population, with annexin V labeled with phycoerythrin (Becton Dickinson, San Jose, CA) by flow cytometry using a FACScan analyzer (Becton Dickinson).

MB435B cells were cotransfected with 1 μg of pEGFP-C1 and 4 μg of either a constitutively active form of Stat3 (RCCMV-Stat3-C; ref. 25) or the empty vector (RCCMV) as described previously (19) and cultured with or without 50 $\mu\text{g/mL}$ trastuzumab. After 36 hours of transfection, EGFP-positive cells were analyzed for the expression of Bcl-2 by flow cytometry, as described (24).

Analysis of Apoptosis. Apoptosis was determined by a quantitative analysis of histone-associated DNA fragments using an ELISA kit (Roche, Mannheim, Germany) and additionally confirmed by DNA content using a FACScan, which correlates the percentage of hypodiploid cells with the extent of apoptosis. When indicated, activation of caspase 9 was determined in cells cultured with YC137 by using FITC-labeled LEHD-FMK peptide (Oncogene Research Products, San Diego, CA), a cell-permeable fluorescent inhibitor of activated caspase 9. After treatment, fluorescent cells were detected by flow cytometry.

In some experiments, 10^5 hematopoietic progenitors were cocultured with the same number of MB435B cells in the presence of YC137, and apoptotic cell death was detected in the CD34⁺ and CD34⁻ cell populations with

phycoerythrin-labeled annexin V and FITC-labeled anti-CD34 antibody (Becton Dickinson) by flow cytometry.

Cytochrome *c* release was analyzed on soluble and heavy membrane fractions, obtained as described (26), by Western blot with specific antibodies against cytochrome *c*.

Electrophoretic Mobility Shift Assay. Nuclear extracts (5 to 10 μg of total protein) were incubated with a ^{32}P -labeled, double-stranded DNA probe containing a consensus sequence for Stat3 as described previously (19). Samples were run on a 5% nondenaturing polyacrylamide gel in 200 mmol/L Tris-borate and 2 mmol/L EDTA. Gels were dried and visualized by autoradiography. Supershifts were performed using rabbit polyclonal antibodies specific for Stat3 (Santa Cruz Biotechnology, Santa Cruz, CA).

Immunoblotting. The expression of Bcl-2, Bcl- x_L , and HER2 proteins were determined by Western blotting as described previously (19). Briefly, proteins (30 to 60 μg) were separated on a 12% polyacrylamide gel and transferred to nitrocellulose. Blots were blocked with 3% bovine serum albumin and incubated with rabbit anti-Bcl- x_L , or mouse anti-Bcl-2, anti-cytochrome *c* (all from Becton Dickinson), anti-HER2 (Ab-3; Oncogene Research Products), and anti- β -tubulin antibodies (Sigma) and then incubated with goat antirabbit or antimouse antibodies conjugated to alkaline phosphatase (Sigma). Bound antibody was detected by a chemiluminescence system (Applied-Biosystems, Foster City, CA).

Reverse Transcription-PCR Analysis. Total RNA was prepared using TRIZOL reagent (Invitrogen, Carlsbad, CA). To assess mRNA expression of Bcl-2, a quantitative real-time PCR was performed in a 7000 Sequence Detection System (Applied-Biosystems) using primers $5'$ catgtgtgtggagagcgtcaaa $3'$ and $5'$ gccggttcaggtactcagca $3'$. The ratio of the abundance of bcl-2 transcripts to that of glyceraldehyde-3-phosphate dehydrogenase transcripts was calculated as described (27). Semiquantitative reverse transcription-PCR conditions and specific primers for HER2, epidermal growth factor receptor (EGFR), and glyceraldehyde-3-phosphate dehydrogenase have been described previously (19).

RESULTS

Characterization of a Small Molecule Inhibitor of Bcl-2. We have used a structure-based computer screening strategy to discover novel small molecule inhibitors that bind to the crucial BH3-binding groove of Bcl-2 protein (16). On the basis of the structure of initial lead compounds, we have designed and synthesized new analogues to improve their binding affinity and cellular activity. One of these compounds, YC137 (Fig. 1A), was selected for binding assays and additional biological studies to determine its specificity and activity.

The binding affinity of YC137 to Bcl-2 protein was analyzed using fluorescence polarization-based methods (Fig. 1B). We determined that YC137 has a K_i value of 1.3 $\mu\text{mol/L}$ when assayed in Bis-Tris buffer. As a positive control, the 21-residue Bid-BH3 peptide was subjected to the same binding assay conditions (Fig. 1B), and we determined a K_i of 54 nmol/L. Using a similar fluorescence polarization-based method for Bcl- x_L , we determined that YC137 has a K_i value >100 $\mu\text{mol/L}$ (data not shown), suggesting that this small molecule preferentially binds to Bcl-2 protein.

A model of Bcl-2 in complex with YC137 was generated by docking YC137 into the BH3 domain-binding pocket of Bcl-2 (Fig. 2). The three-dimensional structure of the Bcl-2 protein (isoform-2) was obtained from the Protein Data Bank (entry 1GJH), and molecular docking was performed with the GOLD program (version 2.1, distributed by Cambridge Crystallographic Data Center). According to this predicted binding model, the tri-cyclic moiety of YC137 occupies two hydrophobic sites in the Bcl-2-binding pocket and forms hydrogen bonds with residues Arg107, Tyr108, Arg146, and Tyr202 of the Bcl-2 protein.

YC137 Induces Apoptosis of Bcl-2-Dependent Cells. To confirm the specificity of YC137 for the Bcl-2 protein in cellular models, we analyzed its activity in cells dependent on Bcl-2 or Bcl- x_L for survival. FL5.12 hematopoietic progenitors and HCD-57 erythroid cells




Fig. 1. Binding of YC137 to Bcl-2 protein *in vitro*. *A*, structure of YC137 [2-methoxycarbonylamino-4-methylsulfanyl-butiric acid 4-(4,9-dioxo-4,9-dihydro-naphtho[2,3-d]thiazol-2-ylamino)-phenyl ester], a small organic compound that binds Bcl-2. In *B*, His-fused soluble Bcl-2 protein was incubated with a fluorescein-labeled Bid-BH3 peptide in the presence of increasing concentrations of YC137 or unlabeled Bid peptide, and the binding affinity for Bcl-2 was measured by a competitive fluorescence polarization assay. Each data point represents the mean \pm SD of at least three independent experiments.

are dependent on IL-3 and erythropoietin, respectively. In the absence of the growth factor, these cells stop proliferation and undergo apoptosis. However, apoptosis induced by IL-3 or erythropoietin deprivation can be inhibited or delayed in cells overexpressing Bcl-2 or Bcl- x_L (20). On the basis of this, FL5.12 and HCD-57 transfected with Bcl-2 or Bcl- x_L were treated with different concentrations of YC137, and then apoptosis was determined by an immunoenzymatic assay. Fig. 3, *A* and *B*, show Western blot analyses of the overexpressed proteins in transfected cells. After 20 hours of culture in the absence of IL-3 or erythropoietin, YC137 induced apoptosis of Bcl-2-transfected cells in a dose-dependent manner (Fig. 3C). At 0.2 $\mu\text{mol/L}$ of YC137, most of the cells were apoptotic. Conversely, the viability of nontransfected control cells and cells transfected with Bcl- x_L remained unaffected (Fig. 3C), even at concentrations $\leq 5 \mu\text{mol/L}$ (data not shown). Of note, all of the other described Bcl-2/Bcl- x_L inhibitors killed tumor cells at concentrations >20 times higher than those needed for YC137 to induce apoptosis.

To demonstrate the specificity for Bcl-2 in a more physiologically relevant model, we transiently cotransfected CD34⁺ progenitors with pSFFV-Bcl-2 and an EGFP-containing vector. After transfection, cells were cultured in the presence or absence of IL-3 and stem cell factor as survival factors. Then, YC137 was added to the culture and incubated for 24 hours. As shown in Fig. 3D, YC137 did not induce apoptosis of Bcl-2-transfected cells (EGFP⁺) cultured with the survival factors, most likely because of the expression of other antiapoptotic proteins, such as Bcl- x_L . On the contrary, the Bcl-2 inhibitor increased the percentage of apoptosis ~ 2 -fold when transfected cells were cultured in the absence of IL-3 and stem cell factor, indicating that YC137 induces apoptosis of CD34⁺ progenitors when viability of these cells is mainly dependent on Bcl-2, which is consistent with our previous data.

Apoptosis of Breast Cancer Cells Induced by Treatment with YC137 Correlates with the Levels of Bcl-2. We have shown previously that a metastatic variant of an estrogen receptor-negative breast cancer cell line, MB435B, expresses high levels of Bcl-2 as compared with the parental cell line MB435 (19). Thus, we studied

the effect of YC137 on these cells and other breast cancer cell lines (MB435L, MB231, SKBR3, MCF7, SUM159, SUM185, and SUM229) with different levels of Bcl-2 expression. Interestingly, the expression level of Bcl-2 correlated with the apoptotic response to the inhibitor (Fig. 4, *A* and *B*). Thus, MB435B and SUM159 cells, which express the highest levels of Bcl-2 protein, showed also the strongest apoptotic response after 20 hours of treatment with YC137. To determine whether YC137 induces apoptosis via caspase activation, we treated MB435B cells with the inhibitor in the presence of a cell-permeable fluorescent peptide that irreversibly binds to activated caspase 9, a caspase strongly associated with chemotherapy-induced apoptosis. As shown in Fig. 4C, after 2 hours of culture with the inhibitor, activated caspase 9 was present in the great majority ($>85\%$) of tumor cells. Furthermore, we studied whether YC137 induces the release of cytochrome *c* from mitochondria, which is needed for caspase 9 activation. Analysis of soluble and heavy membrane fractions by Western blot revealed that YC137 induced cytochrome *c* release (present in soluble fraction) in MB435B cells treated with the inhibitor for 24 hours (Fig. 4D).

Normal Primary Cell Survival Is Not Modified by Treatment with YC137. A key issue to validate a new chemotherapeutic drug is to determine its tumor specificity. For this purpose, we treated normal small intestine epithelial cells (FHs), primary myoblasts, and PBMCs with YC137. Cells were incubated with increasing concentrations of YC137 (0.1 to 0.5 $\mu\text{mol/L}$) for 20 hours and then analyzed for the presence of histone-associated DNA fragments. As shown in Fig. 5A, apoptosis was not detected in these cells at any of the concentrations tested, whereas MB435B, used here as a positive control, reached the highest level of apoptosis at 0.3 $\mu\text{mol/L}$ YC137.

Ex vivo expanded peripheral blood progenitor cells have been proposed as a source of hematopoietic support to decrease or eliminate the period of neutropenia after high-dose chemotherapy (28). However, this procedure will only be advantageous if contaminating tumor cells are not expanded concomitantly. This model served us to additionally analyze the specificity of the Bcl-2 inhibitor. We cocultured defined numbers of MB435B cells with CD34⁺ cells selected from normal donors in the presence of 0.4 $\mu\text{mol/L}$ YC137. By 20 hours of treatment, 27% of MB435B cells (CD34⁻ cell population) were annexin V positive as determined by flow cytometry analysis,





Fig. 2. Modeled three-dimensional structure of Bcl-2 in complex with YC137. The Bcl-2 protein is represented by a colored ribbon model. YC137 is shown in a ball-and-stick model. Note that Bcl-2 has four residues that form hydrogen bonds with YC137.

Fig. 3. Effect of YC137 on the viability of hematopoietic progenitor cells transfected with Bcl-2 or Bcl-x_L. *A* and *B*, Western blot analysis of the overexpressed protein in FL5.12 and HCD-57 cells transfected with either Bcl-2 or Bcl-x_L. The levels of β-tubulin were determined to assure equal loading. In *C*, cells transfected with Bcl-2 or Bcl-x_L and nontransfected control cells were treated with increasing concentrations of YC137 for 20 hours in the absence of growth factor (IL-3 or erythropoietin), and then apoptotic cell death was determined by an ELISA method that quantifies the histone-associated DNA fragments present in the cytosol. A₄₀₅, units of absorbance at 405 nm. In *D*, CD34⁺ progenitors were transfected with pSFFV-Bcl-2 construct and cultured in the presence or absence of stem cell factor and IL-3. After treatment with YC137 for 24 hours, apoptotic cells were determined by flow cytometry with FITC-labeled annexin V. All data points represent the means ± SD of triplicate analyses.




whereas >95% of hematopoietic progenitors (CD34⁺ cell population) were viable annexin V-negative cells (Fig. 5B).

As shown in Fig. 4C and Fig. 5B, there is a difference in the percentage of apoptotic cells between annexin V and activated caspase 9 assays in MB435B cells treated with YC137. This may reflect the fact that activation of caspase 9 measures early apoptosis, whereas annexin V measures late apoptosis, and that both methods assess different cellular responses to apoptosis.

YC137-Resistant Breast Cancer Cells Down-Regulate the Expression of Bcl-2. Resistance to small molecules that inhibit defined tyrosine kinases has been reported in both cell lines and primary cells (17). As a first step to know whether tumor cells may develop

resistance to Bcl-2 inhibitors, we cultured MB435B cells with increasing concentrations of YC137. After 3 months of culture, a MB435B subline growing at 0.225 μmol/L YC137 (a lethal concentration for parental cells) was isolated. Comparative genomic hybridization analysis revealed no chromosome abnormalities (deletions or amplifications) in the resistant variant (data not shown). Interestingly, these cells had low expression of Bcl-2 at the protein and mRNA level (Fig. 6, A and B); however, the expression of Bcl-x_L was the same as that of the parental cell line. We have described previously that Bcl-2 is transcriptionally regulated by Stat3 in MB435B cells (19). Therefore, we analyzed the DNA binding activity of this transcription factor and found a reduced constitutive activation as measured by electrophoretic

Fig. 4. Effect of YC137 on the viability of breast cancer cell lines. In *A*, breast cancer cell lines were treated with 0.5 μmol/L YC137 for 20 hours, and then apoptotic cell death was quantitated. Histograms represent the means ± SD of triplicate analyses. *B*, analysis of the Bcl-2 protein levels by Western blot. The levels of β-tubulin were determined to assure equal loading. In *C*, MB435B cells were treated with 0.25 μmol/L YC137 for 2 hours in the presence of a fluorescently labeled LEHD-FMK peptide, and then activation of caspase 9 was determined by flow cytometry analysis. In *D*, MB435B cells were treated with 0.25 μmol/L YC137 for 24 hours and then analyzed for the presence of cytochrome *c* in soluble (S) and heavy membrane (M) fractions by Western blot.






Fig. 5. Normal cells are resistant to YC137-induced apoptosis. In A, PBMCs, myoblasts, and FHs small intestine epithelial cells were cultured with increasing concentrations of YC137 for 20 hours, and then apoptosis was determined by an ELISA method as described in Fig. 3. MB435B is included for comparison. Data show a representative experiment ($n = 3$). In B, CD34⁺ progenitors and MB435B cells were cocultured in the presence of 0.4 $\mu\text{mol/L}$ YC137 for 20 hours and then analyzed by flow cytometry with phycoerythrin-labeled annexin V and FITC-labeled anti-CD34 antibodies. Numbers, the percentage of apoptotic cells in the CD34⁺ and CD34⁻ (MB435B) populations.

mobility shift assay (Fig. 6C). As Bcl-2 contributes to interfere with the therapeutic action of many chemotherapeutic drugs, we asked whether a reduced expression of Bcl-2 may render YC137-resistant cells more sensitive to chemotherapy-induced apoptosis. To answer this question, we treated parental and resistant MB435B breast cancer cells with either paclitaxel or adriamycin. Fig. 6D shows that after a 16-hour treatment, YC137-resistant MB435B cells were more sensitive to chemotherapy-induced apoptosis than parental cells, although this difference was more pronounced in Adriamycin-treated cells. The level of apoptosis induced by treatment with adriamycin was as much as eight times higher in the resistant variant than in MB435B cells (Fig. 6D). Similar results were obtained with individual YC137-resistant clones (data not shown).

HER2 Is Down-Regulated in YC137-Resistant MB435B Cells. We have described that EGFR is involved in activation of the Stat3-Bcl-2 transcriptional pathway in MB435B cells (19). Thus, we studied the expression of this membrane receptor in the YC137-resistant subline. However, the levels of EGFR mRNA are the same in both parental and resistant cells (Fig. 7A). Constitutively activated Stat3 is also correlated with up-regulation of HER2, another member of the EGFR family. Interestingly, we found that the mRNA and protein levels of HER2 were clearly down-regulated in the resistant variant (Fig. 7, A and B). This result prompted us to analyze whether HER2 may regulate the expression of Bcl-2 via Stat3. To address this issue, we treated MB435B cells with trastuzumab, a blocking antibody against HER2, and then analyzed the DNA-binding activity of Stat3 and the Bcl-2 expression. As shown in Fig. 7C, treatment of MB435B

with trastuzumab for 6 hours significantly reduced the activation of Stat3, as determined by electrophoretic mobility shift assay, which continued to decrease by 12 hours of treatment. This pattern of Stat3-DNA binding complex formation correlated with the protein and mRNA levels of Bcl-2, which were clearly down-regulated after 24 hours of treatment with trastuzumab (Fig. 7D). Furthermore, consistent with our previous data, trastuzumab-treated cells became more resistant to YC137. Pretreatment with trastuzumab for 24 hours reduced ~ 2 -fold the number of apoptotic MB435B cells cultured in the presence of YC137, as determined by flow cytometry with FITC-labeled annexin V (Fig. 7E).

To additionally confirm this transcriptional pathway in a more direct way, we analyzed the effect of a constitutively activated form of Stat3 (Stat3-C) on the Bcl-2 expression in MB435B treated with trastuzumab. Cells were cotransfected with a vector expressing EGFP and vector encoding Stat3-C. After 36 hours of transfection in the presence of trastuzumab, the control cells had significantly reduced the expression of Bcl-2 ($\sim 35\%$), whereas those expressing Stat3-C had no variation in the levels of Bcl-2 as determined by flow cytometry (Fig. 7F).

DISCUSSION

Most of the currently available small molecule inhibitors are designed to block the enzymatic activity of key proteins in cancer cells (17), and there are very few examples of inhibitors that disrupt protein-protein interactions. Bcl-2 family members play key roles in the regulation of apoptosis and may contribute to cancer development and resistance to chemotherapy, which have attracted intensive inter-




Fig. 6. Mechanism of resistance to YC137 in MB435B cells. A, analysis of Bcl-2 and Bcl-x_L proteins in MB435B cells (C) and the YC137-resistant variant (R) by Western blot. B, expression of Bcl-2 by quantitative real-time PCR analysis. In C, an electrophoretic mobility shift assay was performed using a radiolabeled Stat3 probe to analyze the formation of the Stat3-DNA complex. Nuclear extracts from resistant cells were preincubated with antibodies specific for Stat3 (+ αStat3). In D, control MB435B and YC137-resistant cells were treated with 10 nmol/L paclitaxel (P) for 16 hours or 1 $\mu\text{mol/L}$ adriamycin (A) for 20 hours, and then apoptosis was quantitated as described in Fig. 3. U, untreated cells. Histograms represent the means \pm SD of triplicate analyses.




Fig. 7. Regulation of the Stat3-Bcl-2 pathway by HER2 in MB435B. **A**, analysis of HER2 and EGFR mRNAs in MB435B cells (C) and the YC137-resistant variant (R) by semiquantitative reverse transcription-PCR. **B**, expression of HER2 protein by Western blot analysis. SKBR3 (SK) is included as a positive control. In C, MB435B cells were treated with 50 $\mu\text{g}/\text{mL}$ trastuzumab for the indicated times (in hours), and then an electrophoretic mobility shift assay was performed to analyze the formation of the Stat3-DNA complex. Nuclear extracts from untreated cells were preincubated with antibodies to Stat3 (+ αStat3). In D, the expression of Bcl-2 in MB435B cells was analyzed by real-time PCR and Western blot after 24 hours of treatment with trastuzumab. The levels of β -tubulin were also analyzed to assure equal loading. In E, MB435B cells were pretreated or not with trastuzumab (T) for 24 hours and then incubated in the presence of 0.5 $\mu\text{mol}/\text{L}$ YC137 (YC). After 24 hours of culture, the percentage of annexin V-positive cells was determined by flow cytometry. **C**, control untreated cells. In F, cells were cotransfected with pEGFP-C1 and either Stat3-C or the empty vector (C) and cultured with or without trastuzumab (T). Then, Bcl-2 expression was analyzed in the EGFP⁺ population by flow cytometry. The levels of Bcl-2 in cells treated with trastuzumab were expressed relative to those in untreated cells. Histograms represent the mean \pm SD of triplicate analyses.

est in the discovery of small molecules that inhibit the activity of antiapoptotic proteins of this family, mainly Bcl-2 and Bcl-x_L (13–16). Here, we have designed and synthesized an inhibitor of Bcl-2, YC137, which shows high Bcl-2-binding activity. However, it is much less effective in binding to Bcl-x_L. This is consistent with our data, which demonstrate that YC137 selectively induces apoptosis in breast cancer cells with high levels of Bcl-2. Interestingly, the concentration of YC137 needed for the selective killing of tumor cells is >20 times lower than that reported for the other Bcl-2/Bcl-x_L inhibitors. Of note, a complete apoptotic response was achieved at concentrations about four to six times lower than the K_i value of YC137. There are two main explanations for this discrepancy: (1) the concentration of Bcl-2 protein used in our fluorescence polarization-based binding assays is 120 nmol/L, which might be considerably higher than the cellular concentration of Bcl-2, and (2) YC137 has a poor solubility in our binding assay conditions, and consequently, it is possible that the binding affinity of YC137 was underestimated. The effectiveness of this inhibitor in hematopoietic precursors that become dependent on Bcl-2 for survival suggests that the sensitivity to Bcl-2 inhibitors will be given not only by the levels of Bcl-2 but also the contribution of this antiapoptotic protein to the overall survival of particular tumor cells. Consistently, apoptosis can be induced in a number of tumor cell lines by decreasing the amount of Bcl-2 protein (4, 29). Anticancer drugs used in chemotherapy for tumors and leukemias cause severe toxicity in normal tissues, leading to side effects such as myelosuppression (30). Thus, it is of particular interest our finding that normal cells, including a small intestine epithelial cell line (FHs), myoblasts, PBMcs, and CD34⁺ hematopoietic progenitors, are not sensitive to YC137. A likely explanation is that Bcl-2 is either absent or not relevant for survival in these cell systems. Consistently, FHs cells do not express Bcl-2 protein (data not shown), which is in agreement with data showing that Bcl-2 is absent in the

crypts of the small intestine (31). Additionally, it has been described that Bcl-2 expression is limited to a small proportion (1% to 4%) of neonatal and adult primary muscle cells in culture (32) and that other Bcl-2 family members, such as Mcl-1 and Bcl-x_L, instead of Bcl-2, are critical in the regulation of apoptosis in peripheral blood lymphocytes (33, 34). Furthermore, several authors have shown that in quiescent hematopoietic progenitors, Bcl-2 is not expressed or does not contribute significantly to the survival of this cell population (35, 36). This is the first study that shows the specificity of a Bcl-2 inhibitor for tumor cells as compared with a variety of normal primary cells.

Chronic myelogenous leukemia represents the first human malignancy to be successfully treated with a small molecule inhibitor specific for a tyrosine kinase (17). However, clinical resistance to this inhibitor has been reported in a number of patients, which led to the notion that the effectiveness of other selective inhibitors may be hampered by the presence of resistant tumor cells. We have described a variant of the MB435B cell line that is resistant to YC137-induced apoptosis. This resistant subline has reduced levels of Bcl-2 attributable to down-regulation of the HER2-Stat3 transcriptional pathway, which contributes to chemosensitivity. In support of this, reduced expression of Bcl-2 by antisense oligonucleotides has been shown to sensitize tumor cells to chemotherapeutic drugs (3, 37). We demonstrated previously that Bcl-2 is transcriptionally regulated by Stat3 in MB435B cells (19). However, this pathway seems to be triggered by EGFR, and we show here that resistant cells down-regulate the expression of HER2 but not EGFR. This result may be explained by the formation of receptor complexes between these two members of the EGFR family, which allow the participation of HER2 in apoptosis regulation, because no direct ligand appears to exist for HER2 (38). Trastuzumab, used for the treatment of some forms of breast cancer, is another example of a gene-based cancer drug. This humanized

monoclonal antibody binds HER2 and induces receptor internalization and degradation, inhibition of cell cycle progression, and sensitization to chemotherapy-induced apoptosis (39, 40). We show that trastuzumab down-regulates the Stat3-dependent expression of Bcl-2 in MB435B cells, and consistent with our previous data, these cells become more resistant to YC137. This novel result supports the relevance of Bcl-2 levels in the efficacy of Bcl-2 inhibitors and sheds light on the mechanisms by which trastuzumab may modulate the apoptotic response to chemotherapy in tumor cells. It also identifies other molecular targets, such as Stat3 and Bcl-2, that can lead to a rational design of therapies that increase drug sensitivity.

In conclusion, we have described a small molecule inhibitor of Bcl-2, YC137, and showed for the first time that a Bcl-2-specific inhibitor is able to kill selectively tumor cells that overexpress or depend on this antiapoptotic protein for survival but has no effect on a variety of primary cells. We also describe that cancer cells may develop resistance to YC137. Resistant cells reduce the levels of Bcl-2 and become more sensitive to chemotherapy. These data support the rationale for preclinical and clinical studies of small molecule inhibitors specific for Bcl-2 and represent the first example of *in vitro* selection of cancer cells refractory to a Bcl-2 inhibitor.

ACKNOWLEDGMENTS

We thank Felipe Prosper for providing myoblasts and James Darnell for the Stat3-C plasmid.

REFERENCES

- Reed J, Miyashita T, Takayama S, et al. BCL-2 family proteins: regulators of cell death involved in the pathogenesis of cancer and resistance to therapy. *J Cell Biochem* 1996;60:23–32.
- Reed J. Bcl-2 family proteins: regulators of apoptosis and chemoresistance in hematologic malignancies. *Semin Hematol* 1997;34:9–19.
- Konopleva M, Tari A, Estrov Z, et al. Liposomal Bcl-2 antisense oligonucleotides enhance proliferation, sensitize acute myeloid leukemia to cytosine-arabinoside, and induce apoptosis independent of other antiapoptotic proteins. *Blood* 2000;95:3929–38.
- Koty P, Tyurina Y, Tyurin V, Li S, Kagan V. Depletion of Bcl-2 by an antisense oligonucleotide induces apoptosis accompanied by oxidation and externalization of phosphatidylserine in NCI-H226 lung carcinoma cells. *Mol Cell Biochem* 2002;234–5:125–33.
- Gautschi O, Tschopp S, Olie R, et al. Activity of a novel bcl-2/bcl-xL-bispecific antisense oligonucleotide against tumors of diverse histologic origins. *J Natl Cancer Inst (Bethesda)* 2001;93:463–71.
- Guinness ME, Kenney JL, Reiss M, Lacy J. Bcl-2 antisense oligodeoxynucleotide therapy of Epstein-Barr virus-associated lymphoproliferative disease in severe combined immunodeficient mice. *Cancer Res* 2000;60:5354–8.
- Simoës-Wust A, Schurpf T, Hall J, Stahel R, Zangemeister-Wittke U. Bcl-2/bcl-xL bispecific antisense treatment sensitizes breast carcinoma cells to doxorubicin, paclitaxel and cyclophosphamide. *Breast Cancer Res Treat* 2002;76:157–66.
- Liu Q, Gazitt Y. Potentiation of dexamethasone, taxol and Ad-p53-induced apoptosis by Bcl-2 anti-sense oligodeoxynucleotides in drug-resistant multiple myeloma cells. *Blood* 2003;101:4105–14.
- Huang Z. Bcl-2 family proteins as targets for anticancer drug design. *Oncogene* 2000;19:6627–31.
- Diaz J, Oltersdorf T, Horne W, et al. A common binding site mediates heterodimerization and homodimerization of Bcl-2 family members. *J Biol Chem* 1997;272:11350–5.
- Sattler M, Liang H, Nettlesheim D, et al. Structure of Bcl-xL-Bak peptide complex: recognition between regulators of apoptosis. *Science* 1997;275:983–6.
- Petros AM, Medek A, Nettlesheim DG, et al. Solution structure of the antiapoptotic protein bcl-2. *Proc Natl Acad Sci USA* 2001;98:3012–7.
- Degterev A, Lugovskoy A, Cardone M, et al. Identification of small-molecule inhibitors of interaction between the BH3 domain and Bcl-xL. *Nat Cell Biol* 2001;3:173–82.
- Wang J, Liu D, Zhang Z, et al. Structure-based discovery of an organic compound that binds Bcl-2 protein and induces apoptosis of tumor cells. *Proc Natl Acad Sci USA* 2000;97:7124–9.
- Tzung S, Kim K, Basanez G, et al. Antimycin A mimics a cell-death-inducing Bcl-2 homology domain 3. *Nat Cell Biol* 2001;3:183–91.
- Enyedy I, Ling Y, Nacro K, et al. Discovery of small-molecule inhibitors of Bcl-2 through structure-based computer screening. *J Med Chem* 2001;44:4313–24.
- Shah N, Nicoll J, Nagar B, et al. Multiple BCR-ABL kinase domain mutations confer polyclonal resistance to the tyrosine kinase inhibitor imatinib (STI571) in chronic phase and blast crisis chronic myeloid leukemia. *Cancer Cell* 2002;2:117–25.
- Luzzatto L, Melo J. Acquired resistance to imatinib mesylate: selection for pre-existing mutant cells. *Blood* 2002;100:1105.
- Real PJ, Sierra A, De Juan A, et al. Resistance to chemotherapy via Stat3-dependent overexpression of Bcl-2 in metastatic breast cancer cells. *Oncogene* 2002;21:7611–8.
- Sanz C, Benito A, Inohara N, Ekhterae D, Nunez G, Fernandez-Luna JL. Specific and rapid induction of the proapoptotic protein Hrk after growth factor withdrawal in hematopoietic progenitor cells. *Blood* 1997;89:3199–204.
- Sanz C, Benito A, Silva M, et al. The expression of Bcl-2 is downregulated during differentiation of human hematopoietic progenitor cells along the granulocyte but not the monocyte/macrophage lineage. *Blood* 1997;89:3199–204.
- Rajnoch C, Chachques J, Berrebi A, Bruneval P, Benoit M, Carpentier A. Cellular therapy reverses myocardial dysfunction. *J Thorac Cardiovasc Surg* 2001;121:871–8.
- Nikolovska-Coleska Z, Wang R, Fang X, et al. Development and optimization of a binding assay for the XIAP BIR3 domain using fluorescence polarization. *Anal Biochem* 2004;332:261–73.
- Benito A, Grillot D, Nunez G, Fernandez-Luna JL. Regulation and function of Bcl-2 during differentiation-induced cell death in HL-60 promyelocytic cells. *Am J Pathol* 1995;146:481–90.
- Bromberg J, Wrzeszczynska M, Devgan G, et al. Stat3 as an oncogene. *Cell* 1999;98:295–303.
- Pique M, Barragan M, Dalmau M, Bellosillo B, Pons G, Gil J. Aspirin induces apoptosis through mitochondrial cytochrome c release. *FEBS Lett* 2000;480:193–6.
- Gutierrez O, Pipaon C, Inohara N, et al. Induction of Nod2 in myelomonocytic and intestinal epithelial cells via nuclear factor- κ B activation. *J Biol Chem* 2002;277:41701–5.
- McNiece I, Jones R, Bearman S, et al. Ex vivo expanded peripheral blood progenitor cells provide rapid neutrophil recovery after high-dose chemotherapy in patients with breast cancer. *Blood* 2000;96:3001–7.
- Zangemeister-Wittke U, Leech S, Olie R, et al. A novel bispecific antisense oligonucleotide inhibiting both bcl-2 and bcl-xL expression efficiently induces apoptosis in tumor cells. *Clin Cancer Res* 2000;6:2547–55.
- Buschfort-Papewalis C, Moritz T, Liedert B, Thomale J. Down-regulation of DNA repair in human CD34(+) progenitor cells corresponds to increased drug sensitivity and apoptotic response. *Blood* 2002;100:845–53.
- Merritt AJ, Potten CS, Watson AJ, Loh DY, Nakayama K, Hickman JA. Differential expression of bcl-2 in intestinal epithelia. Correlation with attenuation of apoptosis in colonic crypts and the incidence of colonic neoplasia. *J Cell Sci* 1995;108:2261–71.
- Dominov JA, Dunn JJ, Miller JB. Bcl-2 expression identifies an early stage of myogenesis and promotes clonal expansion of muscle cells. *J Cell Biol* 1998;142:537–44.
- Ohta K, Iwai K, Kasahara Y, et al. Immunoblot analysis of cellular expression of Bcl-2 family proteins, Bcl-2, Bax, Bcl-X and Mcl-1, in human peripheral blood and lymphoid tissues. *Int Immunol* 1995;7:1817–25.
- Lomo J, Smeland EB, Krajewski S, Reed JC, Blomhoff HK. Expression of the Bcl-2 homologue Mcl-1 correlates with survival of peripheral blood B lymphocytes. *Cancer Res* 1996;56:40–3.
- Peters R, Leyvraz S, Perey L. Apoptotic regulation in primitive hematopoietic precursors. *Blood* 1998;92:2041–52.
- Park J, Bernstein I, Hockenbery D. Primitive human hematopoietic precursors express Bcl-x but not Bcl-2. *Blood* 1995;86:868–76.
- Simoës-Wust AP, Schurpf T, Hall J, Stahel RA, Zangemeister-Wittke U. Bcl-2/bcl-xL bispecific antisense treatment sensitizes breast carcinoma cells to doxorubicin, paclitaxel and cyclophosphamide. *Breast Cancer Res Treat* 2002;76:157–66.
- Yarden Y. Biology of HER2 and its importance in breast cancer. *Oncology* 2001;61:1–13.
- Lee S, Yang W, Lan K, et al. Enhanced sensitization to taxol-induced apoptosis by hereceptin pretreatment in ErbB2-overexpressing breast cancer cells. *Cancer Res* 2002;62:5703–10.
- Shawver L, Slamon D, Ullrich A. Smart drugs: tyrosine kinase inhibitors in cancer therapy. *Cancer Cell* 2002;1:117–23.

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

Breast Cancer Cells Can Evade Apoptosis-Mediated Selective Killing by a Novel Small Molecule Inhibitor of Bcl-2

Pedro Jose Real, Yeyu Cao, Renxiao Wang, et al.

Cancer Res 2004;64:7947-7953.

Updated version Access the most recent version of this article at:
<http://cancerres.aacrjournals.org/content/64/21/7947>

Cited articles This article cites 39 articles, 20 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/64/21/7947.full#ref-list-1>

Citing articles This article has been cited by 12 HighWire-hosted articles. Access the articles at:
<http://cancerres.aacrjournals.org/content/64/21/7947.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerres.aacrjournals.org/content/64/21/7947>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.