

Type III IFN Interleukin-28 Mediates the Antitumor Efficacy of Oncolytic Virus VSV in Immune-Competent Mouse Models of Cancer

Phonphimon Wongthida¹, Rosa Maria Diaz^{1,2}, Feorillo Galivo¹, Timothy Kottke¹, Jill Thompson¹, Jose Pulido³, Kevin Pavelko², Larry Pease², Alan Melcher⁴, and Richard Vile^{1,2,4}

Abstract

Innate immune effector mechanisms triggered by oncolytic viruses may contribute to the clearance of both infected and uninfected tumor cells in immunocompetent murine hosts. Here, we developed an *in vitro* tumor cell/bone marrow coculture assay and used it to dissect innate immune sensor and effector responses to intratumoral vesicular stomatitis virus (VSV). We found that the type III IFN interleukin-28 (IL-28) was induced by viral activation of innate immune-sensing cells, acting as a key mediator of VSV-mediated virotherapy of B16ova melanomas. Using tumor variants which differentially express the IL-28 receptor, we showed that IL-28 induced by VSV within the tumor microenvironment sensitizes tumor cells to natural killer cell recognition and activation. These results revealed new insights into the immunovirological mechanisms associated with oncolytic virotherapy in immune-competent hosts. Moreover, they defined a new class of tumor-associated mutation, such as acquired loss of responsiveness to IL-28 signaling, which confers insensitivity to oncolytic virotherapy through a mechanism independent of viral replication *in vitro*. Lastly, the findings suggested new strategies to manipulate immune signals that may enhance viral replication, along with antitumor immune activation, and improve the efficacy of oncolytic virotherapies. *Cancer Res*; 70(11): 4539–49. ©2010 AACR.

Introduction

In theory, introduction of even low levels of a fully replication-competent oncolytic virus into a tumor will allow rapid spread of the virus, lysis of the tumor cells, and reductions in tumor burdens (1, 2). Through either natural, or engineered, selectivity for tumor cells, viral replication should be extinguished in normal cells (3–5). Many different oncolytic viruses have been successfully tested in preclinical and clinical models (6–10). However, such studies have, by necessity, often been carried out in models lacking fully functional immune systems (11–14). Therefore, the immune system has largely been perceived to inhibit this form of anticancer therapy by restricting viral replication, spread, and oncolysis (15–18). However, with the increasing availability of immune competent models of oncolytic virotherapy (19), it is becoming clear that the immune system may play a more positive role—both to prevent uncontrolled viral spread and toxicity (20) and as an effector of antitumor therapy (21–26).

We have previously shown that vesicular stomatitis virus (VSV) therapy of B16ova tumors in C57Bl/6 mice depends on host CD8⁺ and natural killer (NK) cells, confirming that viral replication is not the sole mediator of tumor regressions (27). More recently, we showed that it was not possible to detect progressive viral replication within B16ova tumors following intratumoral injection of VSV (26), probably because of a potent innate immune response at the site of the tumor. Moreover, a single-cycle virus was as effective as a fully replication-competent virus in generating antitumor therapy (21), although expression of the viral genome, at least for a single cycle, was required for therapy (21). Therefore, we hypothesized that VSV acts in this model to trigger a potent innate immune response within the tumor, which rapidly clears the virus and has both direct and indirect antitumor activity (21, 23).

The aim of the current study was to characterize the molecules and effector cells that mediate both the antiviral and antitumor innate response induced by VSV in the B16ova/C57Bl/6 model. Here, we identify interleukin-28 (IL-28), induced by viral activation of innate immune sensing cells, as a key mediator of antitumor therapy in this model. Using cell lines that are equally sensitive to VSV *in vitro*, we show that only tumors that can respond to IL-28 are sensitive to VSV-mediated therapy *in vivo*. Although our data here confirms that the innate response inhibits viral replication and spread (28–30), they also indicate that the same response is, in this case, probably the major mediator of the antitumor effects. Therefore, efforts to suppress the innate

Authors' Affiliations: Departments of ¹Molecular Medicine, ²Immunology, and ³Ophthalmology and Ocular Oncology, Mayo Clinic, Rochester, Minnesota; and ⁴Institute of Molecular Medicine, University of Leeds, Leeds, United Kingdom

Corresponding Author: Richard Vile, Mayo Clinic, Guggenheim 18, 200 First Street Southwest, Rochester, MN 55905. Phone: 507-284-3178; Fax: 507-266-2122; E-mail: vile.richard@mayo.edu.

doi: 10.1158/0008-5472.CAN-09-4658

©2010 American Association for Cancer Research.

response to the virus need to be carefully targeted so that viral replication can be enhanced sufficiently to compensate for the loss of positive antitumor immune effects induced by the virus within the tumor.

Materials and Methods

Cell lines and viruses

B16(LIF) cells are murine melanoma cells (H2-K^b) periodically cultured in our laboratory over the previous 12 years. B16ova melanoma cells (H2-K^b), derived from a separate stock of B16 cells transduced with a cDNA encoding the chicken ovalbumin gene (31), were obtained from Dr. Esteban Celis (H. Lee Moffitt Cancer Center, Tampa, FL) and have been cultured periodically over the previous 7 years. Both cell lines were grown in DMEM (Life Technologies) supplemented with 10% (v/v) FCS (Life Technologies) and L-glutamine (Life Technologies). B16ova cells were maintained in G418 (5 mg/mL) to retain the *ova* gene. All cell lines were free of *Mycoplasma* infection.

VSV-GFP (Indiana serotype) was generated by cloning the green fluorescence protein cDNA into the plasmid pVSV-XN2 as described (32). Purification of viruses was by sucrose gradient centrifugation. Virus titers were measured by plaque assays on BHK-21 cells (32).

Bone marrow coculture

Bone marrow cells were removed from the femurs and tibias of mice by flushing with RPMI 1640 and were passed through a 100- μ m filter to prepare single-cell suspensions. B16(LIF) or B16ova cells (1×10^5) were cocultured with 1×10^6 bone marrow cells for 24 hours at 37°C. Cocultures were then infected with VSV [multiplicity of infection (MOI), 0.1 with respect to the tumor cells] for 6 hours; cells were washed thrice with PBS before addition of anti-VSV neutralizing serum (titer of 1:2,560), recombinant IFN- α protein (R&D Systems), or anti-IL-28 antibody (R&D Systems). Twenty-four to 48 hours later, cell killing was assessed using light microscopy, crystal violet staining, or MTT assay (Cell Proliferation kit, Roche Diagnostics GmbH). Bone marrow cultures were depleted of immune subsets by coculture for 24 hours with anti-MAC3, anti-asialo-GM-1 (Cedarlane), anti-CD8 (Lyt2.4.3), anti-CD4 (GK1.5; Monoclonal Antibody Core Facility, Mayo Clinic), or anti-GR-1 antibodies (BioXcell) to deplete macrophages, NK, CD8 T cells, CD4 T cells, and granulocytic cells, respectively. Depletion of cells from bone marrow cultures was confirmed by fluorescence-activated cell sorting analysis to be in excess of 95% of the relevant cell type compared with nontreated control cultures.

Reverse transcription-PCR and mouse IL-28 receptor cloning

Total RNA was extracted from cells using Qiagen RNA extraction kits. Total cellular RNA (1 μ g) was reverse transcribed in 20 μ L using oligo(dT) as primer. A cDNA equivalent of 1 ng RNA was amplified by PCR using choice-taq DNA polymerase (Denville Scientific, Inc.). Full-length mouse IL-28 receptor (AK148133) was amplified from cDNA from B16ova

cells using forward (5'-TGATCATGATCACAGGGACT-GAAATGTGGCGG-3') and reverse (5'-GCGGCCGCGCACTC-TGGGATTTCGGTCAA-3') primers. The cDNA of IL-28R was inserted into the *Bam*HI and *Not*I sites of the pSIN-CSGW-UbEm lenti-vector (Dr. Yasuhiro Ikeda, Mayo Clinic, Rochester, MN).

In vivo studies

All procedures were approved by the Mayo Foundation Institutional Animal Care and Use Committee. C57Bl/6 mice were purchased from The Jackson Laboratory at 6 to 8 weeks of age. IFN- α / β receptor knockout (KO) mice were a gift from Dr. Roberto Cattaneo (Mayo Clinic, Rochester, MN). To establish subcutaneous tumors, 5×10^5 tumor cells in 100 μ L PBS were injected into the flank of mice. Viral injections (50 μ L) were administered intratumorally at days 7, 9, and 11. Animals were examined daily and tumor sizes were measured thrice weekly using calipers. Animals were killed when tumor size was $>1.0 \times 1.0$ cm in two perpendicular directions. NK cells were depleted by i.p. injections (0.1 mg/mouse) of anti-asialo-GM-1 (Cedarlane; ref. 27). For IL-28 depletion, 2 μ g of anti-IL-28 antibody (R&D Systems) or IgG control (ChromPure Rat IgG; Jackson ImmunoResearch) per mouse was given at each injection of virus.

ELISA for IL-28

Supernatants were tested for IL-28 production by ELISA as directed by the manufacturer (PBL Interferonsource).

Flow cytometry analysis

Tumor-draining lymph nodes and tumors were dissociated to single-cell suspensions. Cells (1×10^6) were washed, resuspended in PBS containing 0.1% bovine serum albumin, and incubated with directly conjugated primary antibodies for 30 minutes at 4°C. Cells were washed and resuspended in 500 μ L PBS containing 4% formaldehyde and analyzed by flow cytometry. Data were analyzed using the Flowjo software (Flowjo). Anti-CD11b FITC, anti-PDCA PE, anti-F4/80 APC, anti-GR1 PE-Cy7, anti-B220 Per-CP, and their respective isotype controls were purchased from BD Pharmingen.

NK cell isolation

NK cells were isolated from peripheral blood mononuclear cells using MACS-negative depletion kits following the manufacturer's protocol ($>90\%$ purity; Miltenyi Biotec).

Statistics

Survival data were analyzed by log-rank test using GraphPad Prism 4 (GraphPad Software). *In vitro* experiments were analyzed using the JMP Software (SAS Institute, Inc.). Statistical significance was determined at the level of $P < 0.05$ (27).

Results

VSV-activated bone marrow cells are cytotoxic against B16ova tumor cells

Our hypothesis was that following intratumoral injection of VSV, host-derived immune cells become activated by




Figure 1. VSV activates bone marrow cytotoxicity against B16ova. **A**, *in vitro* coculture between tumor cells and bone marrow cells; NAb: neutralizing anti-VSV immune serum. **B**, light microscopy of cocultures (from **A**) of B16ova cells 24 h following treatment with (i) VSV MOI 0.1 (no bone marrow), (ii) VSV+NAb (no bone marrow), (iii) bone marrow+NAb, (iv) bone marrow cells alone (no VSV or NAb), or (v) bone marrow (BM)+VSV+NAb.

the virus to secrete cytokines, which subsequently recruit/activate the same, or additional, antiviral and antitumor immune effector cells and mechanisms. To test this hypothesis, we established an assay in which host-derived bone marrow cells were cocultured with B16-derived tumor cells in the presence, or absence, of virus (Fig. 1A). Exposure of B16ova tumor cells to VSV, even at low MOI, induced rapid and extensive cytotoxicity (Fig. 1B, i), which could be inhibited by coculturing with anti-VSV neutralizing serum (Fig. 1B, ii). Coculture of B16ova cells with bone marrow cells from C57Bl/6 mice had no significant cytotoxic effects either with (Fig. 1B, iii) or without (Fig. 1B, iv) the neutralizing serum against VSV. In contrast, when cocultures of B16ova and C57-derived bone marrow cells were exposed to VSV in the presence of anti-VSV neutralizing serum, extensive cytotoxicity against the tumor cells was observed (Fig. 1B, v). These data indicated that the VSV-mediated activation of bone marrow cells induced bystander killing

of B16ova tumor cells, distinct from direct viral-mediated oncolysis.

B16 variants show differential sensitivity to VSV-activated bone marrow cytotoxicity

In contrast to B16ova cells, we observed that one of our B16-derived tumor cell lines, B16(LIF), was almost completely resistant to the cytotoxic effects of VSV-activated bone marrow cells (Fig. 2A and B). However, both B16ova and B16(LIF) cells showed almost equal sensitivity to cell-free VSV replication and cytolysis *in vitro* (Fig. 2B). Although B16ova and B16(LIF) cells are both B16 derived, they have been passaged separately for several years in our laboratory (see Materials and Methods). Interestingly, the lack of sensitivity of B16(LIF) cells to VSV-activated bone marrow cell killing *in vitro* was mirrored by a consistent lack of efficacy of antitumor therapy following intratumoral injection of VSV *in vivo* ($P = 0.18$ compared with treatment with heat-inactivated VSV; Fig. 2C, i). In contrast, B16ova cells were sensitive both to VSV-activated bone marrow cell killing *in vitro* (Fig. 2A and B) and to intratumoral VSV treatment *in vivo*, in which we consistently observed significant prolongation of survival times over control treatments ($P < 0.00001$ compared with treatment with heat-inactivated VSV; Fig. 2C, ii).

To investigate further how B16ova and B16(LIF) cells may have diverged over several years in culture, we performed a gene chip analysis. Of the multiple genetic differences that exist between these two cell lines, we focused on genes associated with the innate immune response to viral infection. One such gene, the receptor for the type III IFN IL-28, was overexpressed in B16ova cells by a factor of 23-fold relative to its levels of expression in B16(LIF) cells, as confirmed by reverse transcription-PCR (Fig. 2D). We confirmed also that B16(LIF) cells do not express the IL-28R on infection with VSV at a MOI ranging from 0.1, 1.0, or 10.

IL-28, as well as type I IFNs, mediate VSV-activated bone marrow cytotoxicity

We hypothesized that differences in the expression of the IL-28R between the B16ova and B16(LIF) lines may mediate the differential responses of these cell lines to both VSV-activated bone marrow cytotoxicity *in vitro* as well as VSV-mediated antitumor therapy *in vivo*. Consistent with this hypothesis, antibody-mediated blockade of IL-28 signaling abrogated the VSV-activated bone marrow-mediated killing of B16ova cells *in vitro* (Fig. 3A and B). We also observed that when the bone marrow cytotoxicity assay was repeated, even in the absence of neutralizing VSV serum, components of normal C57Bl/6 bone marrow suppressed viral replication through the B16ova cultures [mean titers reduced from 1.8×10^5 plaque-forming unit (pfu)/mL with no bone marrow cells to ~ 100 pfu/mL in the presence of bone marrow cells; Fig. 3A]. However, although blockade of IL-28 prevented VSV-activated bone marrow-mediated cytotoxicity against B16ova cells, it did not reduce VSV replication/spread through B16ova cultures even in the presence or absence of bone marrow cells (Fig. 3A).





Figure 2. B16(LIF) cells are insensitive to VSV-activated bone marrow cytotoxicity. A and B, B16ova or B16(LIF) cells were cocultured with C57Bl/6 bone marrow with, or without, VSV as in Fig. 1A. Cell survival was assessed by crystal violet staining (A) or MTT assay (B). C, C57Bl/6 (8 mice/group) bearing 7-d subcutaneous (s.c.) B16(LIF) (i) or B16ova (ii) tumors were injected intratumorally every 2 d (three total) with 5×10^8 pfu of heat-inactivated virus (HI) or VSV. Survival with time is shown. D, cDNA from B16ova or B16(LIF) cells was screened for IL-28R. A weak signal was detected in B16(LIF) cells with 20 additional cycles. **, $P < 0.01$; ***, $P < 0.001$; ns; non significant.

Type I IFNs (IFN α/β) exert potent antiviral effects against VSV (11, 14, 33, 34). To identify the factor in bone marrow cocultures that inhibited viral spread/replication through highly permissive B16ova cells, the assays of Fig. 3A and B were repeated using bone marrow cells from IFN α/β receptor KO mice. The inability of bone marrow cells to respond to IFN

α/β completely abolished VSV-activated cytotoxicity against B16ova cells *in vitro* (Fig. 3C, i and ii). In addition, although IFN- α was not itself directly cytotoxic to B16ova tumor cells at 1 or 100 U/mL (data not shown), addition of IFN- α to IFN α/β receptor KO bone marrow/tumor cell cocultures replaced the requirement for VSV to induce *in vitro* cytotoxicity against B16ova



tumor cells (Fig. 3C, v and vi). Finally, IFN α/β receptor KO bone marrow was unable to suppress VSV spread and replication in B16ova cocultures (Fig. 3D) and blockade of IL-28 in these cultures had no inhibitory effect on viral replication (Fig. 3D).

Taken together, these data indicate that type I IFNs are critical for both the VSV-activated bone marrow cytotoxicity against B16ova cells and for the suppression of VSV replication through these cultures; in contrast, although the type III IFN IL-28 mediates VSV-activated bone marrow cytotoxicity against B16ova, it does not exert significant inhibitory effects on viral replication and spread.

GR1⁺ cells sense VSV infection

We confirmed that IL-28 is produced by normal C57Bl/6 bone marrow cultures, but only when activated by VSV (Fig. 4A). Depletion of the bone marrow cultures of both macrophages and GR1⁺ cells reproducibly significantly reduced the levels of VSV-activated IL-28 secretion ($P < 0.01$ for macrophages and $P < 0.001$ for GR-1+ cells compared with no depletion; Fig. 4A). To determine the source of the IL-28 observed in Fig. 4A, this experiment was repeated in the absence of cocultured B16ova tumor cells. Within 48 hours of infection of the bone marrow cell cultures (no added




Figure 4. GR1⁺ cells in bone marrow sense VSV. **A**, bone marrow from C57Bl/6 mice was left undepleted (none) or depleted of macrophages, NK, CD8, CD4, or GR1⁺ cells and cocultured with B16ova cells +/- VSV (MOI 0.1) as in Fig. 1A. Forty-eight hours after addition of VSV, supernatants were assayed for IL-28. **B**, C57Bl/6 (3 mice/group) bearing B16ova tumors were injected intratumorally with 5×10^6 pfu of VSV (filled column) or PBS (opened column). Tumors (i) and draining lymph node (LN; ii and iii) were harvested at times shown and analyzed for (i) neutrophils (CD11b⁺GR1⁺F4/80⁺), CD11b⁺GR1⁺ (ii), or plasmacytoid dendritic cells (CD11b⁺GR1⁺PDCA⁺; iii). Infiltrates in tumors injected intratumorally with heat-inactivated VSV as a negative control were directly comparable with those tumors treated with PBS (data not shown). TM, tumor; DC, dendritic cells. **C**, C57Bl/6 bone marrow was left undepleted (i) or depleted of macrophages (ii) or GR1⁺ cells (iii), and cocultured with B16ova with VSV (MOI, 0.1) and neutralizing anti-VSV antibody (NAb) as in Fig. 1A. Forty-eight hours after the addition of VSV, cell survival was visualized or quantified by MTT assay (**D**). *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

B16ova), levels of IL-28 detected by ELISA were very similar ($\sim 1,100$ pg/mL/ 10^6 bone marrow cells) to those seen in the presence of cocultured B16ova cells. In addition, the dependence of IL-28 production upon both macrophages (Mac+) and GR-1+ cells was identical to that seen on coculture with tumor cells. These data confirm that the IL-28 produced by the B16 tumor cell/GR-1+ bone marrow cultures upon exposure to VSV is derived predominantly from the bone marrow cells, rather than from either the tumor cells, directly or as a result of a factor induced by the tumor cells upon VSV infection that then acts on the bone marrow cells.

VSV activation of bone marrow cells to produce IL-28 was also dependent on type I IFN signaling because bone marrow from IFN α/β KO mice did not secrete IL-28 upon exposure to VSV (<100 pg/mL in the assay of Fig. 4A; data not shown). To relate these *in vitro* data with the *in vivo* mechanisms that mediate the treatment of B16ova tumors (Fig. 2C), we investigated which effector cells respond to VSV. Intratumoral VSV induced significant infiltration into the tumor by neutrophils (CD11b⁺;GR1⁺;F4/80⁻) within 6 hours of virus injection (Fig. 4B, i). In addition, both CD11b⁺;GR1⁺ cells and plasmacytoid dendritic cells (CD11b⁻;GR1⁺;PDCA⁺; Fig. 4B, ii and iii) accumulated in tumor-draining lymph nodes over longer periods (12–72 h) following virus injection. Consistent with our observations of the importance of GR1⁺ cells in sensing VSV infection through the production of IL-28 *in vitro* (Fig. 4A) and of their recruitment *in vivo* (Fig. 4B), depletion of GR1⁺ cells significantly inhibited the *in vitro* cytotoxicity of bone marrow cultures against B16ova tumor cells ($P = 0.0004$ compared with no depletion; Fig. 4C and D). Although we did not observe reproducible increases in macrophage infiltration into VSV-injected B16ova tumors *in vivo* (data not shown), depletion of Mac3⁺ cells from bone marrow cultures also inhibited VSV-activated bone marrow-induced cytotoxicity against B16ova cells *in vitro* ($P = 0.005$ compared with no depletion; Fig. 4C, i and ii, and D). Depletion of CD4, CD8, or B220 cells from bone marrow cultures had no significant effect on the VSV-activated bone marrow cytotoxicity against B16ova cells *in vitro* (data not shown).

Taken together, these data indicate that the VSV-mediated activation of both macrophages and GR1⁺ cells within bone marrow cultures induces IL-28 and cytotoxicity against B16ova tumor cells *in vitro*, consistent with our *in vivo* observations that VSV activates the recruitment of GR1⁺ cells to both B16ova tumors and tumor-draining lymph node *in vivo*.

Expression of IL-28R by B16ova sensitizes B16ova cells for NK recognition

Although NK1.1 cells were not a reproducible source of IL-28 from VSV-activated bone marrow cultures (Fig. 4A), depletion of these cells from bone marrow cultures abolished cytotoxicity against B16ova tumor cells (Fig. 4D) almost to the same extent as blockade of IL-28 (Fig. 3A and B). Therefore, we investigated whether there was a mechanistic link between VSV-induced, IL-28 expression and tumor cell sensitivity to NK cells. Whereas IL-28 induced expression of at least three different NK cell ligands on B16ova cells (Fig. 5A), it had no effect on the expression of NK ligands

on B16(LIF) cells (Fig. 5B). In addition, B16ova cells treated with either recombinant IL-28 or with conditioned medium from VSV-activated bone marrow cultures activated IFN- γ production from NK cells (Fig. 5C). In contrast, B16(LIF) cells were not effective targets for NK cells, even in the presence of IL-28 (Fig. 5C). Consistent with these *in vitro* findings, VSV-mediated therapy of B16ova tumors in mice depleted of NK cells was significantly decreased compared with that seen in VSV-treated, nondepleted animals ($P = 0.01$; Fig. 5D).

Taken together, these data show that IL-28 treatment of B16ova cells induces both the expression of NK ligands and NK cell activation, and that these responses correlate very well with both VSV-activated, IL-28-dependent bone marrow cytotoxicity against B16ova *in vitro*, as well as the NK dependence of VSV-mediated therapy of B16ova tumors *in vivo*.

IL-28/IL-28R expression mediates VSV therapy

Finally, we tested the role of IL-28 in the VSV treatment of tumors *in vivo*. B16ova tumor lysates expressed IL-28 only following exposure to VSV *in vivo* (Fig. 6A). When VSV-induced intratumoral IL-28 was neutralized by antibody blockade, antitumor efficacy was lost and was no better than control treatments ($P = 0.0002$ VSV+ control IgG versus VSV+anti-IL-28Ab; Fig. 6B). Our hypothesis from the gene chip analysis was that B16(LIF) tumors are not amenable to treatment with VSV *in vivo* (Fig. 2C) because of a lack of expression of the IL-28R (Fig. 2D). To test this hypothesis, we transduced B16(LIF) cells with a lentiviral vector expressing the IL-28R. Unlike parental B16(LIF) cells, B16(LIF)-IL-28R populations were as effective at activating NK cells as were B16ova cells, following treatment with IL-28 in the assay of Fig. 5C (inducing >700 pg/mL of IFN- γ from cocultured NK cells when treated with IL-28). As usual, intratumoral injection of VSV into established subcutaneous B16(LIF) tumors was not significantly more effective than control treatments (Fig. 6C). However, although nontreated B16(LIF)-IL-28R tumors grew at an identical rate to B16(LIF) tumors (data not shown), VSV treatment of B16(LIF)-IL-28R tumors induced highly significant delays of tumor growth [$P = 0.0008$ compared with VSV-treated B16(LIF)] and extended the survival times of mice to levels similar to those seen with VSV treatment of B16ova tumors (Figs. 2C and 6B). Therefore, blockade of VSV-induced IL-28 expression *in vivo* inhibits VSV-mediated antitumor therapy of B16ova tumors and, by restoring the expression of the IL-28R to B16(LIF) cells, this VSV-insensitive tumor was converted into one in which VSV has antitumor efficacy.

Discussion

Our data are consistent with a model in which the innate immune response to intratumorally injected VSV mediates therapy in the B16ova/C57Bl/6 immune-competent model, in large part through the activity of the type III IFN IL-28. These findings are consistent with our previous reports that the therapeutic efficacy of oncolytic VSV in this model requires host-derived cells such as NK cells (27); is not dependent on multiple rounds of virus replication; and is

associated with a rapidly evolving, and resolving, innate response to virus in the tumor (21, 23).

We show here that a subset of bone marrow-derived cells sense virus presence/infection by secreting type I and III IFNs. Additional immune subsets then respond by clearing infected as well as uninfected bystander cells, thereby suppressing ongoing viral infection. We believe that this cytotoxicity against tumor cells is largely responsible for the antitumor effects of VSV treatment of B16ova tumors (Fig. 2C).

The B16ova and B16(LIF) B16-derived cell lines have been cultured separately for several years but show almost identi-

cal sensitivity to VSV *in vitro* (Fig. 2). However, although we consistently achieve therapy of B16ova tumors with intratumoral VSV, B16(LIF) tumors are insensitive to VSV-mediated therapy (Fig. 2C). Using gene chip analysis, we identified a possible molecule, the receptor for IL-28, which might explain this difference (Fig. 2D).

IL-28, or IFN- λ , a newly identified class two cytokine receptor ligand (35–37), is produced in response to viral infection and has antiviral activity (35–37). Although IL-28 signaling pathways are similar to those of IFN type I (38), IL-28 interacts with a different receptor complex constituted by




Figure 5. IL-28 activates the NK recognition of B16ova. A and B, B16ova (A) or B16(LIF) (B) were cultured +/- recombinant IL-28 for 24 h and cDNA was screened for expression of NK ligands RAE, H60, and MULT-1. *, signal detected upon additional 20 cycles. GAPDH, glyceraldehyde-3-phosphate dehydrogenase. C, NK cells were cocultured with B16ova (a–i) or B16(LIF) (a'–i') +/- IL-28 or 48-h conditioned medium from C57Bl/6 bone marrow cells exposed to VSV (MOI, 0.1). Twenty-four hours later, supernatants were assayed for IFN- γ . D, mice ($n = 8$ /group) bearing 4d B16ova s.c. tumors were depleted of NK cells by anti-asialo-GM-1 antibody or mock depleted (control IgG). VSV was injected intratumorally on days 7, 9, and 11. **, $P < 0.01$.




Figure 6. IL-28 signaling mediates the VSV therapy of B16ova tumors. A, freeze thaw lysates of *in vitro* cultured B16ova or B16ova infected with VSV (MOI, 0.1) 24 h previously or of 7d-established B16ova tumors from mice treated intratumorally with either PBS or VSV (5×10^8 pfu) 24 h previously were assayed for IL-28. B, C57Bl/6 mice ($n = 8$ /group) bearing 7d-established subcutaneous B16ova tumors were injected intratumorally at days 7, 9, and 11 with 5×10^8 pfu of heat-inactivated virus (HI) or VSV along with anti-IL-28 or control antibody. C, C57Bl/6 mice ($n = 8$ /group) bearing 7d-established B16(LIF) or B16(LIF)-IL-28R tumors were injected intratumorally with 5×10^8 pfu of heat-inactivated VSV (HI-VSV) or VSV on days 7, 9, and 11. ***, $P < 0.001$.

the IL-10 receptor β (IL-10R β) and IL-28R α chains (35, 36). IL-28 also has an antitumor activity in many tumor models (39–41).

B16(LIF) cells express very low levels of IL-28R (Fig. 2D), are not killed by VSV-activated bone marrow cells *in vitro* (Fig. 2A and B), do not respond to IL-28 by inducing NK ligands (Fig. 5B), cannot activate NK cells *in vitro* (Fig. 5C), and are not sensitive to VSV *in vivo* (Fig. 2C). In contrast, B16ova cells express high levels of IL-28R (Fig. 2D), are sensitive to VSV-activated bone marrow cytotoxicity (Figs. 1–3), upregulate NK ligands, activate NK cells in response to IL-28 (Fig. 5), and are sensitive to intratumoral VSV (Fig. 2C). These data explain our previous findings that NK cells mediate the VSV therapy of B16ova tumors (27). Importantly, neutralization of VSV-induced IL-28 abrogated VSV therapy *in vivo* (Fig. 6A and B). Finally, a VSV-insensitive B16(LIF) tumor was converted into a VSV-responsive tumor

by overexpressing the IL-28R in the B16(LIF) parental cells (Fig. 6C).

These data show that innate immune signaling through VSV-induced IL-28, rather than viral replication/oncolysis, is a critical mediator of antitumor therapy with this oncolytic virus. Future efforts aimed at suppressing the innate response as an adjunct to virotherapy should, therefore, take into account that important immune adjuvant benefits may be lost in the process.

Our data also suggest a hierarchy of control and separation of function between the type I and type III IFNs in the response to VSV. Bone marrow from IFN- α/β R KO mice did not express IL-28 in response to VSV, suggesting that the type I IFN response is an upstream controller of the type III response. In addition, coculture of B16ova cells with bone marrow cells of IFN- α/β R KO mice failed to suppress VSV replication *in vitro* (Fig. 3D). Similarly, VSV injected into

B16ova tumors in IFN- α / β R KO mice led to viral replication, dissemination, and rapid fatal toxicity. Thus, type I IFNs are highly inhibitory to spread/replication of VSV, as previously described (14, 28, 42). However, blockade of IL-28 *in vitro* neither increased viral replication (Fig. 3A) nor did it induce viral dissemination/toxicity *in vivo* (Fig. 6B), consistent with reports that IL-28R KO mice survive VSV infections as well as wild-type mice (43). However, IL-28 was critically important for VSV-mediated antitumor therapy (Fig. 6). Ank and colleagues (43) have previously shown that expression, or otherwise, of the IL-28R is a critical determinant of the ability of cells to acquire sensitivity to the antiviral effects of IL-28. Our results here extend this concept into the fields of both tumor immunology and oncolytic virotherapy, and suggest that screening for IL-28R expression may be a critical prognostic feature for predicting effective responses to oncolytic virotherapy.

We have shown that GR1⁺ cells sense intratumoral VSV by secreting IL-28 (Fig. 4A) to trigger antitumor effector mechanisms, including the activation of NK cell recognition of tumor cells (Fig. 5). GR1⁺ plasmacytoid dendritic cells secrete IFNs (44–47), which regulate the acute cellular response to viral infection with VSV. Consistent with this, abrogation of VSV-mediated activation of cytotoxicity against B16ova cells in bone marrow cultures depleted of GR1⁺ cells (Fig. 4C) was rescued by the addition of IFN- α .

Finally, our results show that screening tumor cells for their *in vitro* sensitivity to an oncolytic virus may not reflect their sensitivities *in vivo*, such as the case with B16ova

and B16(LIF) cells (Fig. 2). This will facilitate the evaluation of patient tumors, on an individualized basis, to predict whether oncolytic virotherapy is likely to be effective.

In summary, we show here that viral-mediated activation of innate immune signaling is critical for the VSV therapy of B16ova tumors in immune-competent mice. These studies open new avenues to understand how oncolytic viruses interact with the host immune system and how it may be possible selectively to manipulate key immune signals to enhance both viral replication and antitumor immune activation to improve efficacy of oncolytic virotherapies.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

We thank Toni Higgins for the expert secretarial assistance.

Grant Support

The Richard M. Schulze Family Foundation, the Mayo Foundation, and by NIH grants CA107082, CA130878, and CA132734.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 12/22/2009; revised 02/18/2010; accepted 03/17/2010; published OnlineFirst 05/18/2010.

References

- Kim D, Martuza RL, Zwiebel J. Replication-selective virotherapy for cancer: biological principles, risk management and future directions. *Nat Med* 2001;7:781–7.
- Parato KA, Senger D, Forsyth PA, Bell JC. Recent progress in the battle between oncolytic viruses and tumours. *Nat Rev Cancer* 2005;5:965–76.
- Everts B, van der Poel HG. Replication-selective oncolytic viruses in the treatment of cancer. *Cancer Gene Ther* 2005;12:141–61.
- Martuza RL, Malick A, Markert JM, Ruffner KL, Coen DM. Experimental therapy of human glioma by means of a genetically engineered virus mutant. *Science* 1991;252:854–6.
- Sinkovics JG, Horvath JC. Natural and genetically engineered viral agents for oncolysis and gene therapy of human cancers. *Arch Immunol Ther Exp (Warsz)* 2008;56 Suppl 1:3–59s.
- Aghi M, Martuza RL. Oncolytic viral therapies—the clinical experience. *Oncogene* 2005;24:7802–16.
- Ganly I, Kim D, Eckhardt G, et al. A phase I study of Onyx-015, an E1B attenuated adenovirus, administered intratumorally to patients with recurrent head and neck cancer. *Clin Cancer Res* 2000;6:798–806.
- Hadaschik BA, Zhang K, So AI, et al. Oncolytic vesicular stomatitis viruses are potent agents for intravesical treatment of high-risk bladder cancer. *Cancer Res* 2008;68:4506–10.
- Liu TC, Galanis E, Kim D. Clinical trial results with oncolytic virotherapy: a century of promise, a decade of progress. *Nat Clin Pract Oncol* 2007;4:101–17.
- Stanford MM, McFadden G. Myxoma virus and oncolytic virotherapy: a new biologic weapon in the war against cancer. *Expert Opin Biol Ther* 2007;7:1415–25.
- Balachandran S, Barber GN. Vesicular stomatitis virus (VSV) therapy of tumors. *IUBMB Life* 2000;50:135–8.
- Grote D, Russell SJ, Cornu TI, et al. Live attenuated measles virus induces regression of human lymphoma xenografts in immunodeficient mice. *Blood* 2001;97:3746–54.
- Peng KW, Ahmann GJ, Pham L, Greipp PR, Cattaneo R, Russell SJ. Systemic therapy of myeloma xenografts by an attenuated measles virus. *Blood* 2001;98:2002–7.
- Stojdl DF, Lichty B, Knowles S, et al. Exploiting tumor-specific defects in the interferon pathway with a previously unknown oncolytic virus. *Nat Med* 2000;6:821–5.
- Breitbach CJ, Paterson JM, Lemay CG, et al. Targeted inflammation during oncolytic virus therapy severely compromises tumor blood flow. *Mol Ther* 2007;15:1686–93.
- Haralambieva I, Iankov I, Hasegawa K, Harvey M, Russell SJ, Peng KW. Engineering oncolytic measles virus to circumvent the intracellular innate immune response. *Mol Ther* 2007;15:588–97.
- Nguyen TL, Abdelbary H, Arguello M, et al. Chemical targeting of the innate antiviral response by histone deacetylase inhibitors renders refractory cancers sensitive to viral oncolysis. *Proc Natl Acad Sci U S A* 2008;105:14981–6.
- Wein LM, Wu JT, Kim DH. Validation and analysis of a mathematical model of a replication-competent oncolytic virus for cancer treatment: implications for virus design and delivery. *Cancer Res* 2003;63:1317–24.
- Thomas MA, Spencer JF, La Regina MC, et al. Syrian hamster as a permissive immunocompetent animal model for the study of oncolytic adenovirus vectors. *Cancer Res* 2006;66:1270–6.
- Qiao J, Wang H, Kottke T, et al. Cyclophosphamide facilitates

- antitumor efficacy against subcutaneous tumors following intravenous delivery of reovirus. *Clin Cancer Res* 2008;14:259–69.
21. Galivo F, Diaz RM, Wongthida P, et al. Single-cycle viral gene expression, rather than progressive replication and oncolysis, is required for VSV therapy of B16 melanoma. *Gene Ther* 2009.
 22. Grote D, Cattaneo R, Fielding AK. Neutrophils contribute to the measles virus-induced antitumor effect: enhancement by granulocyte macrophage colony-stimulating factor expression. *Cancer Res* 2003;63:6463–8.
 23. Prestwich RJ, Errington F, Diaz RM, et al. The case of oncolytic viruses versus the immune system: waiting on the judgment of Solomon. *Hum Gene Ther* 2009;20:1119–32.
 24. Thomas DL, Fraser NW. HSV-1 therapy of primary tumors reduces the number of metastases in an immune-competent model of metastatic breast cancer. *Mol Ther* 2003;8:543–51.
 25. Toda M, Rabkin SD, Kojima H, Martuza RL. Herpes simplex virus as an *in situ* cancer vaccine for the induction of specific anti-tumor immunity. *Hum Gene Ther* 1999;10:385–93.
 26. White CL, Twigger KR, Vidal L, et al. Characterization of the adaptive and innate immune response to intravenous oncolytic reovirus (Dearing type 3) during a phase I clinical trial. *Gene Ther* 2008;15:911–20.
 27. Diaz RM, Galivo F, Kottke T, et al. Oncolytic immunovirotherapy for melanoma using vesicular stomatitis virus. *Cancer Res* 2007;67:2840–8.
 28. Barber GN. VSV-tumor selective replication and protein translation. *Oncogene* 2005;24:7710–9.
 29. Friedman A, Tian JP, Fulci G, Chiocca EA, Wang J. Glioma virotherapy: effects of innate immune suppression and increased viral replication capacity. *Cancer Res* 2006;66:2314–9.
 30. Wakimoto H, Ikeda K, Abe T, et al. The complement response against an oncolytic virus is species-specific in its activation pathways. *Mol Ther: J Am Soc Gene Ther* 2002;5:275–82.
 31. Linardakis E, Bateman A, Phan V, et al. Enhancing the efficacy of a weak allogeneic melanoma vaccine by viral fusogenic membrane glycoprotein-mediated tumor cell-tumor cell fusion. *Cancer Res* 2002;62:5495–504.
 32. Fernandez M, Porosnicu M, Markovic D, Barber GN. Genetically engineered vesicular stomatitis virus in gene therapy: application for treatment of malignant disease. *J Virol* 2002;76:895–904.
 33. Balachandran S, Roberts PC, Kipperman T, et al. α/β interferons potentiate virus-induced apoptosis through activation of the FADD/Caspase-8 death signaling pathway. *J Virol* 2000;74:1513–23.
 34. Stojdl DF, Lichty BD, tenOever BR, et al. VSV strains with defects in their ability to shutdown innate immunity are potent systemic anti-cancer agents. *Cancer Cell* 2003;4:263–75.
 35. Kotenko SV, Gallagher G, Baurin VV, et al. IFN- λ s mediate antiviral protection through a distinct class II cytokine receptor complex. *Nat Immunol* 2003;4:69–77.
 36. Sheppard P, Kindsvogel W, Xu W, et al. IL-28, IL-29 and their class II cytokine receptor IL-28R. *Nat Immunol* 2003;4:63–8.
 37. Vilcek J. Novel interferons. *Nat Immunol* 2003;4:8–9.
 38. Dumoutier L, Tounsi A, Michiels T, Sommereyns C, Kotenko SV, Renauld JC. Role of the interleukin (IL)-28 receptor tyrosine residues for antiviral and antiproliferative activity of IL-29/interferon- λ 1: similarities with type I interferon signaling. *J Biol Chem* 2004;279:32269–74.
 39. Lasfar A, Lewis-Antes A, Smirnov SV, et al. Characterization of the mouse IFN- λ ligand-receptor system: IFN- λ s exhibit antitumor activity against B16 melanoma. *Cancer Res* 2006;66:4468–77.
 40. Numasaki M, Tagawa M, Iwata F, et al. IL-28 elicits antitumor responses against murine fibrosarcoma. *J Immunol* 2007;178:5086–98.
 41. Sato A, Ohtsuki M, Hata M, Kobayashi E, Murakami T. Antitumor activity of IFN- λ in murine tumor models. *J Immunol* 2006;176:7686–94.
 42. Barber GN. Vesicular stomatitis virus as an oncolytic vector. *Viral Immunol* 2004;17:516–27.
 43. Ank N, Iversen MB, Bartholdy C, et al. An important role for type III interferon (IFN- λ /IL-28) in TLR-induced antiviral activity. *J Immunol* 2008;180:2474–85.
 44. Asselin-Paturel C, Boonstra A, Dalod M, et al. Mouse type I IFN-producing cells are immature APCs with plasmacytoid morphology. *Nat Immunol* 2001;2:1144–50.
 45. Barchet W, Cella M, Colonna M. Plasmacytoid dendritic cells-virus experts of innate immunity. *Semin Immunol* 2005;17:253–61.
 46. Cella M, Jarrossay D, Facchetti F, et al. Plasmacytoid monocytes migrate to inflamed lymph nodes and produce large amounts of type I interferon. *Nat Med* 1999;5:919–23.
 47. Nakano H, Yanagita M, Gunn MD. CD11c(+)B220(+)Gr-1(+) cells in mouse lymph nodes and spleen display characteristics of plasmacytoid dendritic cells. *J Exp Med* 2001;194:1171–8.

Correction: Online Publication Dates for *Cancer Research* April 15, 2010 Articles

The following articles in the April 15, 2010 issue of *Cancer Research* were published with an online publication date of April 6, 2010 listed, but were actually published online on April 13, 2010:

Garmy-Susini B, Avraamides CJ, Schmid MC, Foubert P, Ellies LG, Barnes L, Feral C, Papayannopoulou T, Lowy A, Blair SL, Cheresh D, Ginsberg M, Varner JA. Integrin $\alpha 4 \beta 1$ signaling is required for lymphangiogenesis and tumor metastasis. *Cancer Res* 2010;70:3042–51. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3761.

Vincent J, Mignot G, Chalmin F, Ladoire S, Bruchard M, Chevriaux A, Martin F, Apetoh L, Rébé C, Ghiringhelli F. 5-Fluorouracil selectively kills tumor-associated myeloid-derived suppressor cells resulting in enhanced T cell-dependent antitumor immunity. *Cancer Res* 2010;70:3052–61. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3690.

Nagasaka T, Rhee J, Kloor M, Gebert J, Naomoto Y, Boland CR, Goel A. Somatic hypermethylation of *MSH2* is a frequent event in Lynch syndrome colorectal cancers. *Cancer Res* 2010;70:3098–108. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3290.

He X, Ota T, Liu P, Su C, Chien J, Shridhar V. Downregulation of HtrA1 promotes resistance to anoikis and peritoneal dissemination of ovarian cancer cells. *Cancer Res* 2010;70:3109–18. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3557.

Fiorentino M, Judson G, Penney K, Flavin R, Stark J, Fiore C, Fall K, Martin N, Ma J, Sinnott J, Giovannucci E, Stampfer M, Sesso HD, Kantoff PW, Finn S, Loda M, Mucci L. Immunohistochemical expression of BRCA1 and lethal prostate cancer. *Cancer Res* 2010;70:3136–9. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-4100.

Veronese A, Lupini L, Consiglio J, Visone R, Ferracin M, Fornari F, Zanesi N, Alder H, D'Elia G, Gramantieri L, Bolondi L, Lanza G, Querzoli P, Angioni A, Croce CM, Negrini M. Oncogenic role of *miR-483-3p* at the *IGF2/483* locus. *Cancer Res* 2010;70:3140–9. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-4456.

Lu W, Zhang G, Zhang R, Flores LG II, Huang Q, Gelovani JG, Li C. Tumor site-specific silencing of *NF- κ B p65* by targeted hollow gold nanosphere-mediated photothermal transfection. *Cancer Res* 2010;70:3177–88. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3379.

Geng H, Rademacher BL, Pittsenbarger J, Huang C-Y, Harvey CT, Lafortune MC, Myrthue A, Garzotto M, Nelson PS, Beer TM, Qian DZ. ID1 enhances docetaxel cytotoxicity in prostate cancer cells through inhibition of p21. *Cancer Res* 2010;70:3239–48. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3186.

Yoo BK, Chen D, Su Z-Z, Gredler R, Yoo J, Shah K, Fisher PB, Sarkar D. Molecular mechanism of chemoresistance by astrocyte elevated gene-1. *Cancer Res* 2010;70:3249–58. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-4009.

Lu ZH, Shvartsman MB, Lee AY, Shao JM, Murray MM, Kladney RD, Fan D, Krajewski S, Chiang GG, Mills GB, Arbeit JM. Mammalian target of rapamycin activator RHEB is frequently overexpressed in human carcinomas and is critical and sufficient for skin epithelial carcinogenesis. *Cancer Res* 2010;70:3287–98. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3467.

Hattermann K, Held-Feindt J, Lucius R, Muerkoster SS, Penfold MET, Schall TJ, Mentlein R. The chemokine receptor CXCR7 is highly expressed in human glioma cells and mediates antiapoptotic effects. *Cancer Res* 2010;70:3299–308. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3642.

Nadiminty N, Lou W, Sun M, Chen J, Yue J, Kung H-J, Evans CP, Zhou Q, Gao AC. Aberrant activation of the androgen receptor by NF- κ B2/p52 in prostate cancer cells. *Cancer Res* 2010;70:3309–19. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3703.

Acu ID, Liu T, Suino-Powell K, Mooney SM, D'Assoro AB, Rowland N, Muotri AR, Correa RG, Niu Y, Kumar R, Salisbury JL. Coordination of centrosome homeostasis and DNA repair is intact in MCF-7 and disrupted in MDA-MB 231 breast cancer cells. *Cancer Res* 2010;70:3320–8. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3800.

McFarlane C, Kelvin AA, de la Vega M, Govender U, Scott CJ, Burrows JF, Johnston JA. The deubiquitinating enzyme USP17 is highly expressed in tumor biopsies, is cell cycle regulated, and is required for G₁-S progression. *Cancer Res* 2010;70:3329–39. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-4152.

Dudka AA, Sweet SMM, Heath JK. Signal transducers and activators of transcription-3 binding to the fibroblast growth factor receptor is activated by receptor amplification. *Cancer Res* 2010;70:3391–401. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3033.

Cho SY, Xu M, Roboz J, Lu M, Mascarenhas J, Hoffman R. The effect of CXCL12 processing on CD34⁺ cell migration in myeloproliferative neoplasms. *Cancer Res* 2010;70:3402–10. Published OnlineFirst April 13, 2010. doi:10.1158/0008-5472.CAN-09-3977.

Published OnlineFirst 05/11/2010.

©2010 American Association for Cancer Research.

doi: 10.1158/0008-5472.CAN-10-1347

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

Type III IFN Interleukin-28 Mediates the Antitumor Efficacy of Oncolytic Virus VSV in Immune-Competent Mouse Models of Cancer

Phonphimon Wongthida, Rosa Maria Diaz, Feorillo Galivo, et al.

Cancer Res 2010;70:4539-4549. Published OnlineFirst May 18, 2010.

Updated version Access the most recent version of this article at:
doi:[10.1158/0008-5472.CAN-09-4658](https://doi.org/10.1158/0008-5472.CAN-09-4658)

Cited articles This article cites 46 articles, 21 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/70/11/4539.full#ref-list-1>

Citing articles This article has been cited by 9 HighWire-hosted articles. Access the articles at:
<http://cancerres.aacrjournals.org/content/70/11/4539.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerres.aacrjournals.org/content/70/11/4539>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.