

MET and KRAS Gene Amplification Mediates Acquired Resistance to MET Tyrosine Kinase Inhibitors

Virna Cepero¹, J. Rafael Sierra¹, Simona Corso¹, Elena Ghiso¹, Laura Casorzo¹, Tim Perera², Paolo Maria Comoglio¹, and Silvia Giordano¹

Abstract

The establishment of the role of MET in human cancer has led to the development of small-molecule inhibitors, many of which are currently in clinical trials. Thus far, nothing is known about their therapeutic efficacy and the possible emergence of resistance to treatment, a problem that has been often observed with other receptor tyrosine kinase (RTK) inhibitors. To predict mechanisms of acquired resistance, we generated resistant cells by treating MET-addicted cells with increasing concentrations of the MET small-molecule inhibitors PHA-665752 or JNJ38877605. Resistant cells displayed *MET* gene amplification, leading to increased expression and constitutive phosphorylation of MET, followed by subsequent amplification and overexpression of wild-type (wt) *KRAS*. Cells harboring *KRAS* amplification progressively lost their MET dependence and acquired *KRAS* dependence. Our results suggest that *MET* and *KRAS* amplification is a general mechanism of resistance to specific MET inhibitors given that similar results were observed with two small inhibitors and in different cell lines of different histotypes. To our knowledge, this is the first report showing that overexpression of wt *KRAS* can overcome the inhibitory effect of a RTK inhibitor. In view of the fact that cellular models of resistance to inhibitors targeting other tyrosine kinases have predicted and corroborated clinical findings, our results provide insights into strategies for preventing and/or overcoming drug resistance. *Cancer Res*; 70(19):7580–90. ©2010 AACR.

Introduction

The frequent alteration of tyrosine kinases in human malignancies led them to be considered as targets for antineoplastic therapies; this resulted in the development of several inhibitors that have shown a strong clinical activity (1–5). The concept of “oncogene addiction” has added further rationale to the use of targeted therapies (6, 7). In general, targeted therapies induce tumor regression in a good percentage of patients. However, almost invariably, responsive patients develop resistance to the treatment and undergo tumor relapse (8). A challenge associated to targeted therapies is, therefore, to predict mechanisms that could cause resistance to the treatment and to find ways to circumvent these hurdles.

Authors' Affiliations: ¹Institute for Cancer Research and Treatment, University of Torino Medical School, Candiolo, Turin, Italy and ²Ortho Biotech Oncology Research and Development, Beerse, Belgium

Note: Supplementary data for this article are available at Cancer Research Online (<http://cancerres.aacrjournals.org/>).

Current address for V. Cepero and J.R. Sierra: University Health Network, Ontario Cancer Institute and Princess Margaret Hospital, Toronto, Ontario, Canada M5G 2M9.

Corresponding Author: Silvia Giordano, Institute for Cancer Research and Treatment, University of Torino Medical School, S.P.142 Km 3.9, 10060 Candiolo, Turin, Italy. Phone: 39-011-9933833; Fax: 39-011-9933225; E-mail: silvia.giordano@unito.it.

doi: 10.1158/0008-5472.CAN-10-0436

©2010 American Association for Cancer Research.

The recent introduction in cancer therapy of several selective tyrosine kinase inhibitors (TKI) has had a dramatic effect in oncology. However, after the first excitement following the initial results, the problem of acquired drug resistance has become more and more important and still represents a crucial limitation. The use of *in vitro* and preclinical models, as well as the evaluation of clinical samples, allowed the identification of several molecular mechanisms responsible for acquired resistance to TKIs (8).

The receptor for hepatocyte growth factor (HGF), encoded by the *MET* gene, which is deregulated in many tumors, is becoming an important candidate for targeted therapies (9, 10). Recent studies by our and other groups have shown that tumor cells displaying *MET* amplification, which results in receptor overexpression and ligand-independent activation, feature oncogenic addiction. In these tumors, inhibition of MET results in impairment of cell growth and in cell death, both *in vitro* and *in vivo* (11–16). In recent times, agents targeting MET—the majority being small-molecule kinase inhibitors—have entered in clinical trials (17–19).

In our work, we have established cell lines resistant to long-term treatment with MET inhibitors and studied the molecular mechanisms involved in the onset and the development of resistance. We show that prolonged treatment with specific MET inhibitors drove amplification, overexpression, and constitutive activation of the receptor. Moreover, we observed that cells progressively amplified

KRAS, resulting in increased expression and activation of wild-type (wt) *KRAS* and in activation of the mitogen-activated protein kinase (MAPK) pathway. We show that amplification of wt *MET* and *KRAS* is found in cells of diverse histotypes, resistant to different inhibitors. Moreover, unexpectedly, we observed that resistance to treatment was reversible and that the alterations leading to resistance were lost after drug withdrawal.

Materials and Methods

Cell lines

GTL16 and EBC-1 cell lines, which harbor *MET* amplification, were cultured in either DMEM or RPMI 1640 supplemented with 10% fetal bovine serum, 100 units/mL penicillin, and 100 µg/mL streptomycin. GTL16 cell line is a clonal cell line established in our lab (as described in refs. 20, 21, derived from the poorly differentiated gastric carcinoma cell line MKN45). After cloning, multiple vials of GTL16 cells were frozen; on thawing, cells were kept in culture for not more than 20 passages. No further characterization of this cell line has been performed, except for the analysis of the karyotype. EBC-1 cell line, derived from a metastatic skin tumor of a patient with lung squamous cell carcinoma, was purchased in 2008 from the Japan Cancer Research Resources Bank. To characterize the cells, the cell bank performed isoenzyme analysis. On arrival, cells were not kept in line for more than 20 passages. Further authentication of the cell line has not been performed by the authors. Resistant cells were generated by exposure to increasing doses of the inhibitor. See Supplementary Materials and Methods for detailed information.

mRNA and genomic DNA analysis

Total mRNA extracts were obtained using Trizol extraction kit (Ambion–Applied Biosystems). RNA (1 µg) was retrotranscribed using M-MLV retrotranscriptase minus H (Promega) in a final volume of 40 µL, according to the manufacturer's protocol. Genomic DNA was extracted using Wizard SV Genomic DNA Purification System (Promega). cDNA (1 µL) or genomic DNA (50 ng) was amplified and analyzed using Power SYBR Green PCR Master Mix (Applied Biosystems) by real-time quantitative PCR using ABI Prism 7900HT (Applied Biosystems). Specific primers were designed using Primer3 web tool (22). Sequences are available on request.

Gene expression and gene copy number fold increase were determined by using the following formula: fold increase = $2^{\Delta - [(C_t \text{ of target gene} - C_t \text{ of control gene})_{\text{resistant}} - (C_t \text{ of target gene} - C_t \text{ of control gene})_{\text{parental}}]}$, where C_t means cycle threshold.

Genomic sequence analysis

Exon sequencing of *MET* (13–20), *EGFR* (18, 19, and 21), *PI3KCA* (9 and 20), *B-RAF* (15), *KRAS* (2–3), and *HRAS* (2–3) was performed by automated sequencing by ABI Prism 3730 (Applied Biosystems).

Western blot analysis

Whole-protein extracts were prepared, quantified using the BCA Protein Assay kit (Pierce), and analyzed using the chemiluminescence ECL kit (Amersham). Primary antibodies anti-phospho-AKT (Ser⁴⁷³), anti-AKT, anti-phospho-p44/42 MAPK, and anti-p44/42 MAPK were purchased from Cell Signaling. Antibodies anti-phospho-MET1234-1235 and anti-phosphotyrosine were from Upstate. Antibodies against *KRAS* (F234) and anti-β-actin (I-9) were from Santa Cruz Biotechnology. Monoclonal antibodies anti-MET DL21 was obtained as described (23). Secondary antibodies were purchased from Amersham.

Pull-down assay for RAS

Pull-down assays were performed using glutathione *S*-transferase (GST)–RAF–CRIB, which specifically binds to the GTP-bound form of RAS, and Western blots were developed using anti-KRAS antibody (F234).

Fluorescence *in situ* hybridization analysis

Fluorescence *in situ* hybridization (FISH) probes for *KRAS* (RP11-295I5) were purchased from Blue Gnone. Probes for the *MET* locus (LSID7S522, spectrum orange), CEP7, and CEP12 (spectrum green) were purchased from Vysis Abbott.

Comparative genomic hybridization analysis

Comparative genomic hybridization (CGH) analyses were performed by NimGenetics. Analyses were performed by comparing genomic DNA from GR150 versus GTL16 wt and GR300 versus GR150 using a 244K CGH array developed by Agilent, Inc.

Biological assays

For growth curve and cell viability assays, cells were seeded in 96-well plastic culture plates (3000 per well) in the presence of the indicated drugs or vehicle (DMSO). Resistant cells were kept in the presence of PHA-665752 (Tocris Bioscience). Cell viability was assessed using a luminescence assay (CellTiter-Glo Luminescent Cell viability Assay, Promega). Anchorage-independent growth in soft agar was performed as described (24), and colony formation was quantified with the Alamar Blue indicator dye (Trek Diagnostic Systems). All the measurements were recorded using a DTX 880 Multimode plate reader (Beckman-Coulter). Small interfering RNA (siRNA) against *MET* [previously described (11)] was electroporated using Amaxa Technology (Instrumentation Laboratory), and short hairpin RNA (shRNA) against *KRAS* sequence (5'-GUGCAAUGAGGGACCAGUA) was cloned into an inducible lentiviral vector as previously described (11). wt *KRAS* cDNA was cloned into the pRRL2 lentiviral vector by following a directional cloning strategy using a *Bam*HI-*Xho*I digestion. MAPK inhibitors were purchased from Sequoia Research Product (sorafenib), Promega (U0126), and Cayman Chemicals (PD0325901).

Xenograft transplantation experiments

All animal procedures were approved by the Ethical Commission of the University of Turin and the Italian Ministry of

Health. GTL16 wt or GR300 cells (3×10^5) were injected s.c. into the right posterior flanks of 6-week-old female CD-1 *nu/nu* mice (Charles River Laboratories). Treatment with JNJ38877605 at half of the maximum tolerated dose or vehicle was initiated after early tumor appearance (approximate volume, 10 mm^3). Tumor volume was monitored every 3 or 4 days for 24 days, calculated as described (25).

Statistical analyses

For experimental data, two-tailed Student's *t* tests were calculated using GraphPad Prism web tool software. A *P* value of <0.05 was considered significant.

Results

Establishment of cell lines resistant to MET inhibitors

GTL16 gastric cancer cells are the prototype of "MET-addicted cells," displaying *MET* amplification (26), overexpression, and constitutive activation (27, 28). Targeting *MET* in GTL16 cells leads to a strong impairment of their viability and growth ability (11, 13–16).

To study possible mechanisms of acquired resistance to *MET* kinase inhibitors, we exposed GTL16 cells to increasing concentrations of the *MET* inhibitor PHA-665752 (29, 30), generating several cell lines resistant to different increasing doses of this inhibitor. Resistant cells were analyzed for their biological and biochemical properties. The growth rate of the resistant cells, which were always kept in the presence of PHA-665752, was only slightly decreased in comparison with wt cells (a representative example is shown in Fig. 1A). Moreover, resistant cells displayed a significant increase in the IC_{50} (half-maximal inhibitory concentration) when compared with wt cells (Fig. 1B), and were able to grow in anchorage-independent conditions in a similar manner as parental cells (Fig. 1C).

To evaluate cross-resistance between PHA-665752 and other *MET* inhibitors, we treated PHA-665752-resistant GTL16 cells with comparable doses of JNJ38877605, a specific *MET* inhibitor of the same class, currently undergoing clinical trials (31), and found that cells showed a similar behavior in terms of cell viability and *MET* status when treated with either compound (Supplementary Fig. S1A and B).

To evaluate if the resistance to PHA-665752 generated *in vitro* was also maintained *in vivo*, we performed xenografts in nude mice. Because PHA-665752 has poor pharmacokinetic properties and low oral bioavailability (17), we daily treated the grafted animals with the orally available JNJ38877605, to which the cells are cross-resistant (Supplementary Fig. S1A and B). As shown in Fig. 1D, mice injected with GTL16 wt cells did not develop tumors in the presence of JNJ38877605 ($P = 0.0015$), whereas resistant cells were able to originate tumors ($P = 0.0222$), which were only slightly smaller than those generated by wt cells in the absence of JNJ38877605 ($P = 0.1659$) or by the same resistant cell growing in the absence of the inhibitor ($P = 0.3347$). These results indicate that resistant cells generated *in vitro* are able to originate tumors *in vivo* in the presence of therapeutic doses of the *MET* inhibitor.

Resistance to MET inhibitors correlates with MET gene amplification

The biochemical analysis of resistant cells showed an increased amount of *MET* protein, which initially paralleled the increase in the concentration of the drug (Fig. 2A, top), thus allowing the maintenance of the basal phosphorylation levels (Fig. 2A, middle). This was concomitant with an increment in *MET* mRNA, evaluated by quantitative reverse transcription-PCR (qRT-PCR; Supplementary Fig. S2A).

Treating the resistant cells with concentrations of inhibitor higher than that used to establish the resistance led to abrogation of *MET* phosphorylation and to the decrease of MAPK and AKT activation (Supplementary Fig. S2B). This is in concordance with previous clinical findings using imatinib, suggesting that, in these conditions, increasing the dose is an effective strategy to overcome resistance (32).

To evaluate if the observed *MET* overexpression was due to gene amplification, a known mechanism of resistance to small inhibitors (8), we performed qRT-PCR on genomic DNA and found an increase in gene copy number in the resistant cells when compared with the parental ones (Fig. 2B). The number of gene copies progressively increased to reach a maximum in cells resistant to 150 nmol/L PHA-665752 (GR150) and started to decrease in cells resistant to higher doses of the drug.

MET overamplification was confirmed by FISH and CGH analysis on PHA-665752-resistant cells. As shown in Supplementary Fig. S2C, CGH analysis of GR150 cells displayed a gain in the genomic area comprising the *MET* locus when compared with wt cells. On the contrary, cells resistant to a higher dose of PHA-665752 (300 nmol/L, GR300) displayed partial loss of the same genomic area. These observations agree with the results obtained by qRT-PCR (see Supplementary Fig. S3 for complete CGH analyses).

In GTL16 wt cells, *MET* is already amplified (11 copies) and the amplicon is localized outside the chromosome 7 (33) on a marker chromosome (M1; ref. 26). Performing FISH analysis, we found that, also in resistant cells, the increased *MET* copies were not located on chromosome 7 (Fig. 2C).

Acquired resistance to MET inhibitors is an adaptive process

To discriminate whether the observed amplification was due to selection of a preexisting subpopulation or if it was acquired during the treatment, we isolated different clones of GTL16 wt cells by limiting dilution and treated two of them (clones 16.1 and 16.2) with increasing doses of the inhibitor to induce resistance (as described in Supplementary Materials and Methods). We found that, in resistant cells obtained from both clones, the number of *MET* copies rose with the increase of the inhibitor concentration, similarly to what was observed in the bulk population of GTL16 wt cells (Supplementary Fig. S4A). This finding reveals that the observed genomic alterations were acquired during the selection.

We also induced resistance in GTL16 wt cells with JNJ38877605, another *MET*-specific kinase inhibitor, using the same procedure described for PHA-665752. We thus obtained cell lines resistant to different concentrations of the




Figure 1. Characterization of PHA-665752-resistant GTL16 cells. Growth ability (A), sensitivity to PHA-665752 (B), and anchorage-independent growth (C) were evaluated by cell viability assays in GTL16 wt cells and resistant to 300 nmol/L PHA-665752 (GR300). Resistant cells were always kept in the presence of the inhibitor. D, mice s.c. injected with either wt or PHA-665752-resistant GTL16 cells were subjected to the indicated treatments. Mean tumor volume \pm SE was evaluated at the indicated times. Vehic, vehicle; 1/2 MTD, half maximum tolerated dose; n, number of tumors per experimental group. *, $P < 0.05$.

drug. Analysis of JNJ38877605-resistant cells revealed an increase in *MET* copy number (Fig. 2D), suggesting that *MET* amplification is not a specific response to PHA-665752 treatment but a more general mechanism of resistance to selective MET TKIs.

A second hit is responsible for resistance to high doses of MET inhibitors

As previously mentioned, MET phosphorylation was maintained up to 150 nmol/L PHA-665752. In cells resistant to higher doses, the level of MET and its phosphorylation started to decrease (Fig. 2A). Moreover, after silencing MET in GR300 cells using siRNAs, we observed that cell viability was almost unaffected compared with what observed in wt and GR150 cells, which are still dependent on MET signaling (Fig. 3A; Supplementary Fig. S2D; $P < 0.0001$). The cells resistant to higher doses of the drug showed a sustained activation of MAPK [extracellular signal-regulated kinase (ERK) 1/2] even if the levels of MET phosphorylation were low (Figs. 2A and 3B).

All these data suggest that in cells resistant to high concentrations of inhibitor, a new molecular event has taken

place and MET activation is no longer the event mediating resistance. To further investigate this possibility, we first discarded the presence of mutations in the receptor by sequencing all the transmembrane and cytoplasmic portions of *MET* in the parental and derived cell lines. In addition, no mutations were found in the signal transducers that are more frequently aberrantly activated in human tumors and are responsible for increased activation of ERK1/2, such as *EGFR*, *KRAS*, *HRAS*, *B-RAF*, and *PI3KCA* (Supplementary Fig. S5).

To evaluate if other tyrosine kinases were differentially activated in resistant cells, we performed anti-phosphotyrosine blots on proteins immunoprecipitated with anti-phosphotyrosine antibodies, finding no qualitative differences in tyrosine phosphorylation patterns (Supplementary Fig. S4C); this result suggests that it is unlikely that other tyrosine kinases (receptor or cytoplasmic) were significantly activated in resistant cells. In particular, we also looked at the activation of epidermal growth factor receptor (EGFR) family members, known to cross talk with MET, and insulin-like growth factor-I receptor, but we did not find increased phosphorylation in GR300 cells (data not shown).

The analysis of the qRT-PCR and CGH data showed the amplification of the genomic region containing *KRAS* in GR300 cells (Fig. 3C; Supplementary Figs. S3 and S6A). FISH analysis showed that the extra copies of *KRAS* were progressively acquired as double minutes outside chromosome 12 (where *KRAS* is positioned; Fig. 3D).


The levels of expression of *KRAS* in GR300 cells were strongly increased, whereas the levels of *HRAS* were unchanged (Fig. 4A, i; $P < 0.0001$). As shown in Fig. 4A (i and ii), *KRAS* protein progressively increased with increasing doses of the drug. Moreover, pull-down experiments conducted with GST-RAF showed that the augmented expression of *KRAS* was paralleled by its enhanced activation, without substantial changes in the ratio of the active molecule (Fig. 4A, iii). Likewise, we found a progressive increase in the number of *KRAS* copies in resistant GTL16 clones (16.1 and 16.2; Supplementary Fig. S4B). This finding reveals that amplification of both *MET* and *KRAS* was acquired during the treatment. Moreover, also cells resistant to high doses of JNJ38877605 displayed an increase in *KRAS* copy number (Fig. 4B).

To prove that the increase of *KRAS*-mediated signaling was involved in resistance, we silenced *KRAS* expression by

transducing the cells with an inducible (Tet-ON) system containing a short-hairpin RNA against *KRAS* (Supplementary Fig. S6B). Although *KRAS* silencing was almost ineffective in wt cells (Supplementary Fig. S6C), it resulted in decreased ability of resistant cells to grow in anchorage-independent manner in soft agar (Fig. 4C; $P < 0.0001$). As a mirror experiment, GTL16 cells overexpressing exogenous wt *KRAS* developed resistance to PHA-665752 (Fig. 4D).

To further validate our observations, we induced resistance to PHA-665752 in the non-small cell lung carcinoma cell line EBC-1, which displays *MET* amplification and high sensitivity to *MET* inhibitors. Resistant cells displayed an increased expression and phosphorylation of *MET* (Fig. 5A, top; Supplementary Fig. S7A), as a result of a further *MET* amplification (Fig. 5B). Moreover, resistance to increasing doses of inhibitor was coupled to high ERK1/2 activation (Fig. 5A, bottom) sustained by increased levels of *KRAS* (Fig. 5A; Supplementary Fig. S7B) due to *KRAS* amplification (Fig. 5B). *MET* and *KRAS* amplification was confirmed by FISH analyses (Fig. 5C and D).

All these results suggest that amplification of both *MET* and *KRAS* can be a general mechanism of resistance to selective *MET* TKIs.






Figure 3. *KRAS* amplification as a second event of resistance. A, to evaluate cell dependence on MET expression, GTL16 wt and resistant cells were transfected with MET-specific siRNAs (siM) or control siRNAs (siC). *, $P < 0.0001$. Cell viability was evaluated after 4 d. B, activation of ERK1/2 and AKT was evaluated in wt and resistant GTL16 cells. pERK1/2, phospho-ERK1/2; pAKT, phospho-AKT. C, evaluation of *KRAS* amplification by qRT-PCR performed on genomic DNA in different cell lines resistant to PHA-665752. D, FISH analysis displays *KRAS* (red) amplification in resistant cells. Green, chromosome 12 centromere (CEP12).

Acquired resistance to MET inhibitors retains dependence on the MAPK pathway

Because resistant cells display a strong activation of ERK1/2, we wondered if they were still responsive to MAPK inhibitors. Therefore, wt and resistant cells were treated with the specific MAPK inhibitors U0126 and PD0325901, or sorafenib (initially identified as a RAF inhibitor and now used in clinics as a multitarget inhibitor), and their ability to grow in either anchorage-dependent or anchorage-independent conditions was evaluated. MAPK inhibitors effectively decreased the anchorage-independent growth ability of resistant cells (Fig. 6A) as well as their viability in anchorage-dependent (adherent) conditions (Supplementary Fig. S8A–C). Western blots to confirm the effects of MAPK inhibitors on MET downstream pathways are shown in Supplementary Fig. S8D. These results suggest that blocking pathways downstream to KRAS

could be an option to overcome the KRAS-mediated resistance to MET inhibitors.

Acquired resistance to MET inhibitors is reversible

To evaluate if resistant cells have acquired a permanent resistant phenotype, we cultured the GR300 cells in the absence of PHA-665752 (GR300-OPHA). Approximately 1 month later, we analyzed these cells and found that they had lost the increased expression of both MET and KRAS (Fig. 6B). The decreased level of protein expression was accompanied by loss of gene copies for both *MET* and *KRAS* (Fig. 6C; $P < 0.0001$), reaching a number of copies similar to that of the wt cells. Likewise, clones of GR300 obtained by single-cell dilution that underwent withdrawal of the inhibitor displayed a significant decrease in *KRAS* copies (Supplementary Fig. S9; $P < 0.0001$). To show that cells kept in the absence

of PHA-665752 had lost their resistance to the inhibitor, we treated them again with the drug. As shown in Fig. 6D, the cells were no longer able to grow in the presence of the same doses of drug to which they had been previously resistant, thus behaving as wt cells. A similar behavior was also observed in cells explanted from tumors resistant to JNJ38877605 *in vivo*. When tumors were excised and cells were put in culture, in the absence of the inhibitor, they lost resistance to the drug.

Discussion

The tyrosine kinase receptor for HGF, encoded by the *MET* gene, has very recently become a target for molecular therapies. Because little is known thus far on the possible mechanisms of resistance to treatment with MET inhibitors, we generated *in vitro* resistant cells by using a protocol similar to those used to establish cells resistant to other TKIs. We thus treated MET-addicted cells with increasing




Figure 4. KRAS is overexpressed and activated in resistant cells. A, i, expression of KRAS and HRAS, evaluated by qRT-PCR performed on RNA extracted from wt, GR150, and GR300 cell lines. *, $P < 0.0001$. ii, Western blot analysis of KRAS expression in resistant cell lines shows an increment of KRAS protein expression correlated with an increase of active KRAS, as shown by the RAF-GST pull-down assay (iii). B, evaluation of KRAS amplification by qRT-PCR performed on genomic DNA in different JNJ38877605-resistant cells. C, GR300 cells were transduced with inducible (Tet-ON) lentiviral vectors containing a short hairpin directed against KRAS (shKRAS) or a control shRNA (shC). Cell ability to survive in anchorage-independent manner was evaluated on induction (doxycycline) of shRNA expression. KRAS silencing severely impaired cell ability to grow in anchorage-independent manner. *, $P < 0.0001$. D, exogenous expression of wt KRAS induced the biological resistance to the MET inhibitor on GTL16 cells when transduced with a lentiviral vector containing wt KRAS exposed to different doses of PHA-665752; expression of KRAS was confirmed by Western blot.




Figure 5. PHA-665752-resistant EBC-1 cells display *MET* and *KRAS* amplification. A, Western blot analysis of *MET* and *KRAS* expression, and activation of *MET*, *ERK1/2*, and *AKT* in EBC-1-resistant cells. B, qRT-PCR analyses showed an increase of *MET* and *KRAS* gene copy number. FISH analysis showed increased copy number of (C) *MET* and (D) *KRAS* (*MET* locus, red; chromosome 7, green; *KRAS*, red; chromosome 12, green). ER: resistant EBC-1, followed by the dose to which they are resistant (in nmol/L).

concentrations of the inhibitor in a stepwise manner, generating cell lines resistant to increasing concentrations of the drug. To investigate the mechanisms responsible for resistance, at first, we ruled out the presence of newly occurring mutations that could potentially interfere with the interaction between the receptor and the drug. Instead, we found that resistant cells show a progressive increase in the amount of *MET* protein that, even in the presence of sustained concentrations of drug, allows the maintenance of a level of tyrosine phosphorylation (and hence pathway activation) comparable with that observed in parental un-

treated cells. By silencing *MET* expression in these resistant cells, we observed a strong impairment of their growth, indicating that these cells were still addicted to *MET*. We found that this increased protein expression was due to *MET* gene amplification, leading to a total copy number of ~60. A similar mechanism of resistance has been observed in the case of other tyrosine kinases, such as *Bcr-Abl* in patients affected by chronic myelogenous leukemia, where the amplification of the *Bcr-Abl* gene is responsible for loss of response to imatinib, and the increase in the dose of the drug can confer again responsiveness to treatment (34).

In our models, FISH analysis showed that the acquired copies of *MET* are not located on chromosome 7 (where the *MET* gene is positioned; ref. 33) but on a marker chromosome (26). This suggests a mechanism of progressive acquisition of additional *MET* copies as a consequence of asymmetric partitioning of the marker chromosome at mitosis. It is likely that cells gaining *MET* extra copies have a selective advantage under the selective pressure of the drug. The observed increase of gene copies is not due to selection of a preexisting population of cells, but rather to an adaptive process, because the *MET* gene underwent amplification also in a clonal population (generated by limiting dilution cloning of wt GTL16 cells) exposed to the drug.

It is puzzling to note that the most frequent genetic alteration of *MET* in human tumors is gene amplification, rather than mutation (35). This behavior is quite different from

that of other tyrosine kinases, such as EGFR or c-Kit, which are more frequently mutated in cancer. Interestingly, *MET* amplification takes place also in tumors treated with inhibitors of other receptor tyrosine kinases (RTK) and mediates resistance to treatment (36–38). An example is the case of some lung cancers treated with gefitinib, where *MET* amplification and overexpression leads to ERBB3 transphosphorylation, thus inducing resistance to inhibitors of the EGFR family (39). One possible explanation for *MET* being frequently amplified relies on the fact that the *MET* gene is located within a known chromosomal common fragile site, FRA7G (33). Hellman and colleagues have shown that conditions that interfere with DNA replication (called “replication stress”) induce a perturbation of chromatin organization of this region, predisposing it to breakage (40). In their work, they hypothesize that this replication stress exerts a selective




Figure 6. PHA-665752-resistant cells are sensitive to multitarget and MAPK inhibitors, and acquired *MET* and *KRAS* amplification is reversible on removal of the inhibitor. A, anchorage-independent growth in soft agar of parental GTL16 and GR300 cells is severely impaired in the presence of U0126, PD0325901, or sorafenib. *, $P < 0.0001$. GR300 cells kept in the absence of PHA-665752 (GR300-0PHA) display *MET* and *KRAS* protein levels (B) and gene copy number (C) similar to those observed in the parental cell line. D, sensitivity to PHA-665752 was evaluated by cell viability assays for GTL16 wt, GR300, and GR300-0PHA cells.

pressure for gene amplification that disappears once the oncogene has attained sufficient amplification. The increased number of *MET* copies, counteracting the activity of the anti-MET drug, rescues cell proliferation.

Interestingly, the number of *MET* copies increased with the dose of the inhibitor up to a threshold, after which it started to decrease. This decrease was accompanied by a reduction of the amount of the MET protein and of its total phosphorylation. At this point, a second unrelated event took place in the cells, which had become independent from MET expression (shown by the fact that MET silencing did not severely impair their viability). We have identified this second event in the progressive amplification of *KRAS*, driving receptor-independent MAPK activation. Amplification of *KRAS* led to an increased expression and activation of the KRAS protein. Moreover, parental cells overexpressing exogenous wt KRAS were capable to overcome the effects of MET inhibitors. In the same line, previous works published by our group showed that overexpression of constitutively active KRAS, BRAF, or AKT could induce the onset of biological resistance to MET inhibitors (41).

The impaired viability of resistant cells on KRAS silencing indicates their dependence on KRAS. On the other hand, parental cells were unaffected by KRAS silencing, likely due to redundant roles of HRAS and KRAS in these cells. This phenomenon, known as “oncogene switch,” has already been reported as a consequence of inhibition of different tyrosine kinases, but it usually involves the activation of other kinases, driving parallel signaling pathways (42). In human tumors, KRAS is almost invariably activated as a consequence of point mutations, with only a few cases where *KRAS* gene amplification has been described (43, 44). It is interesting to note that, as for *MET*, also the *KRAS* gene is located inside a chromosomal common fragile site, on chromosome 12 (45), and thus, it is likely that the same mechanisms responsible for *MET* amplification can account also for this second event.

Because specific anti-RAS drugs are not available, we tested the ability of compounds acting downstream RAS (such as U0126, PD325901, and sorafenib) to impair cell viability. We observed that cells resistant to MET inhibitors that underwent *KRAS* amplification are indeed sensitive to these drugs.

All the molecular events that we have described were not restricted to a single cell line or to a single inhibitor, which indicates that they can represent a more general mechanism to generate cell resistance to small kinase inhibitors against MET.

Finally, we have observed that these cells lost their acquired resistance to MET inhibitors in the absence of the

selective pressure. After culture in the absence of PHA-665752, resistant cells lost the acquired amplification of both *MET* and *KRAS* and once again became sensitive to MET inhibition. One possible explanation for this finding relies on the fact that the amplified oncogenes *MET* and *KRAS* are extrachromosomal in the cells examined. Removal of the inhibitor results in an excess of signal transduction that may induce cellular stress, known to lead to loss of extrachromosomal DNA (46, 47). Cells that underwent loss of *MET* and *KRAS* extra copies have thus an advantage and become the prevalent population in the absence of MET inhibitor. These observations suggest that the mechanism of resistance we observed may be less stable than others already described (such as the appearance of point mutations) and that, possibly, an intermittent therapy, favoring loss of amplified copies, could give better results. Reversibility of the resistant phenotype after drug withdrawal has been previously observed by other groups using other inhibitor for RTKs (48).

This is one of the first preclinical studies highlighting mechanisms of resistance to long-term exposure to selective MET kinase inhibitors (49, 50) and the first report, to our knowledge, showing that amplification of *MET* and *KRAS* genes mediates resistance to MET kinase inhibitors. The prediction of molecular mechanisms of resistance to MET inhibitors may provide insights to strategies aimed at preventing and/or overcoming drug resistance.

Disclosure of Potential Conflicts of Interest

P.M. Comoglio received research support and/or consultation fees from Bayer-Shering, Boehringer-Ingelheim, Johnson & Johnson, and Servier; T. Perera is a full-time employee of Janssen Pharmaceutica and a shareholder of Johnson and Johnson.

Acknowledgments

We thank L. Tamagnone, C. Migliore, A. Petrelli, L. Trusolino, A. Bertotti, and our colleagues for interesting discussions; P. Ferrero, R. Albano, S. Giove, L. Palmas, B. Martinoglio, and R. Porporato for excellent technical assistance; and the staff of the Center for Comparative Oncology for animal care.

Grant Support

Associazione Italiana per la Ricerca sul Cancro (S. Giordano), Ministero dell'Università e della Ricerca (S. Giordano), Regione Piemonte (S. Giordano and S. Corso), and Janssen Pharmaceutica.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received 02/03/2010; revised 06/28/2010; accepted 07/29/2010; published OnlineFirst 09/14/2010.

References

- Vlahovic G, Crawford J. Activation of tyrosine kinases in cancer. *Oncologist* 2003;8:531–8.
- Druker BJ, Guilhot F, O'Brien SG, et al. Five-year follow-up of patients receiving imatinib for chronic myeloid leukemia. *N Engl J Med* 2006;355:2408–17.
- Halatsch ME, Schmidt U, Behnke-Mursch J, Unterberg A, Wirtz CR. Epidermal growth factor receptor inhibition for the treatment of glioblastoma multiforme and other malignant brain tumours. *Cancer Treat Rev* 2006;32:74–89.
- Hurwitz HI, Dowlati A, Saini S, et al. Phase I trial of pazopanib in patients with advanced cancer. *Clin Cancer Res* 2009;15:4220–7.
- Piccart-Gebhart MJ, Procter M, Leyland-Jones B, et al. Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *N Engl J Med* 2005;353:1659–72.

6. Weinstein IB. Cancer. Addiction to oncogenes—the Achilles heel of cancer. *Science* 2002;297:63–4.
7. Weinstein IB, Joe A. Oncogene addiction. *Cancer Res* 2008;68:3077–80, discussion 80.
8. Engelman JA, Settleman J. Acquired resistance to tyrosine kinase inhibitors during cancer therapy. *Curr Opin Genet Dev* 2008;18:73–9.
9. Olivero M, Valente G, Bardelli A, et al. Novel mutation in the ATP-binding site of the MET oncogene tyrosine kinase in a HPRCC family. *Int J Cancer* 1999;82:640–3.
10. Schmidt L, Junker K, Nakaigawa N, et al. Novel mutations of the MET proto-oncogene in papillary renal carcinomas. *Oncogene* 1999;18:2343–50.
11. Corso S, Migliore C, Ghiso E, De Rosa G, Comoglio PM, Giordano S. Silencing the MET oncogene leads to regression of experimental tumors and metastases. *Oncogene* 2008;27:684–93.
12. Lutterbach B, Zeng Q, Davis LJ, et al. Lung cancer cell lines harboring MET gene amplification are dependent on Met for growth and survival. *Cancer Res* 2007;67:2081–8.
13. McDermott U, Sharma SV, Dowell L, et al. Identification of genotype-correlated sensitivity to selective kinase inhibitors by using high-throughput tumor cell line profiling. *Proc Natl Acad Sci U S A* 2007;104:19936–41.
14. Petrelli A, Circo P, Granziero L, et al. Ab-induced ectodomain shedding mediates hepatocyte growth factor receptor down-regulation and hampers biological activity. *Proc Natl Acad Sci U S A* 2006;103:5090–5.
15. Smolen GA, Sordella R, Muir B, et al. Amplification of MET may identify a subset of cancers with extreme sensitivity to the selective tyrosine kinase inhibitor PHA-665752. *Proc Natl Acad Sci U S A* 2006;103:2316–21.
16. Zou HY, Li Q, Lee JH, et al. An orally available small-molecule inhibitor of c-Met, PF-2341066, exhibits cytoreductive antitumor efficacy through antiproliferative and antiangiogenic mechanisms. *Cancer Res* 2007;67:4408–17.
17. Comoglio PM, Giordano S, Trusolino L. Drug development of MET inhibitors: targeting oncogene addiction and expedience. *Nat Rev Drug Discov* 2008;7:504–16.
18. Eder JP, Vande Woude GF, Boerner SA, LoRusso PM. Novel therapeutic inhibitors of the c-Met signaling pathway in cancer. *Clin Cancer Res* 2009;15:2207–14.
19. Toschi L, Janne PA. Single-agent and combination therapeutic strategies to inhibit hepatocyte growth factor/MET signaling in cancer. *Clin Cancer Res* 2008;14:5941–6.
20. Giordano S, Di Renzo MF, Ferracini R, Chiado-Piat L, Comoglio PM. p145, a protein with associated tyrosine kinase activity in a human gastric carcinoma cell line. *Mol Cell Biol* 1988;8:3510–7.
21. Motoyama T, Hojo H, Suzuki T, Oboshi S. Evaluation of the regrowth assay method as an *in vitro* drug sensitivity test and its application to cultured human gastric cancer cell lines. *Acta Med Biol* 1984;27:49–52.
22. Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers. *Methods Mol Biol* 2000;132:365–86.
23. Prat M, Crepaldi T, Pennacchietti S, Bussolino F, Comoglio PM. Agonistic monoclonal antibodies against the Met receptor dissect the biological responses to HGF. *J Cell Sci* 1998;111:237–47.
24. Giordano S, Corso S, Conrotto P, et al. The semaphorin 4D receptor controls invasive growth by coupling with Met. *Nat Cell Biol* 2002;4:720–4.
25. Sierra JR, Corso S, Caione L, et al. Tumor angiogenesis and progression are enhanced by Sema4D produced by tumor-associated macrophages. *J Exp Med* 2008;205:1673–85.
26. Rege-Cambrin G, Scaravaglio P, Carozzi F, et al. Karyotypic analysis of gastric carcinoma cell lines carrying an amplified c-met oncogene. *Cancer Genet Cytogenet* 1992;64:170–3.
27. Bergstrom JD, Hermansson A, Diaz de Stahl T, Heldin NE. Non-autocrine, constitutive activation of Met in human anaplastic thyroid carcinoma cells in culture. *Br J Cancer* 1999;80:650–6.
28. Ponzetto C, Giordano S, Peverali F, et al. c-met is amplified but not mutated in a cell line with an activated met tyrosine kinase. *Oncogene* 1991;6:553–9.
29. Christensen JG, Schreck R, Burrows J, et al. A selective small molecule inhibitor of c-Met kinase inhibits c-Met-dependent phenotypes *in vitro* and exhibits cytoreductive antitumor activity *in vivo*. *Cancer Res* 2003;63:7345–55.
30. Puri N, Khrantsov A, Ahmed S, et al. A selective small molecule inhibitor of c-Met, PHA665752, inhibits tumorigenicity and angiogenesis in mouse lung cancer xenografts. *Cancer Res* 2007;67:3529–34.
31. Perera T. Selective inhibition of Met kinase: JNJ38877605. Spring CTEP Early Drug Development Meeting, Rockville, Massachusetts; 2007.
32. Hochhaus A, La Rosee P. Imatinib therapy in chronic myelogenous leukemia: strategies to avoid and overcome resistance. *Leukemia* 2004;18:1321–31.
33. Miller CT, Lin L, Casper AM, et al. Genomic amplification of MET with boundaries within fragile site FRA7G and upregulation of MET pathways in esophageal adenocarcinoma. *Oncogene* 2006;25:409–18.
34. Beau-Faller M, Ruppert AM, Voegeli AC, et al. MET gene copy number in non-small cell lung cancer: molecular analysis in a targeted tyrosine kinase inhibitor naive cohort. *J Thorac Oncol* 2008;3:331–9.
35. Benvenuti S, Comoglio PM. The MET receptor tyrosine kinase in invasion and metastasis. *J Cell Physiol* 2007;213:316–25.
36. Cappuzzo F, Janne PA, Skokan M, et al. MET increased gene copy number and primary resistance to gefitinib therapy in non-small-cell lung cancer patients. *Ann Oncol* 2009;20:298–304.
37. Puri N, Salgia R. Synergism of EGFR and c-Met pathways, cross-talk and inhibition, in non-small cell lung cancer. *J Carcinog* 2008;7:9.
38. Yano S, Wang W, Li Q, et al. Hepatocyte growth factor induces gefitinib resistance of lung adenocarcinoma with epidermal growth factor receptor-activating mutations. *Cancer Res* 2008;68:9479–87.
39. Engelman JA, Zejnullahu K, Mitsudomi T, et al. MET amplification leads to gefitinib resistance in lung cancer by activating ERBB3 signaling. *Science* 2007;316:1039–43.
40. Hellman A, Zlotorynski E, Scherer SW, et al. A role for common fragile site induction in amplification of human oncogenes. *Cancer Cell* 2002;1:89–97.
41. Bertotti A, Burbridge MF, Gastaldi S, et al. Only a subset of Met-activated pathways are required to sustain oncogene addiction. *Sci Signal* 2009;2:ra80.
42. Pillay V, Allaf L, Wilding AL, et al. The plasticity of oncogene addiction: implications for targeted therapies directed to receptor tyrosine kinases. *Neoplasia* 2009;11:448–58.
43. Hoa M, Davis SL, Ames SJ, Spanjaard RA. Amplification of wild-type K-ras promotes growth of head and neck squamous cell carcinoma. *Cancer Res* 2002;62:7154–6.
44. Mita H, Toyota M, Aoki F, et al. A novel method, digital genome scanning detects KRAS gene amplification in gastric cancers: involvement of overexpressed wild-type KRAS in downstream signaling and cancer cell growth. *BMC Cancer* 2009;9:198.
45. Re A, Cora D, Puliti AM, Caselle M, Sbrana I. Correlated fragile site expression allows the identification of candidate fragile genes involved in immunity and associated with carcinogenesis. *BMC Bioinformatics* 2006;7:413.
46. Durkin SG, Ragland RL, Art MF, Mulle JG, Warren ST, Glover TW. Replication stress induces tumor-like microdeletions in FHIT/FRA3B. *Proc Natl Acad Sci U S A* 2008;105:246–51.
47. Tsantoulis PK, Kotsinas A, Sfrikakis PP, et al. Oncogene-induced replication stress preferentially targets common fragile sites in preneoplastic lesions. A genome-wide study. *Oncogene* 2008;27:3256–64.
48. Sharma SV, Lee DY, Li B, et al. A chromatin-mediated reversible drug-tolerant state in cancer cell subpopulations. *Cell* 2010;141:69–80.
49. McDermott U, Pusapati RV, Christensen JG, Gray NS, Settleman J. Acquired resistance of non-small cell lung cancer cells to MET kinase inhibition is mediated by a switch to epidermal growth factor receptor dependency. *Cancer Res* 2010;70:1625–34.
50. Corso S, Ghiso E, Cepero V, et al. Activation of HER family members in gastric carcinoma cells mediates resistance to MET inhibition. *Mol Cancer* 2010;9:121.

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

***MET* and *KRAS* Gene Amplification Mediates Acquired Resistance to *MET* Tyrosine Kinase Inhibitors**

Virna Cepero, J. Rafael Sierra, Simona Corso, et al.

Cancer Res 2010;70:7580-7590. Published OnlineFirst September 14, 2010.

Updated version	Access the most recent version of this article at: doi: 10.1158/0008-5472.CAN-10-0436
Supplementary Material	Access the most recent supplemental material at: http://cancerres.aacrjournals.org/content/suppl/2010/09/13/0008-5472.CAN-10-0436.DC1

Cited articles	This article cites 49 articles, 22 of which you can access for free at: http://cancerres.aacrjournals.org/content/70/19/7580.full#ref-list-1
-----------------------	--

Citing articles	This article has been cited by 25 HighWire-hosted articles. Access the articles at: http://cancerres.aacrjournals.org/content/70/19/7580.full#related-urls
------------------------	--

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
----------------------	--

Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
-----------------------------------	--

Permissions	To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org .
--------------------	---