

# Long Intergenic Noncoding RNAs: New Links in Cancer Progression

Miao-Chih Tsai, Robert C. Spitale, and Howard Y. Chang

## Abstract

The process of cancer metastasis involves a series of sequential and complex steps. Here we give a perspective on recent results regarding noncoding transcription in cancer progression, focusing on the emerging role of long intergenic noncoding RNAs (lincRNAs). LincRNAs target chromatin modification complexes or RNA-binding proteins to alter gene expression programs. Similarly to miRNAs, lincRNAs exhibit distinct gene expression patterns in primary tumors and metastases. We discuss how lincRNAs can be used for cancer diagnosis and prognosis and serve as potential therapeutic targets. *Cancer Res*; 71(1); 3–7. ©2011 AACR.

## Introduction

Most deaths from cancer are due to metastasis (1), and the main barrier to the treatment of metastases is the biological heterogeneity of cancer cells in the primary neoplasm and in metastases. A full understanding of the gene regulation network during this transition is essential but still far beyond completion. Intensive investigations over the last few decades have focused on the role of protein-coding genes in the pathogenesis of cancer. The human genome contains only 20,000 protein-coding genes, representing <2% of the total genome, whereas a substantial fraction of the human genome can be transcribed, yielding many short or long noncoding RNAs (ncRNAs) with limited protein-coding capacity (2). Are ncRNAs the missing piece of the cancer jigsaw puzzle? One of the emerging themes in the study of noncoding transcripts is microRNAs (miRNAs), a class of small regulatory RNAs that mediate posttranscriptional silencing of specific target mRNAs (3). The identification of the miRNA *miR-34* as the direct target of the tumor suppressor gene *p53* (4) revealed for the first time that ncRNAs function in this crucial tumor suppressor pathway. Numerous miRNA expression profiling and functional studies have also associated miRNAs with cancer progression, diagnosis, prognosis, and treatment (5, 6). Therefore, the interplay between proteins and ncRNAs is an important topic in the field of cancer biology, and ncRNAs may be the missing links in well-known oncogenic and tumor suppressor networks. Long intergenic ncRNAs (lincRNAs), which range in size from several hundred to tens of thousands of bases, comprise another class of newly discovered ncRNAs.

Here we provide an update and perspective on recent advances made in understanding lincRNA mechanisms in cancer progression.

## LincRNA Expression and Cancer

Although >3,000 human lincRNAs have been identified, less than 1% of these have been characterized (7–9). Recent studies showed that lincRNAs are exquisitely regulated during development and in response to diverse signaling cues (8) and can be misexpressed in solid tumors and leukemias (10). Numerous *HOX* lincRNAs were found to be differentially expressed between primary breast carcinomas and distant metastases (11), and many *p53*-dependent lincRNAs were identified in response to DNA damage (12). The finding that several lincRNAs can control transcriptional alteration implies that the difference in lincRNA profiling between normal and cancer cells is not merely the secondary effect of cancer transformation, and that lincRNAs are strongly associated with cancer progression (13). Thus, the differential expression of lincRNAs may be profiled to aid in cancer diagnosis and prognosis and in the selection of potential therapeutics.

Although lincRNAs may have an impact on various human diseases (10, 13, 14), the basis of their molecular mechanisms is still largely unknown. Several lincRNAs can control gene expression by directly recruiting histone-modifying enzymes to chromatin (9). Chromatin modification and DNA methylation are key epigenetic events that are fundamentally disturbed during the development of cancer. Epigenetic alterations are potentially stable and heritable and can occur at a much greater rate than DNA mutations in somatic cells (15). Dysregulated lincRNAs may affect epigenetic information and provide a cellular growth advantage, in that their selection may result in the progressive and uncontrolled growth of a tumor. One such lincRNA, HOTAIR, is highly expressed in breast cancer metastases and in primary tumors predisposed to future metastases. Of importance, enforced expression of HOTAIR can target polycomb repressive complex 2 (PRC2; comprised of histone H3 lysine 27 methylase EZH2, SUZ12,

**Authors' Affiliation:** Howard Hughes Medical Institute and Program in Epithelial Biology, Stanford University School of Medicine, Stanford, California

**Corresponding Author:** Howard Y. Chang, 269 Campus Drive, CCSR 2155, Stanford, CA 94305. Phone: 650-736-0306; Fax: 650-736-0306; E-mail: howchang@stanford.edu

**doi:** 10.1158/0008-5472.CAN-10-2483

©2011 American Association for Cancer Research.

and EED) genome-wide to alter H3K27 methylation and gene expression patterns, resulting in increased cancer invasiveness and metastasis *in vivo*. Loss of HOTAIR or PRC2 components inhibits cancer invasiveness, indicating potentially direct roles for lincRNAs in modulating cancer progression (11). HOTAIR was the first lincRNA to be identified as functioning in *trans* (16); however, other antisense lincRNAs are known to silence genes in *cis*. For example, ANRIL is antisense transcribed from the tumor suppressor genes *CDKN2B* and *CDKN2A*, where ANRIL interacts with CBX7 (a subunit of PRC1), resulting in heterochromatin formation and gene silencing (17–19).

LincRNAs may carry out many of their functions by acting as modular scaffolds for protein–chromatin interactions (20, 21). Tsai and colleagues (21) recently discovered that HOTAIR bridges together the PRC2 complex and the LSD1 H3K4 demethylase complex and recruits both complexes to target genes to coordinately alter several histone modifications and enforce gene silencing. As another example, TLS protein can be allosterically modified by interacting with ncRNA<sub>CCND1</sub>, transcribing from the 5' regulatory regions of *CCND1* in response to DNA damage signals. The conformation change of TLS induces gene-specific TLS–CBP/p300 interactions, which in turn inhibit *CCND1* transcription (22). Another lincRNA, MALAT-1, which is highly expressed in many different tumors, can interact with SR splicing factors and modulates their distribution to nuclear speckles. MALAT1 then regulates alternative splicing by controlling the functional levels of SR splicing factors (23). LincRNAs can also affect global gene changes in response to signals to classic transcription factors. For example, *p53* induces lincRNA-p21, which in turn represses numerous genes globally by recruiting the repressor protein hnRNP-K (12).

Dozens of lincRNAs are differentially expressed in cancers with different clinical courses. In multivariate analyses with clinical and pathologic risk factors, HOTAIR RNA can serve as a significant and independent predictor of eventual metastasis and death (11). Because risk stratification by mRNA or miRNA profiling typically requires dozens if not hundreds of RNA species, the ability to use a single species of lincRNA (HOTAIR) represents a promising approach. These examples suggest that HOTAIR and other lincRNAs may be used as a novel prognostic tool in cancer management.

### Mutations and Alternative RNA Structural Variation

The misexpression of lincRNAs in cancer naturally raises the question as to how structural variations in lincRNA genes, either germline or somatic, may contribute to cancer predisposition. To date, analyses of DNA copy number variations and rearrangements have focused on protein coding genes, although many noncoding sequences are included in recurrent structural aberrations of cancer genomes. Similarly, genome-wide association studies of cancer susceptibility have mainly identified single nucleotide polymorphisms in non-coding portions of the genome, some of which may be


transcribed (24). For instance, a region upstream of the 9p21 locus, encoding the cyclin-dependent kinase inhibitors *CDKN2B* and *CDKN2A* and *p53* activator *ARF*, is associated with cancer as well as cardiovascular disease and diabetes, and this region is transcribed to produce ANRIL, a long noncoding transcript that is implicated in controlling chromatin modification of the locus (17–19). Compared with mutations in protein coding genes, where certain single nucleotide mutations (such as a premature stop codon or frame shift) can completely abrogate protein function, structural variations in lincRNAs may have more subtle effects, which makes it more difficult to verify them experimentally (25). Better annotation of lincRNA expression patterns and primary structures should improve the detection and interpretation of genome aberrations that affect lincRNAs. Moreover, because known oncogenes and tumor suppressor genes are often dysregulated by multiple mechanisms, lincRNAs that are misexpressed in cancers should be the starting point for possible structural alterations in lincRNA genes.

### Targeting Small ncRNAs Associated with Cancer

A major goal in the development of cancer therapeutics is to identify molecular targets that are specific for cancer cells (26). The differential expression and biological importance of specific ncRNAs in cancer suggest that ncRNAs may be useful targets (27). Here we briefly discuss efforts to target miRNAs and then speculate on the potential of targeting long ncRNAs for cancer treatment (Fig. 1).

miRNAs have been effectively targeted in preclinical models by means of small molecules and exogenously introduced complementary RNA sequences (28). High-throughput screens, both *in vitro* and *in vivo*, have identified small molecules that affect several steps in miRNA biogenesis, including pre-miRNA processing, or interact directly with the miRNA, thus blocking miRNA hybridization with its endogenous 3'-UTR sequences (28). For example, a small-molecule inhibitor of *miR-122* can selectively induce apoptosis in liver cancer cells (29). Screens involving small molecules often are initiated with little prior chemical knowledge of a successful binder. Because every RNA sequence is different, it is difficult to generalize the chemical functionality of anti-miRNA molecules. Further, although studies such as those mentioned above are leading to promising compounds, the modest binding affinity of these compounds [often in the micromolar range (30)] suggests a need for improvement. A more targeted approach that exploits the miRNA sequence directly may be more advantageous.

Antagomirs are synthetic RNA molecules that are designed to directly hybridize with miRNAs, thus potentially limiting the availability of the miRNA for Argonaute loading and 3'-UTR hybridization. Antagomirs are chemically modified to prevent premature degradation by RNases, thereby increasing their half-life *in vivo* (31). Krützfeldt and colleagues (32) were the first to show that antagomirs can inhibit specific miRNA function in living mice. Subsequently, Ma and colleagues (33)



**Figure 1.** Targeting cancer-specific ncRNAs. Small molecules and complementary antagomir sequences have been identified that bind to miRNAs, disrupting base pairing. We speculate that targeting the lincRNA structure could inhibit binding to chromatin-modifying complexes without disrupting the binding of different RNAs.

showed the utility of antagomirs in preventing the onset of metastasis. The authors reported that systematic treatment of tumor-bearing mice with antagomirs targeting *miR-10b* increased the expression of its target, *HOXD10*, a tumor suppressor gene that is downregulated in metastasis. In another study (34), intravenous administration of the antagomir did not reduce primary mammary tumor growth, but it did strikingly suppress the formation of lung metastases in a sequence-specific manner. These studies preliminarily suggest that antagomirs may be powerful molecules that can disrupt the miRNA pathway and may be potential chemotherapeutics. Although we are just beginning to understand the potential of lincRNAs as therapeutic targets, it is conceivable that the avenues that have been explored for other ncRNAs can be used to target lincRNA-protein interactions. In the next section, we highlight some of the potential avenues that can be tailored to lincRNAs and their protein partners.

### Potential New Cancer Therapy: lincRNAs

Although efforts to target cancer-specific miRNAs have proven successful, designing molecules to inhibit long ncRNAs presents different opportunities and challenges. The fact that lincRNAs can be depleted by siRNAs provides a straightforward approach to inhibit the function of lincRNAs implicated in cancer. Gupta and colleagues (11) showed that depletion of HOTAIR inhibited the matrix invasiveness of a breast cancer

cell line. In practice, we and others (21) have found that to achieve a successful knockdown of lincRNAs, one must screen more siRNAs compared with mRNAs, possibly because of the extensive secondary structures in lincRNAs.

Structural insights into riboswitches, the ribosome, RNase P, and catalytic RNAs are beginning to be exploited for the design of specific small-molecule inhibitors that can target RNA and ribonucleoprotein (RNP) complexes (35–37). For example, investigators have used the atomic structure of the bacterial ribosome, the most complex RNP structurally characterized to date, to identify functional sites within rRNA for therapeutic targeting by antibiotics (38). These results underscore the importance of structural studies to aid in the design of therapeutic agents against complex RNAs. Detailed structural studies of recently discovered lincRNAs associated with cancer are still forthcoming.

LincRNAs such as HOTAIR serve as a structural scaffold for protein complexes and possess complex RNA structural motifs (21). Because the chromatin complexes that interact with lincRNAs are ubiquitous, targeting the proteins themselves may lead to deleterious phenotypes. However, targeting the RNA-protein interface and focusing on the cancer-specific transcript (e.g., HOTAIR) may provide an inroad to cancer-specific therapeutics. Targeting transcripts the size of lincRNAs may seem like a daunting task; nevertheless, there is a precedent for fragmenting large RNP complexes into more manageable sizes. This strategy has been applied to the design of ligands that can bind expanded rCUG and rCAG repeat RNAs that are expressed in myotonic dystrophy type 1 (DM1) and interact with Muscleblind-like 1 (MBNL1) protein. Ligands identified *in vitro* have also been shown to inhibit the formation of RNA-protein complexes in mouse myoblasts (39). These results, along with other findings, provide strong evidence that examining defined, functionally relevant RNA fragments *in vitro* can lead to the discovery of small-molecule binders, and the information gained can be used in the cellular context to disrupt RNA-protein interactions.

Although one can investigate lincRNAs by applying methods and principles similar to those used to target miRNAs with small molecules, one must use a different strategy for antagomirs. Because the structural content of lincRNAs is expected to be vast, designing antagomirs based on the primary sequence may yield unsuccessful results. However, using a more unbiased approach may yield oligonucleotide sequences that bind to lincRNAs. Methods such as systematic evolution of ligands by exponential enrichment (SELEX) provide an unbiased way to identify high-affinity binders of a selected macromolecule (40). Just recently, SELEX was used to identify RNA sequences that bind to pri-miRNAs (41). pri-miRNAs are larger than their mature products and contain substantial predicted secondary structure. Therefore, a similar approach may yield RNA sequences that can bind to lincRNAs at key lincRNA-protein interfaces, thereby disrupting the interaction. We suggest the term "antagolincs" for this yet to be established class of molecules. Once an RNA sequence is identified through SELEX, the same RNA sequence can be transformed into the more chemically stable construct, in similarity to an

antagomir. The administration of an antagomir against HOTAIR would then lead to competitive inhibition of a chromatin remodeling complex, such as PRC2, by binding to the lincRNA. This RNA–RNA interaction may then normalize the chromatin state to inhibit cancer cell growth and metastasis.

## Conclusions

Just as dysregulation of miRNAs is now recognized as a universal feature of many types of cancer, many lincRNAs that affect cancer initiation, progression, and treatment are surely waiting to be discovered. Cancer-associated lincRNAs may provide new approaches to the diagnosis and treatment of cancer. Systematic identification of lincRNA expression patterns and characterization of lincRNAs and their associated

proteins should pave the way for designer therapeutics that can target lincRNAs for cancer and other disease states.

## Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

## Grant Support

R01-CA118750, National Institutes of Health and California Institute for Regenerative Medicine (RN1-00529-1); American Cancer Society [(RSG 07-084-01-MGO); H.Y. Chang]; T32-AR007422, National Institutes of Health (R.C. Spitale); Susan G. Komen Foundation (M.-C. Tsai).

H. Y. Chang is an Early Career Scientist of the Howard Hughes Medical Institute.

Received July 12, 2010; revised October 14, 2010; accepted October 14, 2010; published online January 3, 2011.

## References

- Fidler IJ. The pathogenesis of cancer metastasis: the 'seed and soil' hypothesis revisited. *Nat Rev Cancer* 2003;3:453–8.
- Birney E, Stamatoyannopoulos JA, Dutta A, Guigo R, Gingeras TR, Margulies EH, et al. Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 2007;447:799–816.
- Fabian MR, Sonenberg N, Filipowicz W. Regulation of mRNA translation and stability by microRNAs. *Annu Rev Biochem* 2010;79:351–79.
- Hermeking H. p53 enters the microRNA world. *Cancer Cell* 2007;12:414–8.
- Lu J, Getz G, Miska EA, Alvarez-Saavedra E, Lamb J, Peck D, et al. MicroRNA expression profiles classify human cancers. *Nature* 2005;435:834–8.
- Ruan K, Fang X, Ouyang G. MicroRNAs: novel regulators in the hallmarks of human cancer. *Cancer Lett* 2009;285:116–26.
- Khalil AM, Guttman M, Huarte M, Garber M, Raj A, Rivea Morales D, et al. Many human large intergenic noncoding RNAs associate with chromatin-modifying complexes and affect gene expression. *Proc Natl Acad Sci USA* 2009;106:11667–72.
- Guttman M, Amit I, Garber M, French C, Lin MF, Feldser D, et al. Chromatin signature reveals over a thousand highly conserved large non-coding RNAs in mammals. *Nature* 2009;458:223–7.
- Ponting CP, Oliver PL, Reik W. Evolution and functions of long noncoding RNAs. *Cell* 2009;136:629–41.
- Calin GA, Liu CG, Ferracin M, Hyslop T, Spizzo R, Sevignani C, et al. Ultraconserved regions encoding ncRNAs are altered in human leukemias and carcinomas. *Cancer Cell* 2007;12:215–29.
- Gupta RA, Shah N, Wang KC, Kim J, Horlings HM, Wong DJ, et al. Long non-coding RNA HOTAIR reprograms chromatin state to promote cancer metastasis. *Nature* 2010;464:1071–6.
- Huarte M, Guttman M, Feldser D, Garber M, Koziol MJ, Kenzelmann-Broz D, et al. A large intergenic noncoding RNA induced by p53 mediates global gene repression in the p53 response. *Cell* 2010;142:409–19.
- Huarte M, Rinn JL. Large non-coding RNAs: missing links in cancer? *Hum Mol Genet* 2010;19:R152–61.
- Yu W, Gius D, Onyango P, Muldoon-Jacobs K, Karp J, Feinberg AP, et al. Epigenetic silencing of tumour suppressor gene p15 by its antisense RNA. *Nature* 2008;451:202–6.
- Chi P, Allis CD, Wang GG. Covalent histone modifications—miswritten, misinterpreted and mis-erased in human cancers. *Nat Rev Cancer* 2010;10:457–69.
- Rinn JL, Kertesz M, Wang JK, Squazzo SL, Xu X, Bruggmann SA, et al. Functional demarcation of active and silent chromatin domains in human HOX loci by noncoding RNAs. *Cell* 2007;129:1311–23.
- Pasmant E, Laurendeau I, Heron D, Vidaud M, Vidaud D, Bieche I. Characterization of a germ-line deletion, including the entire INK4/ARF locus, in a melanoma-neural system tumor family: identification of ANRIL, an antisense noncoding RNA whose expression coclusters with ARF. *Cancer Res* 2007;67:3963–9.
- Gil J, Peters G. Regulation of the INK4b-ARF-INK4a tumour suppressor locus: all for one or one for all. *Nat Rev Mol Cell Biol* 2006;7:667–77.
- Yap KL, Li S, Munoz-Cabello AM, Raguz S, Zeng L, Mujtaba S, et al. Molecular interplay of the noncoding RNA ANRIL and methylated histone H3 lysine 27 by polycomb CBX7 in transcriptional silencing of INK4a. *Mol Cell* 2010;38:662–74.
- Zappulla DC, Cech TR. RNA as a flexible scaffold for proteins: yeast telomerase and beyond. *Cold Spring Harb Symp Quant Biol* 2006;71:217–24.
- Tsai MC, Manor O, Wan Y, Mosammaparast N, Wang JK, Lan F, et al. Long noncoding RNA as modular scaffold of histone modification complexes. *Science* 2010;329:689–93.
- Wang X, Arai S, Song X, Reichart D, Du K, Pascual G, et al. Induced ncRNAs allosterically modify RNA-binding proteins in cis to inhibit transcription. *Nature* 2008;454:126–30.
- Tripathi V, Ellis JD, Shen Z, Song DY, Pan Q, Watt AT, et al. The nuclear-retained noncoding RNA MALAT1 regulates alternative splicing by modulating SR splicing factor phosphorylation. *Mol Cell* 2010;39:925–38.
- Manolio TA, Brooks LD, Collins FSA. HapMap harvest of insights into the genetics of common disease. *J Clin Invest* 2008;118:1590–605.
- Mattick JS. The genetic signatures of noncoding RNAs. *PLoS Genet* 2009;5:e1000459.
- Gewirtz DA, Bristol ML, Yalowich JC. Toxicity issues in cancer drug development. *Curr Opin Investig Drugs* 2010;11:612–4.
- Ryan BM, Robles AI, Harris CC. Genetic variation in microRNA networks: the implications for cancer research. *Nat Rev Cancer* 2010;10:389–402.
- Deiters A. Small molecule modifiers of the microRNA and RNA interference pathway. *AAPS J* 2010;12:51–60.
- Young DD, Connelly CM, Grohmann C, Deiters A. Small molecule modifiers of microRNA miR-122 function for the treatment of hepatitis C virus infection and hepatocellular carcinoma. *J Am Chem Soc* 2010;132:7976–81.
- Gumireddy K, Young DD, Xiong X, Hogenesch JB, Huang Q, Deiters A. Small-molecule inhibitors of microRNA miR-21 function. *Angew Chem Int Ed Engl* 2008;47:7482–4.
- Krützfeldt J, Kuwajima S, Braich R, Rajeev KG, Pena J, Tuschl T, et al. Specificity, duplex degradation and subcellular localization of antagomirs. *Nucleic Acids Res* 2007;35:2885–92.
- Krützfeldt J, Rajewsky N, Braich R, Rajeev KG, Tuschl T, Manoharan M, et al. Silencing of microRNAs in vivo with 'antagomirs'. *Nature* 2005;438:685–9.



33. Ma L, Teruya-Feldstein J, Weinberg RA. Tumour invasion and metastasis initiated by microRNA-10b in breast cancer. *Nature* 2007; 449:682–8.
34. Ma L, Reinhardt F, Pan E, Soutschek J, Bhat B, Marcusson EG, et al. Therapeutic silencing of miR-10b inhibits metastasis in a mouse mammary tumor model. *Nat Biotechnol* 2010;28:341–7.
35. Serganov A, Patel DJ. Ribozymes, riboswitches and beyond: regulation of gene expression without proteins. *Nat Rev Genet* 2007;8:776–90.
36. Kim JN, Blount KF, Puskarz I, Lim J, Link KH, Breaker RR. Design and antimicrobial action of purine analogues that bind guanine riboswitches. *ACS Chem Biol* 2009;4:915–27.
37. Wilson DN. The A-Z of bacterial translation inhibitors. *Crit Rev Biochem Mol Biol* 2009;44:393–433.
38. Wimberly BT. The use of ribosomal crystal structures in antibiotic drug design. *Curr Opin Investig Drugs* 2009;10:750–65.
39. Lee MM, Childs-Disney JL, Pushechnikov A, French JM, Sobczak K, Thornton CA, et al. Controlling the specificity of modularly assembled small molecules for RNA via ligand module spacing: targeting the RNAs that cause myotonic muscular dystrophy. *J Am Chem Soc* 2009;131:17464–72.
40. Djordjevic M. SELEX experiments: new prospects, applications and data analysis in inferring regulatory pathways. *Biomol Eng* 2007;24:179–89.
41. Lunse CE, Michlewski G, Hopp CS, Rentmeister A, Caceres JF, Famulok M, et al. An aptamer targeting the apical-loop domain modulates pri-miRNA processing. *Angew Chem Int Ed Engl* 2010;49: 4674–7.

# Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

## Long Intergenic Noncoding RNAs: New Links in Cancer Progression

Miao-Chih Tsai, Robert C. Spitale and Howard Y. Chang

*Cancer Res* 2011;71:3-7.

**Updated version** Access the most recent version of this article at:  
<http://cancerres.aacrjournals.org/content/71/1/3>

**Cited articles** This article cites 41 articles, 6 of which you can access for free at:  
<http://cancerres.aacrjournals.org/content/71/1/3.full.html#ref-list-1>

**Citing articles** This article has been cited by 40 HighWire-hosted articles. Access the articles at:  
</content/71/1/3.full.html#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, contact the AACR Publications Department at [permissions@aacr.org](mailto:permissions@aacr.org).