

Title:

Effect of depo-medroxyprogesterone acetate on breast cancer risk among women 20-44 years of age

Running title:

DMPA use and breast cancer risk

Authors:

Christopher I. Li,¹ Elisabeth F. Beaber,¹ Mei-Tzu Chen Tang,¹ Peggy L. Porter,^{1,2} Janet R. Daling,¹ Kathleen E. Malone¹

¹ Division of Public Health Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA

² Division of Human Biology, Fred Hutchinson Cancer Research Center, Seattle, WA

Corresponding author:

Christopher I. Li, MD, PhD

Fred Hutchinson Cancer Research Center

1100 Fairview Ave. N., M4-C-308

P.O. Box 19024

Seattle, WA 98109-1024

Telephone: 206.667.7444

Fax: 206.667.5948

E-mail: cili@fhcrc.org

Financial support:

This study was funded by the National Cancer Institute (R01-CA105041, ARRA supplement to R01-CA10541, and T32-CA09168) and the Department of Defense Breast Cancer Research Program (W81XWH-05-1-0482).

Potential conflicts of interest:

None of the authors have any potential conflicts of interest.

PRECIS

Findings suggest that the injectable contraceptive medroxyprogesterone acetate appears to increase risk of breast cancer in young women.

ABSTRACT

Depo-medroxyprogesterone acetate (DMPA) is an injectable contraceptive that contains the same progestin as the menopausal hormone therapy regimen found to increase breast cancer risk among postmenopausal women in the Women's Health Initiative clinical trial. However, few studies have evaluated the relationship between DMPA use and breast cancer risk. Here we conducted a population-based case-control study among 1028 women 20-44 years of age to assess the association between DMPA use and breast cancer risk. Detailed information on DMPA use and other relevant covariates was obtained through structured interviewer administered in-person questionnaires, and unconditional logistic regression was used to evaluate associations between various aspects of DMPA use and breast cancer risk. We found that recent DMPA use for 12 months or longer was associated with a 2.2-fold (95% CI: 1.2-4.2) increased risk of invasive breast cancer. This risk did not vary appreciably by tumor stage, size, hormone receptor expression, or histological subtype. Although breast cancer is rare among young women and the elevated risk of breast cancer associated with DMPA appears to dissipate after discontinuation of use, our findings emphasize the importance of identifying the potential risks associated with specific forms of contraceptives given the number of available alternatives.

INTRODUCTION

Taken together, the results of the Women's Health Initiative (WHI) randomized controlled trials of postmenopausal hormones strongly suggest that progestational agents, and medroxyprogesterone acetate (MPA) in particular, increase a woman's risk of breast cancer. Specifically, MPA in combination with conjugated estrogen was observed to increase breast cancer incidence by 24%,⁽¹⁾ while users of unopposed estrogen hormone therapy (EHT) had a non-statistically significant reduced risk.⁽²⁾ Thus, the progestin component of combined estrogen and progestin menopausal hormone therapy (CHT) appears to play a central role in elevating breast cancer risk.

The injectable contraceptive depo-medroxyprogesterone acetate (DMPA) is another progestin containing preparation that is widely used by women throughout the world. It contains the same progestin that was evaluated in the WHI trial. In 1992 DMPA received FDA approval for use as a contraceptive, and since this time rates of DMPA use have steadily increased in the U.S.⁽³⁾ However, there are limited data on the relationship between DMPA and breast cancer incidence. Results across international case-control studies conducted in Costa Rica, New Zealand, Kenya, Mexico, Thailand, and South Africa are somewhat mixed, with one analysis showing that ever use of DMPA increases breast cancer risk 2.6-fold,⁽⁴⁾ one finding it increases risk 1.2-fold,⁽⁵⁾ and two observing no association between ever use of DMPA and breast cancer risk.^(6, 7) However, the three studies evaluating recency of use all consistently found that current DMPA use was associated with a 1.5 to 1.65-fold increased risk of breast cancer.⁽⁵⁻⁷⁾ Additional studies of the relationship between DMPA use and breast cancer incidence are needed because previous reports have been limited by small

numbers of cases <45 years of age, none evaluated risk according to breast cancer subtype, and the generalizability of these results to other populations such as those in more developed countries is uncertain given differences in breast cancer incidence rates, demographics, and reproductive patterns across women worldwide.

METHODS

We conducted a large population-based case-control study of breast cancer among women 20 to 44 years of age living in the three county Seattle-Puget Sound metropolitan area (King, Pierce, and Snohomish counties) specifically designed to assess the relationship between DMPA use and breast cancer risk.

Cases were women 20 to 44 years old diagnosed with a primary invasive breast cancer between June 2004 and June 2010 with no prior history of *in situ* or invasive breast cancer. Potentially eligible cases residing in King, Pierce, and Snohomish counties were identified through the Cancer Surveillance System, the population-based tumor registry that serves the 13 counties of western Washington state and participates in the Surveillance, Epidemiology, and End Results program of the National Cancer Institute. Since controls were ascertained via random digit dialing of landline home telephone numbers, to be eligible all cases were also required to have a landline home telephone (160 potentially eligible cases without a landline telephone were identified and excluded). Of the 1,359 eligible cases identified, 1,056 (78%) were interviewed. Of those not enrolled (n=303), 82% refused to be interviewed, 10% could not be located, and 8% died before an interview could be conducted. In addition to basic information on breast cancer diagnosis, we obtained information on tumor characteristics from the

cancer registry and from a centralized review of pathology reports. This includes data on estrogen receptor (ER), progesterone receptor (PR), and HER2-neu (HER2) status; and tumor stage, size, and histology.

Using a combination of list-assisted (purchased randomly generated telephone numbers) and Mitosky-Waksberg (telephone numbers randomly generated ourselves using a clustering factor of 5)(8) random digit dialing methodologies, controls from the general population of female residents of King, Pierce, and Snohomish counties were identified. Controls were frequency matched within 5-year age groups to the cases. A 1:1 ratio of controls to cases was employed for participants with reference dates from 2004-2007 and was then switched to 0.7:1 for those with reference dates from 2008-2010 as a result of additional funding that was obtained to increase the number of cases enrolled during the latter years of our data collection effort. A total of 90,488 random telephone numbers were pursued with multiple rounds of contacts attempted as needed. 66,844 numbers were nonworking, business, cellular, paging, dedicated facsimile, or data line numbers. 3,570 numbers were never answered, and thus their residential status could not be determined. Prior studies suggest that only about 20% of such numbers are indeed residential.(9) Of the 20,074 residential or presumed residential numbers, 14,130 were successfully screened for eligibility. Of the remainder, 3,105 were answering machines, 2,351 reached a respondent who refused to answer the screening questions, and for 488 there were language or other communication barriers. Of the 1,489 eligible controls identified, 943 (63%) were interviewed.

For this analysis, 24 controls and 28 cases missing data on use of injectable contraceptives were excluded. Thus, our final analytic sample size consisted of 919 controls and 1,028 cases.

Data Collection

The study protocol was approved by the Fred Hutchinson Cancer Research Center Institutional Review Board, and written informed consent was obtained from all study subjects. Cases and controls were interviewed in-person and asked about their reproductive history, body size, medical history, and family history of cancer. Additionally, detailed histories of all episodes of hormonal contraceptive use, including beginning and ending dates, brand, dose, route of administration, and pattern of use (number of days per month) were obtained. Our questioning was limited to exposures that occurred before each participant's reference date (month and year). The reference date used for each woman with breast cancer was her diagnosis date. As described above, controls were frequency matched to cases on reference year. The reference months assigned to controls reflected the distribution of reference months among the cases.

Statistical Analysis

The primary referent category consisted of women who never used any type of injectable hormonal contraceptive. We also conducted sensitivity analyses considering alternative reference categories. These included defining women who never used any type of hormonal contraceptive as the reference category, and conducting analyses limited to ever users of hormonal contraceptives where the reference category consisted of ever users of a non-injectable hormonal contraceptive. Since even a single

DMPA shot has been shown to result in MPA measurable in serum for as long as 7.5 to 9 months after injection, ever users of DMPA were defined as women who ever received even a single DMPA shot.(10, 11) Given these pharmacokinetics, recent DMPA use was defined as having received one or more DMPA shots within five years of reference date similar to how recency of hormone use has been defined by the Collaborative Group on Hormonal Factors.(12) Duration of use was calculated by attributing three months of “use” (exposure) for each DMPA shot a woman received as the recommended prescribed regimen is shots that are administered every three months. Analyses focus primarily on recent DMPA use for at least 12 months because in pooled data from 54 epidemiological studies conducted world-wide the positive relationship between oral contraceptive use and breast cancer risk was only observed among recent users for at least 12 months duration.(13)

We used unconditional logistic regression to calculate odds ratios (ORs) and their associated 95% confidence intervals (CIs) to compare breast cases to controls.(14) All analyses were conducted using Stata/SE version 11.2 (StataCorp LP, College Station, TX). All models were adjusted for age (five year categories) and reference year (continuous) since controls were matched to cases on these factors and additionally adjusted for first degree family history of breast cancer (no / yes / missing), body mass index one year prior to reference date [$<25.0 \text{ kg/m}^2$ / $(25.0-29.9 \text{ kg/m}^2$ / $\geq 30.0 \text{ kg/m}^2$ / missing (based on WHO categories)], number of full-term pregnancies (0 / 1-3 / ≥ 3 / missing), duration of oral contraceptive (OC) use (never / <5 years / 5-9.9 years / ≥ 10 years / missing), and screening mammography (ever / never). These latter five covariates were selected *a priori* as potential confounders. Other variables

evaluated as potential confounders included: education, income, race/ethnicity (based on self-report), and age at first live birth. None of these potential confounders changed our risk estimates by more than 10% and there was no statistically significant change ($p < 0.05$) in the fit of the model with the addition of any of these potential confounders. Thus none were added to our final statistical models.

RESULTS

Cases and controls had generally similar distributions with respect to age, education, and income (Table 1). Cases were somewhat more likely to be African American and Asian and less likely to be Hispanic white compared to controls. Cases were also more likely to have a first degree family history of breast cancer, to be somewhat leaner, to have used OCs for 10 years or longer, to be nulliparous, to have a younger age at first birth, and to have ever had a screening mammogram.

Among control women, compared to never users of hormonal contraception, ever users of DMPA were somewhat more likely to be younger, African American, Native American, less educated, to not have a first-degree family history of breast cancer, and to be obese, and were somewhat less likely to be Asian/Pacific Islander and nulliparous (Table 2). Compared to never users of hormonal contraception, ever users of hormonal contraception who never used DMPA were somewhat older, more educated, and were less likely to be Asian/Pacific Islander and nulliparous.

In our primary analysis we compared DMPA users to all non-users of DMPA and observed that neither ever use nor recent use of DMPA was associated with breast

cancer risk using multivariate adjusted statistical models (OR=1.2, 95% CI: 0.9-1.6 and OR=1.5, 95% CI: 0.9-2.7, respectively) (Table 3). However, recent users of DMPA for 12 months or longer had a 2.2-fold increased risk of breast cancer (95% CI: 1.2-4.2). There was also some suggestion that age at first use of DMPA influenced risk as women who first used DMPA at age ≥ 35 years had a non-statistically significant 2.0-fold (95% CI: 0.9-4.6) increased risk of breast cancer. Timing of DMPA use in relation to either first or most recent full-term pregnancy was not related to risk.

To further evaluate the relationship between DMPA use and breast cancer risk we considered alternative reference groups. Using a reference group of never users of any type of hormonal contraceptive (included oral contraceptives, contraceptive patches, implants, and hormonal intra-uterine devices) throughout a woman's life, recent use of DMPA for ≥ 12 months was associated with a 2.8-fold (95% CI: 1.3-5.9) increased risk of breast cancer. While this risk estimate was statistically significant, it is important to note that this analysis was constrained to a small numbers of cases (n=105) and controls (n=91) comprising the reference group of never users of any hormonal contraceptive. A second approach restricted the analysis to women who had ever used some type of hormonal contraceptive and used a reference group of women who had ever used a hormonal contraceptive but had never used DMPA. Using this approach, recent use of DMPA for ≥ 12 months was associated with a 2.1-fold (95% CI: 1.1-4.0) increased risk of breast cancer.

We also evaluated the association between recent DMPA use for ≥ 12 months by breast cancer subtype (Table 4). While some variation in risk was observed across

tumors defined by clinical factors, hormone receptor expression, proxies of molecular subtype, or tumor histology, none of these differences were statistically significant.

DISCUSSION

This is the first large scale U.S. study specifically designed to evaluate the relationship between DMPA use and breast cancer risk. Recent users of DMPA for ≥ 12 months were observed to have a 2.2-fold increased risk of breast cancer in this population of women 20-44 years of age. The relevance of both recency and duration of use were supported by no observed increase risk among either former users of DMPA (those who last used DMPA more than 5 years ago) or among recent users of DMPA for less than 12 months. Despite being conducted in a U.S. population that is demographically and culturally quite different from the diverse populations included in the previously published studies of DMPA and breast cancer risk (which included women living in Costa Rica, New Zealand, Kenya, Mexico, Thailand, and South Africa), our results are quite similar to these prior studies which found that recent DMPA use was associated with 1.5 to 1.65-fold increased risks of breast cancer.(5-7)

Oral contraceptives have been shown across numerous studies to confer an increased risk of breast cancer only among recent users for at least 12 months, with a pooled analysis of the world's literature showing that among young women current users have a 24% increased risk of breast cancer, but that this risk also dissipates once oral contraceptive use ceases.(13) The stronger magnitude of the association with DMPA use seen in prior studies and in our own U.S.-based study compared to that seen in relation to oral contraceptives (based on the results of our analysis restricted to

users of hormonal contraceptives) may be attributable to the hormonal composition of these formulations and their different pharmacokinetics. The most common forms of oral contraceptives almost all contain both estrogen and progestin, and the dosing of these hormones can either be constant (monophasic preparations) or variable (biphasic or triphasic) over the course of a monthly cycle. They most typically also involve a one week hormone free break each month in between cycles. In contrast, DMPA is a progestin only contraceptive that involves a single injection every three months. With respect to its pharmacokinetics, serum concentrations of MPA are maintained at approximately 1.0 ng/mL for at least three months following a DMPA injection. MPA levels then decline to 0.2 ng/mL in the fifth and sixth months, and become undetectable 7.5 to 9 months after injection.⁽¹¹⁾ Ovulation resumes once MPA levels fall below 0.1 ng/mL.⁽¹⁰⁾ So based on its pharmacokinetics, even a single dose of DMPA results in a relatively lengthy and sustained exposure. There are also progestin only oral contraceptives, but they are rarely used so their relationship to breast cancer risk is not well known.

This is the first study to evaluate associations between DMPA use and risk of different breast cancer subtypes. While we observed a substantial 3.3-fold increased risk of poor prognosis triple-negative breast cancer associated with DMPA use, this result must be interpreted cautiously given the sample size limitations of this analysis and that the risk estimates for triple-negative and ER+ breast cancer were not statistically different. Relevant to this finding though is the observation in the only published study of oral contraceptive use and risk of different breast cancer subtypes defined by ER, PR, and HER2 status among premenopausal women that OC users for

at least one year have a 2.5-fold (95% CI: 1.4-4.3) increased risk of triple-negative breast cancer, but no increased risk of non-triple-negative breast cancer (case-case comparison p-value<0.01).(15) In this study the risks conferred by longer durations of OC use and fewer years since last use were also greater for triple-negative than for non-triple-negative breast cancer. The biological mechanisms underlying the potentially stronger relationships between hormonal contraceptives and risk of triple-negative breast cancer among premenopausal women are essentially unknown, but they certainly warrant further study given the relatively poorer prognosis of this more aggressive breast cancer subtype.(16-21)

Another relevant comparison is to the relationship between menopausal hormone therapy and breast cancer risk. One commonly used form of combined estrogen and progestin menopausal hormone therapy, including the form most commonly used in the United States, consists of oral conjugated estrogen in combination with MPA. This specific regimen was shown in the WHI trials to increase breast cancer risk. Studies examining use of menopausal hormones demonstrate that the risk of breast cancer returns to baseline within a few years of cessation of estrogen and progestin use.(12) A primary implication of these results is the importance of exogenous progestin use, and MPA use in particular, as having a promotional rather than initiatory role with respect to breast cancer risk. Though we had limited statistical power to fully assess various aspects of the timing and duration of DMPA use in relation to breast cancer risk, our results are generally consistent with other studies of exogenous progestin use in this regard and they add to the evidence that MPA use specifically confers an increased risk of breast cancer.

While few studies have evaluated the relationship between DMPA use and breast cancer risk, the evidence related to recent use is remarkably consistent. However, all of these studies are observational and therefore susceptible to different forms of bias. Given our case-control design, recall bias is a potential concern. DMPA is a unique exposure though given that is in an injection and serves as a contraceptive for a limited time period. Thus, difficulty in recall of this exposure in a population restricted to young adults is likely not a problem appreciably enough to bias our risk estimates. Confounding is also a potential concern given the differences in women who do and do not use DMPA as shown in Table 2. However, all analyses were adjusted for multiple a priori confounders and additional potential confounders were also carefully assessed. Though relatively large in its overall sample size, a limitation of this study was that only 10.9% of controls and 11.8% of cases had ever used DMPA. Consequently, our statistical power to assess more detailed aspects of patterns of DMPA use was limited. Lastly, the exclusion of women without a landline telephone could potentially bias our results. However, a recent study comparing women in the Seattle-Puget Sound region with and without a landline telephone found no differences in their frequencies of ever use of either injectable contraceptives or oral contraceptives(22) suggesting that any bias resulting from this exclusion is likely to have minimal impact.

Since breast cancer is relatively rare among young women existing clinical trial data are insufficient to evaluate this relationship and launching a new trial is not feasible. Consequently, the highest level of evidence will come from observational studies such as this one purposefully designed to address the association between DMPA use and breast cancer. With the addition of the results reported here there are

now five studies conducted over a diverse group of countries that have observed that recent DMPA use is associated with a 1.5 to 2.3-fold increased risk of breast cancer.(4-7) Mitigating the clinical and public health impact of this risk is the fact that breast cancer is rare among premenopausal women. However, there are numerous contraceptive options and so further clarifying the benefits and risks associated with each option is important as women consider what choices might be best for them.

ACKNOWLEDGEMENTS

The authors wish to acknowledge the substantial contributions of Kristine Wicklund, Christabel Fowler, and Anne Oswald in the conduct of this research study. Other staff members making important contributions to this work are: Nancy Blythe, Ann Bradshaw, Dante del Pilar, Nicole Donovan, Fran Fleck, Joia Hicks, Amy Hoffman, Dick Jacke, Kristine Maddux, Evan McKay, Susan McKeeth, Sarah Moore, Kathryn Nord, Patty Pride, Georgene Ranney, Tiffany Silver-Brace, Camille Taylor, Jodi Thiel, Loni Tipton, and Margaret Trzyna. Lastly, we want to acknowledge the time and generosity of all of the women who participated in this research.

REFERENCES

1. Chlebowski RT, Hendrix SL, Langer RD, Stefanick ML, Gass M, Lane D, et al. Influence of estrogen plus progestin on breast cancer and mammography in healthy postmenopausal women: the Women's Health Initiative Randomized Trial. *JAMA* 2003;289:3243-53.
2. The Women's Health Initiative Steering Committee. Effects of conjugated equine estrogen in postmenopausal women with hysterectomy: the Women's Health Initiative randomized controlled trial. *JAMA* 2004;291:1701-12.
3. Frost JJ, Jones RK, Woog V, Singh S, Darroch JE. Teenage sexual and reproductive behavior in developed countries: Country report for the United States. Occasional Report No. 8. New York, NY: The Alan Guttmacher Institute; 2001.
4. Lee NC, Rosero-Bixby L, Oberle MW, Grimaldo C, Whatley AS, Rovira EZ. A case-control study of breast cancer and hormonal contraception in Costa Rica. *J Natl Cancer Inst* 1987;79:1247-54.
5. Breast cancer and depot-medroxyprogesterone acetate: a multinational study. WHO Collaborative Study of Neoplasia and Steroid Contraceptives. *Lancet* 1991;338:833-8.
6. Paul C, Skegg DC, Spears GF. Depot medroxyprogesterone (Depo-Provera) and risk of breast cancer. *BMJ* 1989;299:759-62.
7. Shapiro S, Rosenberg L, Hoffman M, Truter H, Cooper D, Rao S, et al. Risk of breast cancer in relation to the use of injectable progestogen contraceptives and combined estrogen/progestogen contraceptives. *Am J Epidemiol* 2000;151:396-403.
8. Waksberg J. Sampling methods for random digit dialing. *J Am Stat Assoc* 1978;73:40.
9. Voigt LF, Davis S, Heuser L. Random digit dialing: the potential effect on sample characteristics of the conversion of nonresidential telephone numbers. *Am J Epidemiol* 1992;136:1393-9.

10. Mishell DR, Jr. Pharmacokinetics of depot medroxyprogesterone acetate contraception. *J Reprod Med* 1996;41:381-90.
11. Ortiz A, Hirol M, Stanczyk FZ, Goebelsmann U, Mishell DR. Serum medroxyprogesterone acetate (MPA) concentrations and ovarian function following intramuscular injection of depo-MPA. *J Clin Endocrinol Metab* 1977;44:32-8.
12. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormone replacement therapy: collaborative reanalysis of data from 51 epidemiological studies of 52,705 women with breast cancer and 108,411 women without breast cancer. *Lancet* 1997;350:1047-59.
13. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and hormonal contraceptives: collaborative reanalysis of individual data on 53,297 women with breast cancer and 100,239 women without breast cancer from 54 epidemiological studies. *Lancet* 1996;347:1713-27.
14. Begg CB, Gray R. Calculation of polychotomous logistic regression parameters using individualized regressions. *Biometrika* 1984;71:11-8.
15. Dolle JM, Daling JR, White E, Brinton LA, Doody DR, Porter PL, et al. Risk factors for triple-negative breast cancer in women under the age of 45 years. *Cancer Epidemiol Biomarkers Prev* 2009;18:1157-66.
16. Carey LA, Perou CM, Livasy CA, Dressler LG, Cowan D, Conway K, et al. Race, breast cancer subtypes, and survival in the Carolina Breast Cancer Study. *JAMA* 2006;295:2492-502.
17. Kim MJ, Ro JY, Ahn SH, Kim HH, Kim SB, Gong G. Clinicopathologic significance of the basal-like subtype of breast cancer: a comparison with hormone receptor and Her2/neu-overexpressing phenotypes. *Hum Pathol* 2006;37:1217-26.

18. Lund MJ, Trivers KF, Porter PL, Coates RJ, Leyland-Jones B, Brawley OW, et al. Race and triple negative threats to breast cancer survival: a population-based study in Atlanta, GA. *Breast Cancer Res Treat* 2009;113:357-70.
19. Sorlie T, Perou CM, Tibshirani R, Aas T, Geisler S, Johnsen H, et al. Gene expression patterns of breast carcinomas distinguish tumor subclasses with clinical implications. *Proc Natl Acad Sci U S A* 2001;98:10869-74.
20. Brown M, Tsodikov A, Bauer KR, Parise CA, Caggiano V. The role of human epidermal growth factor receptor 2 in the survival of women with estrogen and progesterone receptor-negative, invasive breast cancer: the California Cancer Registry, 1999-2004. *Cancer* 2008;112:737-47.
21. Kaplan HG, Malmgren JA. Impact of triple negative phenotype on breast cancer prognosis. *Breast J* 2008;14:456-63.
22. Voigt LF, Schwartz SM, Doody DR, Lee SC, Li CI. Feasibility of including cellular telephone numbers in random digit dialing for epidemiologic case-control studies. *Am J Epidemiol* 2011;173:118-26.

Table 1: Distribution of selected characteristics among controls and cases

Characteristic	Controls (n=919)		Cases (n=1,028)	
	n	%	n	%
Age (years)				
20-29	24	2.7	23	2.3
30-34	81	8.8	84	8.2
35-39	261	28.4	287	27.9
40-44	553	60.2	634	61.7
Reference year				
2004-2005	302	32.8	295	28.7
2006-2007	350	38.1	357	34.7
2008-2010	267	29.1	376	36.6
Race/ethnicity				
Non-Hispanic white	752	82.2	804	79.2
African American	31	3.4	51	5.0
Asian/Pacific Islander	82	9.0	115	11.3
Native American	19	2.1	26	2.6
Hispanic White	31	3.4	19	1.9
Missing	4		13	
Education				
High school or less	88	9.6	121	11.9
Post high school/some college	300	32.8	338	33.1
College graduate	347	37.9	372	36.4
Post college	181	19.8	190	18.6
Missing	3		7	
Annual household income				
<\$25,000	72	7.9	79	7.8
\$25,000-49,999	120	13.2	154	15.3
\$50,000-89,999	335	36.8	325	32.3
\$90,000+	383	42.1	449	44.6
Missing	9		21	
First-degree family history of breast cancer				
No	794	89.6	800	80.3
Yes	92	10.4	196	19.7
Missing	33		32	
BMI one year prior to reference date (kg/m²)				
<25.0	522	57.2	617	60.7
25.0-29.9	229	25.1	230	22.6
≥30	162	17.7	170	16.7
Missing	6		11	
Duration of oral contraceptive use (years)				
Never	101	11.0	118	11.6
<5.0	332	36.2	358	35.1
5.0-9.9	213	23.3	212	20.8
≥10	270	29.5	333	32.6
Missing	3		7	
Numer of full-term pregnancies				
Never	186	20.2	266	25.9
1-2	540	58.8	585	57.0
≥3	193	21.0	176	17.1
Missing	0		1	

Table 1, continued

Characteristic	Controls (n=919)		Cases (n=1,028)	
	n	%	n	%
Age at first full-term pregnancy (among parous women)				
<25	210	28.6	245	32.2
25-29	223	30.4	246	32.4
30-34	201	27.4	183	24.1
≥35	100	13.6	86	11.3
Missing	0		1	
Ever had a screening mammogram				
Never	463	50.4	439	42.8
Ever	455	49.6	586	57.2
Missing	1		3	

Table 2: Distribution of selected characteristics among controls who never used hormonal contraception, never used DMPA, and ever used hormonal contraception but never used DMPA

Characteristic*	Never used hormonal contraception (n=91)		Ever used DMPA (n=100)		Ever used hormonal contraception but never used DMPA (n=728)	
	n	%	n	%	n	%
Age (years)						
20-29	7	7.7	5	5.0	12	1.6
30-34	9	9.9	17	17.0	55	7.6
35-39	22	24.2	39	39.0	200	27.5
40-44	53	58.2	39	39.0	461	63.3
Race/ethnicity						
Non-Hispanic white	54	59.3	76	76.0	622	85.9
Asian/Pacific Islander	26	28.6	7	7.0	49	6.8
Other	11	12.1	17	17.0	53	7.4
Education						
High school or less	10	11.0	17	17.0	61	8.4
Post high school/some college	25	27.5	43	43.0	232	32.0
College graduate	42	46.2	27	27.0	278	38.3
Post college	14	15.4	13	13.0	154	21.2
First-degree family history of breast cancer						
No	78	87.6	88	93.6	628	89.3
Yes	11	12.4	6	6.4	75	10.7
Body mass index one year prior to reference date (kg/m²)						
<25.0	54	60.0	49	49.0	419	58.0
25.0-29.9	23	25.6	26	26.0	180	24.9
≥30.0	13	14.4	25	25.0	124	17.2
Duration of oral contraceptive use (years)						
Never	91	100.0	4	4.0	6	0.8
<5.0	0	0.0	42	42.4	290	39.9
5.0-9.9	0	0.0	28	28.3	185	25.5
≥10	0	0.0	25	25.3	245	33.7
Number of full-term pregnancies						
Never	38	41.8	11	11.0	137	18.8
1-2	41	45.1	67	67.0	432	59.3
≥3	12	13.2	22	22.0	159	21.8
Ever had a screening mammogram						
Never	48	52.7	57	57.0	358	49.2
Ever	43	47.3	43	43.0	369	50.8

Table 3: Relationship between depot-medroxyprogesterone acetate (DMPA) use and risk of invasive breast cancer

DMPA use category	Controls (n=919)		Cases (n=1,028)		OR*	95% CI
	n	%	n	%		
Ever use of DMPA						
Never use	819	89.1	907	88.2	1.0	ref
Ever use	100	10.9	121	11.8	1.2	0.9-1.6
Recency of DMPA use						
Recent use (last use <5 years ago)	24	2.6	36	3.5	1.5	0.9-2.7
Former use (last use ≥5 years ago)	76	8.3	85	8.3	1.1	0.8-1.5
Last use 5-10 years ago	42	4.6	34	3.3	0.8	0.5-1.3
Last use ≥10 years ago	34	3.7	51	5.0	1.4	0.9-2.1
Duration of DMPA use among recent users						
<12 months	9	1.0	4	0.4	0.5	0.1-1.6
≥12 months	15	1.6	32	3.1	2.2	1.2-4.2 [†]
Age at first DMPA use						
<25	40	4.4	43	4.2	1.1	0.7-1.8
25-29	27	2.9	39	3.8	1.4	0.8-2.3
30-34	24	2.6	22	2.1	0.8	0.4-1.5
≥35	9	1.0	17	1.7	2.0	0.9-4.6
Timing of use in relation to first full-term pregnancy						
Never had a full-term pregnancy	11	1.2	21	2.0	1.4	0.6-3.0
First used DMPA before first full-term pregnancy	30	3.3	24	2.3	0.8	0.4-1.3
First used DMPA after first full-term pregnancy	59	6.4	76	7.4	1.4	0.9-2.0
Timing of use in relation to most recent full-term pregnancy						
Never had a full-term pregnancy	11	1.2	21	2.0	1.4	0.6-3.0
First used DMPA before most recent full-term pregnancy	60	6.5	57	5.5	1.0	0.7-1.4
First used DMPA after most recent full-term pregnancy	29	3.2	43	4.2	1.5	0.9-2.5

* Odds ratios (ORs) are adjusted for age, year, body mass index, duration of oral contraceptive use, number of full-term pregnancies, family history of breast cancer, and history of screening mammography.

[†] p-value <0.05.

Table 4: Relationship between recent DMPA use for ≥ 12 months and risk of invasive breast cancer by clinical, molecular, and histopathologic subtype

	Never use		Recent use for ≥ 12 months				p-value for difference compared to the reference case group
	n	%	n	%	OR*	95% CI	
Controls	819	89.0	15	1.6	1.0	reference	
AJCC stage							
1	346	88.9	14	3.6	2.7	1.3-5.9 [†]	reference
2, 3, or 4	537	88.3	17	2.8	1.9	0.9-3.9	0.27
Tumor size, cm							
≤ 2.0	481	88.4	16	2.9	2.2	1.1-4.5 [†]	reference
> 2.0	395	88.6	14	3.1	2.1	0.99-4.4	0.87
ER status							
ER+	673	88.3	22	2.9	2.1	1.1-4.2 [†]	reference
ER-	226	87.9	10	3.9	2.5	1.1-6.4 [†]	0.63
ER/PR status							
ER+/PR+	611	88.3	20	2.9	2.1	1.1-4.3 [†]	reference
ER-/PR-	209	88.2	9	3.8	2.4	1.0-5.8 [†]	0.69
ER/PR/HER2 status							
ER+	673	88.3	22	2.9	2.2	1.1-4.3 [†]	reference
ER-/PR-/HER2-	154	86.5	9	5.1	3.3	1.4-7.8 [†]	0.28
ER-/HER2+	56	93.3	0	0.0	-	-	-
Histology							
Ductal	791	88.4	26	2.9	2.0	1.0-3.9 [†]	reference
Lobular	48	94.1	2	3.9	3.2	0.7-15.2	0.61
Other	58	85.3	3	4.4	3.7	1.0-13.8 [†]	0.30

* Odds ratios (ORs) are adjusted for age, year, body mass index, duration of oral contraceptive use, first-degree family history of breast cancer, and history of screening mammography.

[†] p-value < 0.05 .

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

Effect of depo-medroxyprogesterone acetate on breast cancer risk among women 20-44 years of age

Christopher I Li, Elisabeth F Beaber, Mei-Tzu C Tang, et al.

Cancer Res Published OnlineFirst February 27, 2012.

Updated version Access the most recent version of this article at:
doi:[10.1158/0008-5472.CAN-11-4064](https://doi.org/10.1158/0008-5472.CAN-11-4064)

Author Manuscript Author manuscripts have been peer reviewed and accepted for publication but have not yet been edited.

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, contact the AACR Publications Department at permissions@aacr.org.