

Loss of P27^{Kip1} Expression Correlates with Tumor Grade and with Reduced Disease-free Survival in Primary Superficial Bladder Cancers¹

Alessandro Sgambato,² Mario Migaldi, Beatrice Faraglia, Lorella Garagnani, Gianpiero Romano, Carmela De Gaetani, Paolo Ferrari, Giovanni Capelli, Gian Paolo Trentini, and Achille Cittadini

Centro di Ricerche Oncologiche "Giovanni XXIII"—Istituto di Patologia Generale [A. S., B. F., G. R., A. C.] and Istituto di Igiene [G. C.], Catholic University, 00168 Rome; Dipartimento di Scienze Morfologiche e Medico Legali, Sezione di Anatomia Patologica, University of Modena, Modena [M. M., L. G., C. D. G., G. P. T.]; and Divisione di Urologia, Azienda Unità Sanitaria Locale Modena 41100 [P. F.], Italy

ABSTRACT

p27^{Kip1} is a member of the Cip1/Kip1 family of cyclin-dependent kinase inhibitors and is a potential tumor suppressor gene. We previously reported a deregulated expression of p27^{Kip1} in a series of human cancer cell lines and in primary breast and colon cancers. Moreover, p27^{Kip1} has been reported as an important prognostic factor in primary lung, breast, colon, and prostate cancers. In this study, we evaluated the prognostic value of p27^{Kip1} in a series of 96 superficial (pTa-1) human bladder carcinomas. High (>50% positive cells), moderate (25–50%), and low (<25%) p27^{Kip1} staining was observed in 39 (41%), 19 (20%), and 38 (39%) of the 96 primary superficial bladder cancers, respectively. No significant association was found between the expression level of p27^{Kip1} and tumor stage. Decreased p27^{Kip1} staining correlated with higher tumor grade ($P = 0.001$). Interestingly, a significant association was observed between increased expression of p27^{Kip1} and positivity for p53 (>20% positive cells; $P < 0.001$). A significant correlation was also observed between low expression of p27^{Kip1} and decreased disease-free survival ($P = 0.0003$ by log-rank test) and overall survival ($P = 0.01$ by log-rank test). Furthermore, on multivariate analysis, low p27^{Kip1} protein expression was an independent predictor of reduced disease-free survival ($P = 0.018$; relative risk = 1.95) second only to tumor stage. These data indicate that p27^{Kip1} protein is frequently expressed at low level in poorly differentiated tumors and suggest that this protein might represent a useful prognostic marker for disease recurrence and overall survival in superficial bladder carcinomas.

INTRODUCTION

Alterations in genes involved in the regulation of normal cell cycle progression are frequent events in human cancers. The CDI³ proteins are potent negative regulators of the cell cycle and are potential tumor suppressor genes, the loss of which might play an important role in the development of human tumors (1, 2).

Indeed, loss of the CDIs p15^{Ink4b}, p16^{Ink4a}, and p18^{Ink4c} by gene mutation, deletion, and/or methylation has been observed in a variety of human tumors (3, 4), and p16^{Ink4a}-deficient mice develop spontaneous tumors at an early age and are highly sensitive to carcinogens (5).

On the other hand, several independent studies have found that alterations in the integrity of the human p27^{Kip1} and p21^{Cip1} genes, which belong to the Kip/Cip family of CDIs, are rare events in a variety of human primary tumors and cancer cell lines (6–13). However, it has been reported that reduced expression of p27^{Kip1} predicts

poor survival in breast, colon, prostate, esophagus, and lung cancer patients (14–18). The decreased expression of p27^{Kip1} is due to posttranscriptional events and has been associated with increased proteasome-mediated degradation of the protein (15, 16).

Alterations of the p53 tumor suppressor gene have been reported for a variety of tumors, including bladder carcinomas. The p53 protein is produced at a very low level in normal cells and can act as a transcriptional factor involved in many cellular functions, including the regulation of cell proliferation and apoptosis. In the absence of a functional p53 protein, cells fail to repair DNA damage or to undergo to apoptosis and are more susceptible to neoplastic transformation (19). Mutant p53 has, usually, an increased half-life and is more easily detected by immunohistochemistry than the wild-type protein. In addition to gene mutations, several viral and cellular proteins (*i.e.*, mdm2) can bind to and stabilize the p53 protein, thus, inhibiting its activity (20). p53 overexpression has been shown to correlate with tumor grade and stage in primary bladder carcinomas (21), and it has been suggested that p53 mutations might play an important role in bladder cancer progression (22).

In this study, we have evaluated by immunostaining the expression of p27^{Kip1} in 96 primary superficial bladder carcinomas and have correlated the results obtained with tumor grade and stage. The expression of p27^{Kip1} showed a significant relationship with tumor grade and with the expression of the p53 protein. p27^{Kip1}, but not p53, expression correlated with the prognosis of patients, both in terms of disease-free and overall survival, in our series. On a multivariate Cox regression analysis, loss of p27^{Kip1} expression was an independent predictor of early recurrence. The implications of these findings are discussed.

MATERIALS AND METHODS

Patient Characteristics and Tissue Samples. A cohort of 96 patients who underwent routine surgery for bladder cancer at the Division of Urology, County Hospital (Modena, Italy) between April 1990 and December 1995 were used for this study. They included 83 men (86%) and 13 women (14%), with a mean age at diagnosis of 68 years (range, 29–92) and a mean follow-up of 50 months (range, 24–102). All of the patients underwent transurethral resection with curative intent, and none of them had received any therapy before surgery. The follow-up of patients was conducted according to standard clinical practice, and all of the patients were treated by intravesical *Bacillus Calmette-Guerin* instillations, in case of recurrence. Tissue specimens were obtained from the files of the Service of Anatomical Pathology, University of Modena. For molecular analysis, fresh tissues were frozen and stored at -140°C immediately after surgery. The corresponding formalin-fixed paraffin-embedded specimens were used for immunohistochemical analyses. Histological grading and staging were assessed according to WHO (23) and tumor-node-metastasis (24) classification, respectively. Other clinic-pathological parameters, such as age and gender, were also recorded and are shown in Table 1. The samples were coded, and the names of the patients were not revealed.

Immunohistochemistry. All immunohistochemical analyses were performed on routinely processed, formalin-fixed, paraffin-embedded tissues using an avidin-biotin complex immunoperoxidase technique. Successive 5- μm tissue sections were cut from blocks selected for the presence of representative

Received 12/18/98; accepted 5/3/99.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported in part by grants from "Co-finanziamento MURST" 1998 and the University of Modena and by an award from Associazione Angela Serra for Cancer Research (Modena, Italy). A. S. was supported by a fellowship from the Italian Association for Cancer Research (AIRC).

² To whom requests for reprints should be addressed, at Centro di Ricerche Oncologiche "Giovanni XXIII," Università Cattolica del Sacro Cuore, Largo Francesco Vito 1, 00168 Rome, Italy. Phone: 39-06-3016619; Fax: 39-06-3012753; E-mail: crogxxiii@rm.unicatt.it.

³ The abbreviations used are: CDI, cyclin-dependent kinase inhibitor; CI, confidence interval; RR, relative risk.

Table 1 p27^{Kip1} expression and clinicopathological parameters in 96 primary superficial bladder carcinomas

	Total	p27 ^{Kip1} expression			P ^a
		High (%)	Moderate (%)	Low (%)	
Age (yr)					
≥65	62	24 (39%)	10 (16%)	28 (45%)	
<65	34	15 (44%)	9 (26%)	10 (30%)	0.26
Sex					
Male	83	33 (40%)	16 (19%)	34 (41%)	
Female	13	6 (46%)	3 (23%)	4 (31%)	0.78
Tumor grade					
Well differentiated	13	3 (23%)	6 (46%)	4 (31%)	
Moderately differentiated	51	30 (59%)	6 (12%)	15 (29%)	0.001 ^b
Poorly differentiated	32	6 (19%)	7 (22%)	19 (59%)	
Tumor stage					
A	42	21 (50%)	8 (19%)	13 (31%)	
I	54	18 (33%)	11 (20%)	25 (47%)	0.22 ^c
p53 expression ^c					
Positive (≥20%)	52	30 (58%)	9 (17%)	13 (25%)	
Negative (<20%)	44	9 (20%)	10 (23%)	25 (57%)	<0.001

^a Statistical analyses were performed by the Pearson χ^2 test. $P < 0.05$ was considered significant.

^b The same values were also obtained when the Fisher's exact test was used for the analysis.

^c The values refer to the results obtained using the monoclonal Pab1801 antibody. See text for more details.

tumor tissue and mounted on poly-L-lysine-coated slides. After incubation in an oven at 43°C for 12 h to prevent sections from lifting from the slide during the antigen-retrieval step, sections were dewaxed and rehydrated. They were then submerged in a citrate buffer (10 mM; pH 6.0) and microwaved for a total of 10 min at 750 W. Sections were then treated to block endogenous peroxidase and, after blocking with goat (for the polyclonal anti-p27^{Kip1} antibody) or horse (for the monoclonal anti-p53 antibodies) serum for 1 h at room temperature, the primary antibodies were applied overnight at 4°C in a high-humidity chamber. After washing, immunostaining was performed using the Vectastain ABC and diaminobenzidine kits (Vector Laboratories, Burlingame, CA), as described. Sections were then counterstained with 1% modified Harris hematoxylin, dehydrated, and mounted.

The polyclonal anti-p27^{Kip1} antibody, raised against a peptide corresponding to amino acids 181–198 mapping in the COOH-terminus of the human p27^{Kip1} protein (Santa Cruz Biotechnology, Santa Cruz, CA), was used at a concentration of 1 μ g/ml (in PBS with 10% goat serum) because this gave good nuclear staining with minimal background. As a negative control, a duplicate section of each tissue sample was immunostained in the absence of the primary antibody. A breast carcinoma with known positive immunostaining for p27^{Kip1} served as a positive control. A strong nuclear staining of lymphocytes provided a useful internal positive control for preservation of the p27^{Kip1} immunogenicity in most sections examined. Within tumor cells, the staining reaction for p27^{Kip1} was often localized both in the nucleus and in the cytoplasm, but the nuclear staining was predominant and, in most of the cases, exclusive. Some tumor cells also showed a clear nucleolar signal. The significance of this finding is unknown. The specificity of the reaction for p27^{Kip1} was demonstrated by inhibition of the immunohistochemical staining in positive samples by preincubating the antibody with a 100-fold excess of the immunizing peptide (Santa Cruz Biotechnology) for 1 h at room temperature (data not shown). Confirmatory results were obtained when a different monoclonal anti-p27^{Kip1} antibody (NCL-p27, clone 1B4; Novocastra, Newcastle, United Kingdom) was used to stain selected cases for which duplicate sections were available. The cytoplasmic staining was markedly reduced with this antibody (data not shown). The monoclonal PAb1801 (DAKO, Milan, Italy) and DO7 (Novocastra) anti-p53 antibodies were used diluted 1:750 in PBS. A p53-positive breast carcinoma with known positive nuclear reaction for p53 was used as a positive control. The immunostaining of duplicate sections in the absence of the primary antibody was used as a negative control.

The number of cells with positive nuclear reaction for both p27^{Kip1} and p53 was calculated semiautomatically by means of a computer-assisted cellular image analyzer on a total of 1000 nuclei/case. All scoring and interpretations of the results were made by two of the authors independently (A. S. and M. M.) without knowledge of clinical outcome or other clinic-pathological variables. The few cases with discrepant scoring were reevaluated jointly on a second

occasion, and agreement was reached. Nuclei were considered positive when they showed a distinct brown color in the absence of background staining. Positive and negative control slides were included within each batch of slides. Tumors showing undetectable or negligible levels of p27 expression were scored as negative. Positives were categorized as low (<25%), moderate (25–50%), and high (>50%; Refs. 14, 16–18). The cutoff value of p53 immunostaining used to define p53 overexpression for statistical analysis was 20% of positive tumor nuclei (25, 26).

Statistical Analyses. The association between p27^{Kip1} and p53 expression and other variables shown in Table 1 were calculated using contingency table methods and tested for significance using the Pearson's χ^2 test. Interrater agreement of the results obtained with the two anti-p53 antibodies was evaluated by Cohen's κ coefficient (27). Survival curves were calculated using the Kaplan-Meier method, and the log-rank test was used for the analysis. Patients who died of other causes during the follow-up period were treated as censored data in the survival analyses. Univariate and multivariate RRs were calculated using Cox proportional hazards regression. The RR for age represents the hazard increase/1-year increase in age. For sex, the RRs are given as male *versus* female. For grade and stage, the G1 grade and the pTa stage were used as baseline, respectively. All calculations were performed using the STATA 5.0 statistical software package (Stata Corporation, College Station, TX), and the results were considered statistically significant when the P was <0.05.

RESULTS

Expression of p27^{Kip1} Correlates with Tumor Grade in Primary Superficial Bladder Cancers. To investigate the significance of p27^{Kip1} in human bladder cancer, the expression of this protein was evaluated by immunostaining in a series of 96 primary human superficial bladder carcinomas. Within tumor cells, the staining was predominantly nuclear. However, a concomitant diffuse, less intense cytoplasmic staining was also observed in 37 (38.5%) of the 96 tumors (Fig. 1 and data not shown). Only cells with a clear nuclear staining were considered positive. The percentage of positive tumor cells within a microscopic field ranged from 0–90%, and intratumoral heterogeneity of p27^{Kip1} nuclear immunostaining was common (Fig. 1). When present, the normal mucosa was also assessed for p27^{Kip1} staining. We found that normal urothelium constantly demonstrated a weak nuclear reaction for p27^{Kip1} (data not shown).

High (>50% positive cells), moderate (25–50%), and low (<25%) p27^{Kip1} staining were observed in 39 (41%), 19 (20%), and 38 (39%) of the 96 primary superficial bladder cancers, respectively. High or moderate p27^{Kip1} expression was found in 9 of 13 (69%) well-differentiated tumors (G1), 36 of 51 (71%) moderately differentiated tumors (G2), and 13 of 32 (41%) poorly differentiated tumors (G3). Thus, p27^{Kip1} expression was strongly associated with tumor grade in superficial bladder cancer ($P = 0.001$). No correlation was found between p27^{Kip1} expression and tumor stage in superficial bladder cancers ($P = 0.2$; Table 1).

Prognostic Significance of p27^{Kip1} Expression in Primary Superficial Bladder Cancer. Thirty-two of 38 superficial tumors (84%) with low expression of p27^{Kip1}, 10 of 19 moderate expressors (53%), and 20 of 39 high expressors (51%) recurred in our series of tumors during the period of follow-up. This difference was highly significant ($P = 0.005$). Similarly, 10 of 38 low expressor tumors (26%), 2 of 19 moderate expressors (11%), and 2 of 39 tumors (5%) expressing high levels of p27^{Kip1} died of disease during the period of follow-up. This difference was also significant ($P < 0.03$). Hence, recurrence and death occurred more frequently when the patients with primary bladder cancers in our series showed a reduced expression of p27^{Kip1} compared with tumors showing high expression of p27^{Kip1}.

The Kaplan-Meier curves of disease-free survival and overall survival within patients with low (<25% of cells) *versus* moderate or high expression of p27^{Kip1} showed a highly significant separation ($P = 0.0003$ and 0.01 by log-rank test, respectively). When all of the




Fig. 1. Examples of p27^{Kip1} immunostaining in three representative cases of bladder carcinomas. A, more than 90% of neoplastic cells show a strong, positive staining of the nuclei (high expressor) and a diffuse, faint cytoplasmic reaction. B, less than 50% of nuclei (moderate expressor) can be regarded as positive. C, only few nuclei can be regarded as positive (low expressor). A strong nuclear staining of lymphocytes (arrows) provided a useful internal positive control for preservation of the p27^{Kip1} immunogenicity. ($\times 100$)

groups (*i.e.*, low, moderate, and high expressor tumors) were included in the analysis, there was a clear trend for a statistically significant association between reduced expression of p27 and worse prognosis both in terms of disease-free survival ($P = 0.001$) and overall survival ($P = 0.03$; Fig. 2).

Positive Correlation between p27^{Kip1} and p53 Expression in Primary Bladder Cancer. The expression of p53 protein was evaluated by immunostaining in the same series of bladder tumors using the Pab1801 and the DO7 monoclonal antibodies. As mentioned in "Materials and Methods," a tumor was considered positive for p53 only if definite nuclear staining was present. Tumor samples were divided in two categories, positive or negative, by estimating the percentage of stained nuclei. The presence of $>20\%$ positive cells was

used as a cutoff to define positive tumors because, in previous studies, it gave the best prognostic value for p53 (25, 26). Normal urothelium demonstrated no immunoreactivity for p53 (data not shown).

We observed a good, although not absolute, agreement between the results obtained with the two antibodies. The Cohen's K coefficient was 0.65, which corresponded to a substantial (82.41%) amount of agreement. In fact, 52 (54%) and 41 (43%) of 96 tumors were positive for p53 immunostaining when tested with the Pab1801 and the DO7 antibodies, respectively. Positivity for p53, as assessed with the Pab1801, was correlated with the differentiation status. In fact, overexpression of p53 was observed in 6 of 13 (46%) well-differentiated tumors, 39 of 51 (77%) moderately differentiated tumors, and 7 of 32 (22%) poorly differentiated tumors ($P < 0.0001$). No correlation was observed with tumor stage. In fact, overexpression of p53, as assessed with the Pab1801 antibody, was found in 25 of 42 T_a (59%) and 27 of 54 T₁ (50%) tumors ($P = 0.3$). When assessed with the DO7 antibody, overexpression of p53 was observed in 4 of 13 G1 (31%), 29 of the 51 G2 (57%), and 8 of 32 G3 (25%) tumors ($P = 0.01$). No significant association was observed between p53 overexpression and tumor stage when the results obtained with the DO7 antibody were used for the analysis ($P = 0.4$).

Interestingly, we observed a significant relationship between the expression levels of p27^{Kip1} and positivity for p53 in our series of patients. In fact, 30 of 39 high (77%), 9 of 19 moderate (47%), and 13 of 38 low (34%) p27^{Kip1}-expressing tumors were positive for p53, as assessed by the Pab1801 antibody ($P = < 0.001$; Table 1). This association was also true, although weaker, when p53 expression was assessed using the DO7 antibody. With this antibody, positivity for p53 was found in 23 of 39 high (59%), 7 of 19 moderate (37%), and 11 of 38 low (29%) p27-expressing tumors ($P = 0.02$).

We then examined whether positivity for p53, as assessed with the Pab1801, was associated with disease-free survival and overall survival after surgery of the 96 cases of superficial bladder tumors in our series. The Kaplan-Meier survival curve showed a tendency for patients with positive tumors to have a poorer prognosis than patients with negatively stained tumors. However, no statistically significant association of the p53 staining was observed with both disease-free survival ($P = 0.07$ by log-rank test) and overall survival ($P = 0.05$ by log-rank test). When assessed with the DO7 antibody, overexpression of p53 was not associated with disease-free survival ($P = 0.6$ by log-rank test), nor with overall survival ($P = 0.6$ by log-rank test).

p27^{Kip1} Expression Is an Independent Prognostic Factor of Disease-free Survival in Superficial Bladder Cancers. As expected, in an univariate analysis, tumor stage was significantly associated with disease-free survival ($P = 0.0001$ by log-rank test) and overall survival ($P = 0.01$) in our series of patients. On the other hand, tumor grade was significantly associated with disease-free survival ($P = 0.01$) but not overall survival ($P = 0.07$).

When a Cox proportional hazards model was constructed that included age of patients at the diagnosis, sex, tumor grade and stage, and p53 and p27^{Kip1} expression, p27^{Kip1} was a strong independent predictor of disease-free survival ($P = 0.018$; RR = 1.95), second only to tumor stage (Table 2). For overall survival, age at diagnosis was the only independent prognostic marker ($P = 0.038$) in our series of patients when all of the variables were included in the analysis (Table 3). As shown in Table 3, the power of this variable was also not very strong (RR = 1.084; 95% CI = 1.005- 1.170) and needs to be confirmed by a study that includes larger patient numbers and a longer follow-up period. It is to be noted, however, that the RR given represents the hazard ratio/1-year increase in age. We also observed a tendency for patients with low p27^{Kip1}-expressing tumors to have a shorter survival than did subjects with moderate or high p27^{Kip1} expression (RR = 2.38), but this association was not significant




Fig. 2. Kaplan-Meier curve for disease-free survival (A) and overall survival (B) in 96 patients who underwent surgery for primary superficial bladder carcinomas, stratified according to p27^{Kip1} expression. Decreased p27^{Kip1} expression was significantly associated with early recurrence ($P = 0.001$ by log-rank test) and dead of disease ($P = 0.03$ by log-rank test). Significance was stronger when negative or low expressor tumors (<25% of positive tumor cells) were plotted against both moderate (25–50%) and high (>50%) expressor tumors, both in terms of disease-free survival ($P = 0.0003$ by log-rank test) and overall survival ($P = 0.01$ by log-rank test).

Table 2. Contribution of various potential prognostic factors to disease-free survival by Cox regression analysis in 96 superficial bladder cancers

Variable	RR	95% CI	P
Age	1.005	0.976–1.035	0.741
Sex	1.04	0.455–2.378	0.926
Grade	1.112	0.399–3.098	0.839
Stage	2.489	1.293–4.791	0.006
p27 ^a	1.951	1.122–3.391	0.018
p53 ^b	0.917	0.516–1.629	0.767

^a The RR is given as low (<25% positive cells) versus moderate or high expression.

^b The RR is given as positive ($\geq 20\%$ positive cells) versus negative tumors.

($P = 0.17$; 95%CI = 0.69–8.18). Age was included in the model because it is essential in this type of analysis. It is noteworthy that when age was not included in the model, only tumor stage and p27^{Kip1} expression were statistically significant predictors of overall survival ($P = 0.006$ and 0.009, respectively).

DISCUSSION

p27^{Kip1} belongs to the Cip1/Kip1 family of CDIs and it likely plays an important role in regulating cell proliferation during vertebrate development and tumorigenesis. In fact, p27-deficient mice exhibit multiorgan hyperplasia and develop pituitary tumors. Moreover, although alterations in the integrity of the human p27^{Kip1} gene are rare events in human primary tumors and cancer cell lines (6–13), it has been reported that reduced expression of p27^{Kip1} predicts poor survival in breast, colon, prostate, esophagus, and lung cancer patients (14–18). In this study, expression of p27^{Kip1} was evaluated by immunostaining in 96 superficial bladder cancers. Superficial bladder cancers are localized in the mucosal layer (pTa) or invade the lamina propria without extension into the muscularis propria (pT1) and have mostly a good prognosis, being that metastasis is a rare event. However, the majority of them recur after surgery and require additional treatments.

To our knowledge, this is the first study investigating the expression of the p27^{Kip1} protein in a large series of superficial bladder

carcinomas. We found that a reduced expression of p27^{Kip1} is associated with tumor grade ($P = 0.001$) and with disease-free survival ($P = 0.0003$ by log-rank test) and overall survival ($P = 0.01$ by log-rank test) in superficial bladder cancers. In a multivariate analysis, low expression of p27^{Kip1} (<25% positive cells) was a powerful predictor of recurrence ($P = 0.018$; RR = 1.95), second only to tumor stage ($P = 0.006$; RR = 2.48), which is currently the strongest independent prognostic parameter in superficial bladder cancers. These results are in agreement with previous data on the correlation between p27^{Kip1} expression and tumor grade and on the prognostic role of p27^{Kip1}, and they further support the hypothesis that loss of p27^{Kip1} confers a selective growth advantage to tumor cells and might play an important role in the development of a variety of human tumors (14–18).

In preliminary studies, we observed that the frequency of loss of p27^{Kip1} in invasive tumors is almost identical to that observed in superficial tumors (data not shown). Regardless of the underlying molecular mechanism, these findings strongly suggest that alterations in the expression of p27^{Kip1} may be an early event in the multistep process of bladder carcinogenesis. Previous studies have demonstrated that the reduced expression of p27^{Kip1} observed in tumor cells may result from an accelerated degradation of the protein mediated by the ubiquitin-proteasome pathway (16, 28). Further *in vitro* and *in vivo* studies are required to elucidate whether alterations in the ubiquitin pathway are also responsible for the reduced expression of p27^{Kip1} observed in bladder cancer cells.

Table 3. Contribution of various potential prognostic factors to overall survival by Cox regression analysis in 96 superficial bladder cancers

Variable	RR	95% CI	P
Age	1.084	1.005–1.170	0.038
Sex	1.626	0.2–13.229	0.649
Grade	0.497	0.032–7.793	0.618
Stage	5.333	0.763–37.282	0.092
p27 ^a	2.381	0.693–8.178	0.168
p53 ^b	1.784	0.541–5.878	0.951

^a The RR is given as low (<25% positive cells) versus moderate or high expression.

^b The RR is given as positive ($\geq 20\%$ positive cells) versus negative tumors.

An unexpected finding of the present study was the significant relationship between the expression levels of p27^{Kip1} and that of the p53 protein in our series of superficial bladder cancers. This relationship was true when the level of the p53 protein was assessed using both the Pab1801 and the DO7 monoclonal antibodies ($P < 0.001$ and 0.02 , respectively). Analysis of the *p53* gene by PCR-single strand conformation polymorphism analysis and sequencing was able to detect mutations only in 13 tumors. Most of these tumors (9 of 13) were G3 tumors, and there was no significant relationship between the level of expression of p27^{Kip1} and the status of the *p53* gene (data not shown). Thus, this relationship only occurs at a posttranscriptional level for both proteins.

Accumulation of the p53 protein is associated with stabilization and functional inactivation of the protein, which can be due not only to gene mutations but also to binding of the p53 protein to several cellular and/or viral proteins (*i.e.*, mdm2; 20). Our results suggest that when p53 protein is overexpressed, and thus presumably inactive, cells most frequently retain p27^{Kip1} expression which, on the other hand, is most frequently lost in the presence of a functional p53 protein. Both p53 and p27^{Kip1} regulate cell cycle progression through a similar mechanism (29–32). In fact, a functional p53 can block cell cycle progression by inducing increased expression of p21^{Cip1}, which belongs to the same family of CDKI than p27^{Kip1} and targets the same cyclin/CDK complexes (30). Thus, it can be assumed that cells would not get any growth advantage from alterations (*i.e.*, inactivation of p53 and loss of p27^{Kip1}) that target the same pathway of cell growth regulation (29–32). This would explain why loss of p27^{Kip1} and accumulation of p53 are mutually exclusive in our series of tumors. A similar correlation has been observed between other important regulators of cell cycle progression, such as overexpression of cyclin D1 and loss of the pRb protein (33, 34) and loss of pRb and loss of p16^{Ink4a} (32, 35, 36), which also occur in distinct subsets of human tumors.

On the other hand, our data suggest that the relationship between p53 and p27^{Kip1} expression only occurs at a posttranscriptional level. In fact, there was no relationship between p27^{Kip1} expression and the status of the *p53* gene (data not shown). Moreover, most of the p53 overexpressing tumors retained a normal *p53* gene, as is the case for the *p27^{Kip1}* gene, which is rarely mutated in tumor cells (6–13). An alteration in a common mechanism regulating the abundance of both proteins at a posttranscriptional level might be responsible for this relationship. (16, 28, 37, 38). It is also possible that a more general deregulation of the mechanisms controlling protein stability occurs in tumor cells and might involve several proteins other than p53 and p27^{Kip1}. Studies are ongoing to test this hypothesis in *in vitro* studies.

We could not find any significant association between p53 expression and the clinical outcome of the patients in our series, both in terms of disease-free survival and overall survival. Conflicting data have been reported in the literature regarding the prognostic significance of p53 accumulation in superficial bladder carcinomas (22, 25, 26). Our results are in agreement with previous studies that also found that overexpression of p53 is not a prognostic marker in superficial bladder carcinomas (25, 39, 40). We could not find any association between p53 expression and prognosis in our series of patients, even when different cutoffs were used to discriminate between negative and positive tumors (data not shown). These findings are consistent with the hypothesis that nuclear accumulation of the p53 protein becomes more important and apparent in the late stages of bladder carcinogenesis (25).

We previously reported a significant relationship between the level of expression of p27^{Kip1} and that of cyclin D1 and cyclin E in a series of human cancer cell lines including breast, colon, prostatic, and esophageal cancer (41). A significant association between the expres-

sion levels of p27^{Kip1} and that of cyclin D1 has been also observed in primary breast (42) and colon (42, 43) carcinomas and in esophageal cancer cell lines (44, 45). This is the first study evaluating the levels of expression of the p27^{Kip1} protein in primary bladder cancer. It has been reported that the expression level of the cyclin D1 protein correlates with disease-free survival in superficial bladder cancers (46). It will be of interest to evaluate whether any relation exists between the expression levels of p27^{Kip1} and that of cyclin D1, cyclin E and/or other cell cycle-related proteins, such as p21^{Cip1}, in these tumors. Studies are ongoing to evaluate the expression levels of these molecules in our series of superficial bladder cancers and to extend these studies on a larger series of patients with a longer follow-up.

In conclusion, we demonstrated a clear correlation of p27^{Kip1} expression with tumor grade and p53 expression in superficial bladder carcinomas. Decreased p27^{Kip1} expression was also shown to be associated with an increased risk of recurrence and death for disease in patients with superficial bladder carcinomas who undergo potential curative surgery. Bladder cancer is the eighth most common cancer worldwide in men, and over 50,000 new cases are diagnosed each year in the United States. Fortunately, about 80% of the patients present with superficial tumors and respond to transurethral resection. However, even patients with tumor confined to the mucosa (pTa) frequently recur (more than 50% during the 1st year) and can progress into invasive disease or metastases (10–15% of the cases; Ref. 47). Thus, our findings will certainly be of interest, considering the scarcity of useful prognostic factors able to accurately predict the clinical outcome of patients with superficial bladder cancers. Low p27^{Kip1} expression might help to identify patients at higher risk of recurrence and to select candidates for an adjuvant treatment. Additional studies on a larger series of cases are required, however, to confirm these results and to elucidate the possible functional significance of the observed relationship between p27^{Kip1} and p53.

ACKNOWLEDGMENTS

We gratefully acknowledge P. Manni and R. Iotti for technical assistance and all of the members of the Institute of General Pathology, Catholic University (Rome, Italy) for helpful discussions and suggestions.

REFERENCES

- MacLachlan, T. K., Sang, N., and Giordano, A. Cyclins, cyclin-dependent kinases and Cdk inhibitors: implications in cell cycle control and cancer. *Crit. Rev. Eukaryot. Gene Expr.*, 5: 127–156, 1995.
- Weinstein, I. B., and Zhou, P. Cell cycle control gene defects and human cancers. *In: J. R. Bertino (ed.), Encyclopedia of Cancer*, Vol. 1, pp. 256–267. New York: Academic Press, 1997.
- Guan, K.-L., Jenkins, C. W., Li, Y., Nichols, M. A., Wu, X., O'Keefe, C. L., Matera, A. G., and Xiong, Y. Growth suppression by p18, a p16^{INK4/MTS1} and p14^{INK4/MTS2}-related CDK6 inhibitor, correlates with wild-type Rb function. *Genes Dev.*, 8: 2939–2952, 1994.
- Herman, J. G., Jen, J., Merlo, A., and Baylin, S. B. Hypermethylation-associated inactivation indicates a tumor suppressor role for p15^{INK4B1}. *Cancer Res.*, 56: 722–727, 1996.
- Serrano, M., Lee, H.-W., Chin, L., Cordon-Cardo, C., Beach, D., and Depinto, R. A. Role of the *INK4a* locus in tumor suppression and cell mortality. *Cell*, 85: 27–37, 1996.
- Bullrich, F., MacLachlan, T. K., Sang, N., Druck, T., Veronese, M. L., Allen, S. L., Chiorazzi, N., Koff, A., Heubner, K., Croce, C. M., and Giordano, A. Chromosomal mapping of members of the *cdc2* family of protein kinases, *cdk3*, *cdk6*, *PISSLRE*, and *PITALRE*, and a cdk inhibitor, p27^{Kip1}, to regions involved in human cancer. *Cancer Res.*, 55: 1199–1205, 1995.
- Ferrando, A. A., Balbin, M., Pendas, A. M., Vizoso, F., Velasco, G., and Lopez-Otin, C. Mutational analysis of the human cyclin-dependent kinase inhibitor p27^{Kip1} in primary breast carcinomas. *Hum. Genet.*, 97: 91–94, 1996.
- Kawamata, N., Morosetti, R., Miller, C. W., Park, D., Spirin, K. S., Nakamaki, T., Takeuchi, S., Hatta, Y., Simpson, J., Wilczynski, S., Lee, Y. Y., Bartram, C. R., and Koeffler, H. P. Molecular analysis of the cyclin-dependent kinase inhibitor gene *p27/Kip1* in human malignancies. *Cancer Res.*, 55: 2266–2269, 1995.
- Morosetti, R., Kawamata, N., Gombart, A. F., Miller, C. W., Hatta, Y., Hiramata, T., Said, J. W., Tomonaga, M., and Koeffler, H. P. Alterations of the *p27^{Kip1}* gene in

- non-Hodgkin's lymphomas and adult T-cell leukemia/lymphoma. *Blood*, 86: 1924–1930, 1995.
10. Pietenpol, J. A., Bohlander, S. K., Sato, Y., Papadopoulos, N., Liu, B., Friedman, C., Trask, B. J., Roberts, J. M., Kinzler, K. W., Rowley, J. D., and Vogelstein, B. Assignment of the human *p27Kip1* gene to 12p13 and its analysis in leukemias. *Cancer Res.*, 55: 1206–1210, 1995.
 11. Ponce-Castaneda, M. V., Lee, M-H., Latres, E., Polyak, K., Lacombe, L., Montgomery, K., Mathew, S., Krauter, K., Sheinfeld, J., Massague, J., and Cordon-Cardo, C. *p27^{KIP1}*: chromosomal mapping to 12p12–12p13.1 and absence of mutations in human tumors. *Cancer Res.*, 55: 1211–1214, 1995.
 12. Shiohara, M., El-Deiry, W. S., Wada, M., Nakamaki, T., Takeuchi, S., Yang, R., Chen, D-L., Vogelstein, B., and Koeffler, H. P. Absence of WAF1 mutations in a variety of human malignancies. *Blood*, 84: 3781–3784, 1994.
 13. Spirin, K. S., Simpson, J. F., Takeuchi, S., Kawamata, N., Miller, C. W., and Koeffler, H. P. *p27/Kip1* mutations found in breast cancer. *Cancer Res.*, 56: 2400–2404, 1996.
 14. Catzavelos, C., Bhattacharya, N., Ung, Y. C., Wilson, J. A., Roncari, L., Sandhu, C., Shaw, P., Yeger, H., Morava-Protzner, I., Kapusta, L., Franssen, E., Pritchard, K. I., and Slingerland, J. M. Decreased levels of the cell-cycle inhibitor *p27^{KIP1}* protein: prognostic implications in primary breast cancer. *Nat. Med.*, 3: 227–230, 1997.
 15. Esposito, V., Baldi, A., De Luca, A., Groger, A. M., Loda, M., Giordano, G. G., Caputi, M., Baldi, F., Pagano, M., and Giordano, A. Prognostic role of the cyclin-dependent kinase inhibitor *p27* in non-small cell lung cancer. *Cancer Res.*, 57: 3381–3385, 1997.
 16. Loda, M., Cukor, B., Tam, S. W., Lavin, P., Fiorentino, M., Draetta, G. F., Jessup, J. M., and Pagano, M. Increased proteasome-dependent degradation of the cyclin-dependent kinase inhibitor *p27* in aggressive colorectal carcinomas. *Nat. Med.*, 3: 231–233, 1997.
 17. Porter, P. L., Malone, K. E., Heagerty, P. J., Alexander, G. M., Gatti, L. A., Firpo, E. J., Daling, J. R., and Roberts, J. M. Expression of cell-cycle regulators *p27* and cyclin E, alone and in combination, correlate with survival in young breast cancer patients. *Nat. Med.*, 3: 222–225, 1997.
 18. Tsihlias, J., Kapusta, L. R., DeBoer, G., Morava-Protzner, I., Zbieranowski, I., Bhattacharya, N., Catzavelos, G. C., Klotz, L. H., and Slingerland, J. M. Loss of cyclin-dependent kinase inhibitor *p27^{KIP1}* is a novel prognostic factor in localized human prostate adenocarcinoma. *Cancer Res.*, 58: 542–548, 1998.
 19. Lane, D. P. *p53*, guardian of the genome. *Nature (Lond.)*, 358: 15–16, 1992.
 20. Momand, J., Zambetti, G. P., Olson, D. C., George, D., and Levine, A. J. The *mdm-2* oncogene product forms a complex with the *p53* protein and inhibits *p53*-mediated transactivation. *Cell*, 26: 1237–1245, 1992.
 21. Esrig, D., Spruck, C. H., Nichols, P., Chaiwun, B., Steven, K., Groshen, S., Chen, S. C., Skinner, D. G., Jones, P. A., and Cote, R. J. *p53* nuclear protein accumulation correlates with mutations in the *p53* gene, tumor grade and stage in bladder cancer. *Am. J. Pathol.*, 143: 1389–1397, 1993.
 22. Sarkis, A. S., Dalbagni, G., Cordon-Cardo, C., Zhang, Z. F., Sheinfeld, J., Fair, W. R., Herr, H. W., and Reuter, V. E. Nuclear overexpression of *p53* protein in transitional cell bladder carcinoma: a marker for disease progression. *J. Natl. Cancer Inst.*, 85: 53–59, 1993.
 23. Mostofi, F. H., Sobin, L. H., and Torloni, H. Histological typing of urinary bladder tumors. *In: International Classification of Tumors*, No. 10. Geneva: World Health Organization, 1973.
 24. Sobin, L. H., and Wittekind, C. *TNM Classification of Malignant Tumors*, Ed. 5. New York: Wiley-Liss, 1997.
 25. Lipponen, P. K. Over-expression of *p53* nuclear oncoprotein in transitional-cell bladder cancer and its prognostic value. *Int. J. Cancer*, 53: 365–370, 1993.
 26. Serth, J., Kuczyk, M. A., Bokemeyer, C., Hervatin, C., Nafe, R., Tan, H. K., and Jonas, U. *p53* immunohistochemistry as an independent prognostic factor for superficial transitional cell carcinoma of the bladder. *Br. J. Cancer*, 71: 201–205, 1995.
 27. Fleiss, J. L. *Statistical methods for rates and proportions*, Ed. 2. New York: John Wiley and sons, 1981.
 28. Pagano, M., Tam, S. W., Theodoras, A. M., Beer-Romero, P., Del Sal, G., Chau, V., Yew, P. R., Draetta, G. F., and Rolfe, M. Role of the Ubiquitin-proteasome pathway in regulating abundance of the cyclin-dependent kinase inhibitor *p27*. *Science (Washington DC)*, 269: 682–685, 1995.
 29. Dulic, V., Kaufmann, W. K., Wilson, S. J., Tlsty, T. D., Lees, E., Harper, J. W., Elledge, S. J., and Reed, S. I. *p53*-dependent inhibition of cyclin-dependent kinase activities in human fibroblasts during radiation-induced *G₁* arrest. *Cell*, 76: 1013–1023, 1994.
 30. El-Deiry, W. S., Tokino, T., Velculescu, V. E., Levy, D. B., Parsons, R., Trent, J. M., Lin, D., Mercer, W. E., Kinzler, K. W., and Vogelstein, B. WAF1, a potential mediator of *p53* tumor suppression. *Cell*, 75: 817–825, 1993.
 31. Polyak, K., Lee, M-H., Erdjument-Bromage, A., Koff, A., Roberts, J. M., Tempst, P., and Massague, J. Cloning of *p27^{KIP1}*, a cyclin-dependent kinase inhibitor and a potential mediator of extracellular antimitogenic signals. *Cell*, 78: 59–66, 1994.
 32. Sherr, C. J., and Roberts, J. M. Inhibitors of mammalian *G₁* cyclin-dependent kinases. *Genes and Dev.*, 9: 1149–1163, 1995.
 33. Bartkova, J., Lukas, J., Muller, H., Lutzhoft, D., Strauss, M., and Bartek, J. Cyclin D1 protein expression and function in human breast cancer. *Int. J. Cancer*, 57: 353–361, 1994.
 34. Jiang, W., Zhang, Y-J., Kahn, S. M., Hollestein, M. C., Santella, R. M., Lu, S. H., Harris, C. C., Montesano, R., and Weinstein, I. B. Altered expression of the cyclin D1 and retinoblastoma genes in human esophageal cancer. *Proc. Natl. Acad. Sci. USA*, 90: 9026–9030, 1993.
 35. Aagaard, L., Lukas, J., Bartkova, J., Kjerulf, A. A., Strauss, M., and Bartek, J. Aberrations of *p16^{Ink4}* and retinoblastoma tumor-suppressor gene occur in distinct sub-sets of human cancer cell lines. *Int. J. Cancer*, 61: 115–120, 1995.
 36. Shapiro, G. I., Edwards, C. D., Kobzik, L., Godleski, J., Richards, W., Sugarbaker, D. J., and Rollins, B. J. Reciprocal Rb inactivation and *p16^{INK4}* expression in primary lung cancer and cell lines. *Cancer Res.*, 55: 505–509, 1995.
 37. Lane, D. P., and Hall, P. A. MDM2-arbiter of *p53*'s destruction. *Trends Biochem. Sci.* 22: 372–374, 1997.
 38. Scheffner, M., Huibregtse, J. M., Vierstra, R. D., and Howley, P. M. The HPV-16 E6 and E6-AP complex functions as a ubiquitin-protein ligase in the ubiquitination of *p53*. *Cell*, 75: 495–505, 1993.
 39. Gardiner, R. A., Walsh, M. D., Allen, V., Rahman, S., Samarantunga, M. L. T. H., Seymour, G. J., and Lavin, M. F. Immunohistological expression of *p53* in primary pT1 transitional cell bladder cancer in relation to tumour progression. *Br. J. Urol.*, 73: 526–532, 1994.
 40. Nakopoulou, L., Constantinides, C., Papandropoulos, J., Theodoropoulos, G., Tzounou, A., Giannopoulos, A., Zervas, A., and Dimopoulos, C. Evaluation of overexpression of *p53* tumor suppressor protein in superficial and invasive transitional cell bladder cancer: comparison with DNA ploidy. *Urology*, 46: 334–340, 1995.
 41. Sgambato, A., Zhang, Y-J., Arber, N., Hibshoosh, H., Doki, Y., Ciaparrone, M., Santella, R. M., Cittadini, A., and Weinstein, I. B. Deregulated expression of *p27^{KIP1}* in human breast cancers. *Clin. Cancer Res.*, 3: 1879–1887, 1997.
 42. Fredersdorf, S., Burns, J., Milne, A. M., Packham, G., Fallis, L., Gillett, C. E., Royds, J. A., Peston, D., Hall, P. A., Hanby, A. M., Barnes, D. M., Shousha, S., O'Hare, M. J., and Lu, X. High level expression of *p27^{KIP1}* and cyclin D1 in some human breast cancer cells: inverse correlation between the expression of *p27^{KIP1}* and degree of malignancy in human breast and colorectal cancers. *Proc. Natl. Acad. Sci. USA*, 94: 6380–6385, 1997.
 43. Ciaparrone, M., Yamamoto, H., Yao, Y., Sgambato, A., Cattoretti, G., Tomita, N., Monden, T., Rotterdam, H., and Weinstein, I. B. Localization and expression of *p27^{KIP1}* in multistage colorectal carcinogenesis. *Cancer Res.*, 58: 114–122, 1997.
 44. Doki, Y., Imoto, M., Han, E. K-H., Sgambato, A., and Weinstein, I. B. Increased expression of the *p27^{KIP1}* protein in human esophageal cancer cell lines that overexpress cyclin D1. *Carcinogenesis (Lond.)*, 18: 1139–1148, 1997.
 45. Kitahara, K., Yasui, W., Yokozaki, H., Semba, S., Hamamoto, T., Hisatsugu, T., and Tahara, E. Expression of cyclin D1, CDK4 and *p27Kip1* is associated with the *p16^{MTS1}* gene status in human esophageal carcinoma cell lines. *J. Exp. Ther. Oncol.* 1: 7–12, 1996.
 46. Shin, K. Y., Kong, G., Kim, W. S., Lee, T. Y., Woo, Y. N., and Lee, J. D. Overexpression of cyclin D1 correlates with early recurrence in superficial bladder cancers. *Br. J. Cancer*, 75: 1788–1792, 1997.
 47. Boring, C. C., Squires, T. S., and Tong T. Cancer Statistics, 1994. *CA Cancer J. Clin.*, 44: 7–26, 1994.

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

Loss of P27^{Kip1} Expression Correlates with Tumor Grade and with Reduced Disease-free Survival in Primary Superficial Bladder Cancers

Alessandro Sgambato, Mario Migaldi, Beatrice Faraglia, et al.

Cancer Res 1999;59:3245-3250.

Updated version Access the most recent version of this article at:
<http://cancerres.aacrjournals.org/content/59/13/3245>

Cited articles This article cites 43 articles, 18 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/59/13/3245.full#ref-list-1>

Citing articles This article has been cited by 11 HighWire-hosted articles. Access the articles at:
<http://cancerres.aacrjournals.org/content/59/13/3245.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerres.aacrjournals.org/content/59/13/3245>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.