

DNA Sequence Copy Number Changes in Gastrointestinal Stromal Tumors: Tumor Progression and Prognostic Significance¹

Wa'el El-Rifai, Maarit Sarlomo-Rikala, Leif C. Andersson, Sakari Knuutila,² and Markku Miettinen

Department of Medical Genetics, Haartman Institute and Helsinki University Central Hospital, University of Helsinki [W. E.-R., S. K.], Department of Pathology, Haartman Institute, University of Helsinki [M. S.-R., L. C. A.], FIN-00029 HUCH Helsinki, Finland; Department of Soft Tissue Pathology, Armed Forces Institute of Pathology, Washington, DC 20306-6000 [M. M.]; and Department of Human Genetics, National Research Center, Cairo, Egypt [W. E.-R.]

ABSTRACT

To identify genetic changes related to tumor progression and find out diagnostic and prognostic genetic markers in gastrointestinal stromal tumors (GISTs), 95 tumor samples (24 benign GISTs, 36 malignant primary GISTs, and 35 GIST-metastases) from 60 patients were studied using comparative genomic hybridization. DNA copy number changes were detected in all samples. Benign GISTs had a mean of 2.6 aberrations/sample (losses:gains, 5:1) and significantly fewer DNA copy number changes and fewer gains than malignant primary and metastatic GISTs ($P < 0.01$). High-level amplifications were not seen in benign GISTs. Malignant primary GISTs had a mean of 7.5 aberrations/tumor (losses:gains, 1.6:1), whereas the mean number of aberrations/metastatic GIST was 9 (losses:gains, 1.8:1). Frequent changes observed in all GIST groups included losses in chromosome arms 1p (51%), 14q (74%), and 22q (53%). Gains and high-level amplifications at 8q and 17q were significantly more frequent in metastatic GISTs (57 and 43%) than in benign GISTs (8 and 0%; $P < 0.001$) and malignant primary GISTs (33 and 25%; $P < 0.05$). Gains and high-level amplifications at 20q were only seen in malignant primary and metastatic GISTs ($P < 0.01$), and gains at 5p were not detected in benign GISTs ($P < 0.01$). Losses in chromosome arm 9p were never seen in benign tumors ($P < 0.001$), and they were more frequent in metastatic GISTs than in malignant primary GISTs (63 and 36%; $P < 0.05$). Losses in 13q were less frequent in benign GISTs than in malignant primary ($P < 0.05$) and metastatic ($P < 0.01$) GISTs. Our results show that several DNA copy number changes are related to the behavior of GISTs and can be used as prognostic markers for tumor progression.

INTRODUCTION

GISTs³, classified previously as smooth muscle tumors, constitute the most important group of primary mesenchymal tumors of the gastrointestinal tract. These tumors occur at all levels of the gastrointestinal tract and usually present between the sixth and eighth decades. Immunohistochemically, GISTs are usually positive for KIT (CD117) and CD34, variably positive for smooth muscle actin, and usually negative for desmin, in contrast to true smooth muscle tumors. GISTs are negative for S100 protein, in contrast to schwannomas (1, 2).

Clinically and pathologically, GISTs represent a spectrum of tumors including benign and malignant variants, the latter of which are generally identified by mitotic activity (>1 mitosis/10 high-power field). In some cases, the prediction of biological potential is difficult, because larger tumors with lower mitotic activity may also occasionally metastasize.

CGH enables screening of entire tumor genomes for gains and losses of DNA copy number and consequent mapping of aberrations to chromosomal subregions (reviewed in Refs. 3 and 4). Recently, we reported using CGH that loss of genetic material at chromosome arm 14q is the most frequently occurring aberration in both benign and malignant primary GISTs (5). The DNA copy number changes seen in GISTs were not detected in leiomyomas and leiomyosarcomas (6). Therefore, the immunophenotypic characteristics and the genetic profile of GISTs have clearly placed it as a separate tumor entity different from other mesenchymal tumors of the gastrointestinal tract. However, the prognostic evaluation of GISTs remained a difficult issue, requiring a complex multiparametric approach.

This study was performed to analyze the CGH changes in benign and malignant primary GISTs and to investigate whether progressive DNA copy number changes occur in recurrent and metastatic GISTs. We also investigated whether any DNA copy number change has prognostic significance for GISTs.

MATERIALS AND METHODS

Tumors. Ninety-five GIST samples from 60 patients were included in the study. The samples consisted of 24 benign GISTs, 36 malignant primary GISTs, and 35 GIST metastases that were available from 15 of the malignant primary GISTs. Immunohistochemically identified leiomyomas and leiomyosarcomas (desmin-positive, CD117-negative) and schwannomas (S100-protein positive, CD117-negative) were excluded from the study. Most GISTs were CD117 and CD34 positive (92 and 72%, respectively). The categorization of GISTs into benign and malignant tumors was based on mitotic counts. Benign GISTs had <2 mitoses/10 high-power field. Malignant or potentially malignant GISTs had >1 mitosis/10 high-power fields. Follow-up data for 6–209 months (mean, 41 months) were available for all patients except four (nos. 25, 26, 29, and 48). None of the patients with benign tumors showed recurrence or metastases.

CGH. Because CGH sensitivity requires at least 50% of tumor material within a sample, paraffin-embedded tissue sections were dissected to obtain at least 70% of tumor cells. DNA from paraffin-embedded tissue sections was extracted as described earlier (7).

CGH was performed according to standard procedures with a modification using a mixture of fluorochromes conjugated to dCTP and dUTP nucleotides for nick translation (8). Hybridizations, washings, and ISIS digital image analysis (Metasystems GmbH, Altlussheim, Germany) were performed as described elsewhere (5).

Controls. In each CGH experiment, a negative control (peripheral blood DNA from a healthy donor) and a positive control were included. The positive control was a gastric tumor with known DNA copy number changes. On the basis of our earlier reports and the control results, we used 1.17 and 0.85 as cutoff levels for gains and losses, respectively. All of the CGH results were confirmed using a 99% confidence interval.

Statistical Analysis. All of the CGH results were confirmed using a 99% confidence interval. Briefly, intra-experiment SDs for all positions in the CGH ratio profiles were calculated from the variation of the ratio values of all homologous chromosomes within the experiment. Confidence intervals for the ratio profiles were then computed by combining them with an empirical inter-experiment SD and by estimating error probabilities based on the t distribution. For the analysis of the frequencies of DNA copy number changes in primary and metastatic GISTs, we used Fisher's exact two-tailed test. $P_s < 0.05$ were considered significant.

Received 12/30/99; accepted 5/17/00.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported by grants from the Finnish Cancer Society and the University of Helsinki in Finland.

² To whom requests for reprints should be addressed, at Department of Medical Genetics, Helsinki University Central Hospital, P. O. Box 404 (Haartmanink. 3, 4th floor), FIN-00029 HUCH, Helsinki, Finland. Phone: 358-9-1912-6527; Fax: 358-9-1912-6788; E-mail: Sakari.Knuutila@Helsinki.FI.

³ The abbreviations used are: GIST, gastrointestinal stromal tumor; CGH, comparative genomic hybridization.

Table 1 Clinical, histological, and CGH karyotype findings of 95 gastrointestinal stromal tumors obtained from 60 patients

Metastases from the same patient have the same number as the primary tumor but with a different letter, e.g. b, c, d.

No.	Age/sex/ code	Histology	Site ^a	Size ^b	Follow-up in months	CGH karyotype ^c	
						Losses	Gains and high-level amplifications
1	59/m G1	Benign	Jejunum	5, 5	NED, 51	1p, 15, 18, 22	
2	45/m G2	Benign	Stomach	4, 5	NED, 45	14	
3	50/m G12	Benign	Stomach	3, 3	NED, 38	1p, 14	8q
4	75/m G25	Benign	Stomach	3	NED, 87	14, 22	
5	63/m G9	Benign	Stomach	3, 5	NED, 41	14q11-q13, 14q23-qter, 22	
6	61/f G10	Benign	Stomach	7	NED, 40	1p, 14	
7	81/m G13	Benign	Stomach	5, 5	NED, 110	15	
8	80/m G14	Benign	Stomach	5	NED, 124	14	
9	9/73/f G18	Benign	Stomach	3, 5	NED, 81	11p, 14	5q22-qter
10	63/f G19	Benign	Jejunum	2	NED, 31	1p, 11, 14, 15, 21	1q, 5, 7p, 18q
11	58/m G21	Benign	Stomach	4	NED, 210	14q13-qter, 22	
12	49/f G23	Benign	Stomach	16	NED, 209	22q12-qter	
13	65/f G24	Benign	Stomach	3	NED, 28	14, Xq	
14	73/m G26	Benign	Stomach	2	NED, 82	14, 15, 21, 22	
15	79/f G29	Benign	Stomach	2	NED, 133	22q12-qter	
16	78/m G32	Benign	Stomach	2, 5	NED, 75	14, 22	
17	80/m G35	Benign	Jejunum	4, 5	NED, 24	1p, 6q, 13q21-qter, 14	19
18	85/m G38	Benign	Stomach	5	NED, 21	14, 22	
19	69/m G41	Benign	Jejunum	2	NED, 14	1p, 11p, 13, 14q11-q12, 15, 22	
20	67/f G44	Benign	Stomach	9	NED, 6	14	
21	53/m G48	Benign	Stomach	6	NED, 31	14	
22	34/m G49	Benign	Stomach	6, 5	NED, 32		7, 8, 11
23	41/m G52	Benign	Jejunum	7	NED, 13	1p31.2-pter, 6q, 22	
24	68/m G60	Benign	Stomach	2	NED, 33	14	
25	74/f G15	Malignant	Stomach	NA	NA	14q13-qter, 15q15-qter, 18	16, 17
26	60/f G22	Malignant	Stomach	4	NA	10q	3q, 5, 7, 12, 13, 18
27	59/m G47	Malignant	Colon	18	NED, 12	1p, 6q, 8p22-pter, 10, 11q14-qter, 12, 14, 15, 18q11-q23	2q22-qter
28	52/m G6	Malignant	Stomach	15	DOD, 5	3cen-p21, 14	1q32-qter, 3p22-pter, 3q (3q26-q29) 8p12-qter
29	72/m G17	Malignant	Jejunum	15	NA	2q31-qter, 3p, 9p	
30	60/m G20	Malignant	Jejunum	10	DOD, 9	1p, 13, 14cen-q23, 15, 21, 22	3q, 8, 16p, 17
31	68/f G27	Malignant	Stomach	15	DOD, 71	1cen-p31, 3p, 14q21-q23, 15q21-qter	2q34-q37
32	88/f G31	Malignant	Stomach	6	DOD, 26	1cen-p31, 8p, 9q, 10p15-q22, 22, X	1p32-pter, 5p, 9p, 17q21-q25
33	63/m G36	Malignant	Colon	16	NED, 24	8p21-pter, 10pter-q22, 14q21-q32	1p, 5, 7p, 8q22-qter, 9q, 19, 20
34	63/m G11	Malignant	Rectum	7	NED, 39	8p, 9, 13q21-qter, 14, 15q15-q22, 22	3, 5, 8q, 11, 12, 17, 18, X
35	86/f G16	Malignant	Small intestine	6, 5	DOD, 19	1p, 2p, 15	
36	43/f G28	Malignant	Jejunum	8	NED, 78	1cen-p31, 13q21-qter, 15q15-qter, 18q	4, 17, 18p, 19
37	69/m G30	Malignant	Stomach	11	DOD, 32		11q14-q22, 19, X
38	41/m G42	Malignant	Stomach	18	NED, 12	1p, 2p13-pter, 9p, 14, 15, 22	11q
39	44/m G43	Malignant	Stomach	17	NED, 9	22	5, 7
40	73/f G45	Malignant	Stomach	4	NED, 8		5, 6p, 7
41	71/f G46	Malignant	Jejunum	9	NED, 27	1p, 4, 10, 15, 22	
42	73/f G50	Malignant	Jejunum	9	Alive, 9	1p, 2q, 9p, 13, 14, 15, 18q, 22	5p14-pter
43	59/m G58	Malignant	Stomach	12	NED, 27	9, 14	4p, 5, 7
44	37/f PR28	Malignant	Small intestine	NA	DOD, 50	1p, 8p, 9p, 11p, 14, 18q, 22	1q11-q31, 8q, 9q, 16q, 19p, 20q
45	73/f G65	Malignant	Duodenum	6, 5	NED, 24	1p, 6q21-qter, 9p, 11, 13, 14, 15	7
46	46/f K1	Malignant	Stomach	10	Alive, 35	7p15-p22, 7q32-qter, 9p21.2-pter, 14, 15, 18p, 22q13	2q, 4q, 8q23-qter
46b	K2	Metastases	Omentum			7p15-p22, 7q32-qter, 9p21.2-pter, 13q13-qter, 14, 15, 19q13.2-qter, 22q13	2q, 6p, 8q23-qter, 17q
46c	K3	Metastases	Colon mesentery			7p15-pter, 7q32-qter, 9p21.2-pter, 14, 15, 19q13.2-qter, 22q13	2q, 4q, 8q23-qter, 17q
46d	K4	Metastases	Omentum			4p, 7p15-p22, 7q32-qter, 9p21.2-pter, 10, 11, 13q13-qter, 14, 15, 19q13.2-qter, 22q13	8q23-qter, 17q
46e	K5	Metastases	Liver			4p, 7p15-p22, 7q32-qter, 9p21.2-pter, 11p, 13q13-qter, 14, 15, 19q13.2-qter, 22q13	2q, 8q23-qter, 17q
46f	K6	Metastases	Liver			1p34.3-pter, 4, 6q14-qter, 7p15-p22, 7q32-qter, 9, 14, 18, 22q13	1q, 5
46g	K7	Metastases	Colon mesentery			7p15-p22, 7q32-qter, 9p21.2-pter, 14, 15, 18p, 19q13.2-qter, 22q13	2q, 4q, 8q23-qter, 17q
46h	K8	Metastases	Colon mesentery			7p15-p22, 7q32-qter, 9p21.2-pter, 13q13-qter, 14, 15, 19q13.2-qter, 22q13	2pter-q13, 4q, 8q22-qter, 17q
46i	K9	Metastases	Colon mesentery			1p, 7p15-p22, 7q32-qter, 9p21.2-pter, 11, 13q, 14, 15, 19q13.2-qter, 22q13, X	2q, 4q, 8q22-qter (8q23-qter), 17q
46j	K10	Metastases	Colon mesentery			7p15-p22, 7q32-qter, 9p21.2-pter, 13q13-qter, 14, 15, 19q13.2-qter, 22q13	2pter-q13, 4q, 8q22-qter, 17q
46k	K11	Metastases	Ovary			7p15-p22, 7q32-qter, 9p21.2-pter, 11, 13, 14, 15, 19q13.2-qter, 22q13, X	2q, 4q, 8q22-qter, 17q
46l	K12	Metastases	Colon mesentery			7p15-p22, 7q32-qter, 9p21.2-pter, 13q13-qter, 14, 15, 19q13.2-qter, 22q13	2pter-q13, 8q23-qter, 17q
46m	K14	Metastases	Colon mesentery			6q14-qter, 7p11-p15, 7q32-qter, 9pter-q34.1, 14, X	
47	67/m PR1	Malignant	Jejunum	7	DOD, 35	1p, 2, 11, 13, 14, 15, 18	
47b	PR2	Metastases	Abdomen			1p, 2, 11, 13, 14, 15	1q, 3, 5, 6p, 7, 8, 9
47c	PR48	Metastases	Abdomen			1p, 2, 11, 13, 14, 15, 18	1q11-q24, 3p14-p22, 6p, 7, 8
47d	PR3	Metastases	Abdomen			1p, 2p22-pter, 11, 13, 14, 15, 18	1q, 3, 6p, 7, 8, 9
48	44/m G8	Malignant	Stomach	17	NA	1q32-qter, 14	8q22-qter
48b	PR5	Metastases	Abdomen			1p, 2q, 5p14.3-pter, 12q15-qter, 14, 19q	8
49	64/m PR6	Malignant	Small intestine	12	Alive, 24	1p, 6q, 13, 14, 15, 18, 22	5p, 6p, 7, 12q
49b	PR7	Metastases	Liver			1p, 6q, 13, 14, 15, 18, 22	5p, 6p, 7, 12q
50	59/f PR8	Malignant	Stomach	15	Alive, 96	9pter-q31, 10p, 13, 14, 19q13.3-qter, X	2p14-pter, 4p15-pter, 6p, 8, 9q32-qter, 10q, 12, 17q, 19pter-q13.2, 20q
50b	PR9	Metastases	Liver			9pter-q31, 10p, 13, 14, 19q13.3-qter, X	2p14-pter, 7, 8, 12, 17, 19pter-q13.2, 20q

Table 1 Continued.

No.	Age/sex/ code	Histology	Site ^a	Size ^b	Follow-up in months	CGH karyotype ^a	
						Losses	Gains and high-level amplifications
51	54/m PR10	Malignant	Jejunum	15	Alive, 22	1p, 2, 7p, 11, 15, 18, 22	4q, 7q, 9p, 12q
51b	PR11	Metastases	Abdomen			1p, 2, 4p, 11, 12p, 13, 15, 17p, 18, 22	5q, 7q, 8q, 9p , 21
51c	PR12	Metastases	Abdomen			1p, 2p, 13, 15, 18, 22	5q, 10p
52	59/m PR13	Malignant	Small intestine	20	DOD, 49	1p, 8, 13, 14, 15, 22	1q, 2q22-qter, 4p, 5p, 17, 20p
52b	PR14	Metastases	Abdomen			1p, 3pter-q13, 4q, 8, 9pter-q21, 10, 13, 14, 15, 22	1q, 2q22-qter, 4p, 5p, 17, 20p
52c	PR15	Metastases	Abdomen			1p, 3pter-q13, 4q, 8, 9pter-q21, 13, 14, 15, 22	1q, 2q22-qter, 5p, 17, 20p
53	62/f PR16	Malignant	Stomach	15	Alive, 60	6, 9, 12pter-q22, 14, 22	17
53b	PR17G5	Metastases	Abdomen		Alive	9, 14, 22	17q
53c	PR18	Metastases	Abdomen			9, 14, 22	4, 17q, 20
54	59/m PR19G3	Malignant	Ileum	12	DOD, 14	13, 14q11-q12	5p
54b	PR20	Metastases	Liver			12p, 14	5p, 8q
55	38/f PR21	Malignant	Jejunum	10	DOD, 59	1p, 11q14-qter, 13, 14, 15, 18, 22, X	1q, 5p, 8
55b	PR22	Metastases	Abdomen			1p, 15, 18, 22q13	1q, X
55c	PR23	Metastases	Abdomen			1p, 15, 18, 22q13	1q
55d	PR24	Metastases	Abdomen			1p, 13, 15, 18, 22	1q
56	52/f PR33	Malignant	Small intestine	10	DOD, 12	1p, 14, 15	7, 8, 12q22-qter
56b	PR34	Metastases	Abdomen			1p, 14, 15	7, 8, 12q22-qter
57c	PR35	Metastases	Abdomen			1p, 14, 15	7, 8, 12q22-qter
58	61/m PR36	Malignant	Stomach	25	DOD, 19	1p, 11, 13, 14, 15	2q, 8q23-qter, 17q, X
58b	PR37	Metastases	Abdomen			1p, 9	20
58c	PR38	Metastases	Abdomen			1p, 9	17q22-qter, 20
59	39/m PR39	Malignant	Stomach	20	DOD, 14	4q27-q34, 9, 14, 22	
59b	PR40	Metastases	Abdomen			4q27-q34, 9, 14, 22	
59c	33/m PR42	Metastases	Abdomen			1p11-p32, 4, 9pter-q33, 14	20q13.2-qter
60	41/f PR45	Malignant	Stomach	6	Alive, 6	1p35-pter, 4, 9, 15, 22	5, 8p12-pter, 8q23-qter, 12q14-qter, 13, 18q
60b	PR46	Metastases	Abdomen			1p35-pter, 4, 6q, 9, 14q11-q21, 15, 22	5, 8p12-pter, 8q23-qter, 12 (12q22-qter), 13, 18q, 20q
60c	PR47	Metastases	Abdomen			1p35-pter, 4, 9, 10, 14, 15, 22	5p, 5q32-qter, 8p12-pter, 8q23-qter, 12 (12q22-qter), 13, 20q

^a All metastases were intra-abdominal.

^b Maximum size in cm, high.

^c High-level amplifications are in bold; NED, no evidence of disease; NA, not available; DOD, died of disease.

RESULTS

Changes in DNA copy numbers were detected in all samples. DNA copy number changes were seen in all chromosomes in malignant primary and metastatic GISTs, whereas fewer chromosomes were involved in benign GISTs (Figs. 1 and 2). Benign GISTs had a mean of 2.6 aberrations/case (losses:gains, 5:1) and contained significantly fewer DNA copy number changes and fewer gains than malignant primary and metastatic GISTs ($P < 0.01$). High-level amplifications were not seen in benign GISTs. In malignant primary GISTs, the mean number of aberrations/case was 7.5 (losses:gains, 1.6:1), whereas the mean was 9 in metastatic GISTs (losses:gains, 1.8:1). The CGH profiles in metastatic GISTs and malignant primary tumors were similar, but metastatic GISTs showed a larger number of high-level amplifications. There was no correlation between the location and any specific DNA copy number changes.

Table 2 Summary of significant DNA copy number changes in GISTs

DNA copy number changes seen in primary malignant and metastatic tumors were compared to benign lesions. *Ps* are shown in parentheses. *Ps* shown in italics indicate comparison between metastatic and primary malignant tumors. Gains are indicated with + and losses with -.

Chromosomal region	Benign GIST	Malignant GIST	Metastatic GIST
8q+	8%	33% (<0.001)	57% (<0.001), (<0.05)
17q+	0	25% (<0.05)	43% (<0.001), (<0.05)
20q+	0	11% (<0.001)	26% (<0.001), (<0.05)
5p+	0	31% (<0.001)	29% (<0.001)
9p-	0	36% (<0.001)	63% (<0.001), (<0.05)
13q-	8%	36% (<0.05)	46% (<0.001)

Changes that were frequently detected in all GISTs with no statistical differences between benign, malignant primary, and metastatic tumors included losses in chromosome arms 1p (51%), 14q (74%), and 22q (53%).

Gains and high-level amplifications at 8q and 17q were significantly more often detected in metastatic GISTs (57 and 43%) than in benign GISTs (8 and 0%; $P < 0.001$) and malignant primary GISTs (33 and 25%; $P < 0.05$). Gains and high-level amplifications at 20q were only seen in malignant primary and metastatic GISTs ($P < 0.01$), and gains at 5p were not detected in benign GISTs ($P < 0.01$). Losses in 9p were never seen in benign tumors ($P < 0.001$), and they were more frequent in metastatic GISTs than in malignant primary tumors (63 and 36%; $P < 0.05$). The losses in 13q were less frequent in benign GISTs than in malignant primary ($P < 0.05$) and metastatic ($P < 0.01$) GISTs. In addition, several other changes were seen more frequently in malignant primary and metastatic GISTs than in benign GISTs. The details of DNA copy number changes are shown in Table 1 and Fig. 1. Fig. 2 shows the relative frequencies of the aberrations in all GISTs, and Table 2 summarizes the significant DNA copy number changes.

DISCUSSION

On the basis of a large series of clinically benign and malignant GISTs and their metastases, we investigated the possible value of CGH to identify genetic markers of tumor behavior. The observation that benign tumors, almost exclusively, contain losses rather than gains, whereas malignant primary GISTs and their metastases contain




Fig. 1. Summary of DNA copy number gains and losses detected by CGH in 60 GIST patients. Each bar represents one patient, regardless of the number of samples available, and the minimal common overlapping regions are presented. Gains are on the right hand side, losses on the left. Green broken lines, benign tumors; green continuous lines, malignant primary tumors that had no metastases samples. Blue lines, the aberration was seen in the malignant primary tumor and in at least one metastatic sample from the same patient. Red lines, changes that were seen on metastatic samples but not in their primary tumors for the same patient.




Fig. 2. Comparative frequencies of losses (left) and gains (right) detected in 24 benign GISTs (green), 36 malignant primary GISTs (blue), and 35 metastatic GISTs (red). Chromosomal numbers are shown in the middle, and the minimal common overlapping regions, when applicable, are added to each bar.

gains and high-level amplifications, presents an apparently clear criterion for categorization of these tumors. Accordingly, losses are more likely related to the development of GISTs, whereas accumulation of additional genetic alterations, especially gains/amplifications, is required for malignant transformation and metastatic behavior in GISTs. Absence of gains correlated with benign GISTs with verified good prognosis, whereas presence of gains was more frequent in malignant primary GISTs and their metastases ($P < 0.01$). Therefore, absence of gains can be considered as a good prognostic parameter, which can be used as a new complementary diagnostic criterion for characterization of GISTs.

Gains and high-level amplifications at 5p, 8q, 17q, and 20q and losses in 9p, 13q, 15q, and 19q correlated with malignant primary GISTs and metastatic GISTs. Moreover, the gains at 8q, 17q, and 20q and the losses in 9p were more frequent in metastatic GISTs than in malignant primary GISTs ($P < 0.05$). Therefore, clones with this genetic make-up have a potential capability of developing metastases.

Losses in 1p, 14q, and 22q seen in benign GISTs seem to be unique because they have been reported rarely in other tumors at frequencies as high as observed in GISTs (9). These changes were frequently seen in benign GISTs, indicating that they may be early events in GIST development. The fact that these losses were maintained with tumor progression suggests that they are required for the maintenance of GIST phenotype. The present extended study also supports the notion that these changes may be unique for GIST development. Recently, we reported high-resolution deletion mapping of chromosome 14 in GISTs and identified two possible tumor suppressor loci in 14q11.2 and 14q23 (10).

The gains/high-level amplifications at 5p, 8q, 17q, and 20q as well as the losses in 9p, 13q, 15q, and 19q were detected in many malignant GISTs and their metastases. Such changes have also been reported in several other tumors, such as carcinomas of lung, breast, and ovary, and in squamous cell carcinomas of the head with variable frequencies (reviewed in Refs. 3 and 4). Gains and high-level amplifications at 8q22–q24 and 17q21–qter have been implicated as poor prognostic indicators in breast cancer, ovarian cancer, osteosarcomas, and neuroblastomas (11–14). The gains and high-level amplifications at 8q and 20q correlate with invasiveness in breast cancer (15, 16).

Although these chromosomal regions are known to contain several oncogenes, such as *CMYC* at 8q and *ERBB2* at 17q, the possibility of yet undiscovered oncogenes cannot be ruled out (14). At 20q, several genes have been implicated in breast cancer, and the region is known to harbor specific amplified genes (*AIB1*, *AIB3*, and *AIB4*; Ref. 17). Several candidate genes are located at 20q, e.g., the *PTP1B/PTPN1* gene (20q12), which is involved in growth regulation, and the *MYBL2* gene (20q13), which plays an important role in cell cycle progression. Moreover, the human *cellular apoptosis susceptibility* (*CAS*) gene has been mapped to this same region (18). There are no known oncogenes that map to 5p.

Losses in 9p were associated with recurrences and unfavorable outcome in squamous cell carcinoma of the head and neck (19) and in astrocytic tumors (20). Losses in 13q12–q13 are associated with poor prognosis in familial and sporadic breast cancer (21). Losses in 15q have been observed rarely upon CGH. LOH studies have suggested the presence of a putative tumor suppressor gene in 15q in small cell lung cancer (22). However, there are no data related to its potential prognostic significance or identified tumor suppressor genes.

In summary, our results show that genetic changes can be used as a complementary diagnostic tool for the prognostication of GISTs and for the differentiation between benign and malignant tumors. The increased number of changes and/or increased number of gains correlate with malignant behavior. Furthermore, the genetic changes seen

as gains at 5p, 8q, 17q, and 20q and losses in 9p, 13q, 15q, and 19q are new prognostic parameters for GISTs.

REFERENCES

- Sarlomo-Rikala, M., Kovatich, A., Barusevicius, A., and Miettinen, M. CD117: a sensitive marker for gastrointestinal stromal tumors that is more specific than CD34. *Mod. Pathol.*, *11*: 728–734, 1998.
- Miettinen, M., Sarlomo-Rikala, M., and Lasota, J. Gastrointestinal stromal tumors: recent advances in understanding of their biology. *Hum. Pathol.*, *30*: 1213–1220, 1999.
- Knuutila, S., Björkqvist, A.-M., Autio, K., Tarkkanen, M., Wolf, M., Monni, O., Szymanska, J., Larramendy, M. L., Tapper, J., Pere, H., El-Rifai, W., Hemmer, S., Wasenius, V.-M., Vidgren, V., and Zhu, Y. DNA copy number amplifications in human neoplasms. Review of comparative genomic hybridization studies. *Am. J. Pathol.*, *152*: 1107–1123, 1998.
- Knuutila, S., Aalto, Y., Autio, K., Björkqvist, A.-M., El-Rifai, W., Hemmer, S., Huhta, T., Kettunen, E., Kiuru-Kuhlefelt, S., Larramendy, M. L., Lushnikova, T., Monni, O., Pere, H., Tapper, J., Tarkkanen, M., Varis, A., Wasenius, V.-M., Wolf, M., and Zhu, Y. DNA copy number losses in human neoplasms. *Am. J. Pathol.*, *155*: 683–694, 1999.
- El-Rifai, W., Sarlomo-Rikala, M., Miettinen, M., Knuutila, S., and Andersson, L. C. DNA copy number losses in chromosome 14: an early change in gastrointestinal stromal tumors. *Cancer Res.*, *56*: 3230–3233, 1996.
- El-Rifai, W., Sarlomo-Rikala, M., Knuutila, S., and Miettinen, M. DNA copy number changes in development and progression in leiomyosarcoma of soft tissues. *Am. J. Pathol.*, *153*: 985–990, 1998.
- Miller, S. A., Dykes, D. D., and Polesky, H. F. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res.*, *16*: 1215, 1988.
- El-Rifai, W., Larramendy, M. L., Björkqvist, A.-M., Hemmer, S., and Knuutila, S. Optimization of comparative genomic hybridization using fluorochrome conjugated to dCTP and dUTP nucleotides. *Lab. Invest.*, *77*: 699–700, 1997.
- Mitelman, F. *Catalog of Chromosome Aberrations in Cancer*, Ed. 4, 1987. New York: Wiley-Liss, Inc., 1991.
- El-Rifai, W., Sarlomo-Rikala, M., Miettinen, M., Andersson, L. C., and Knuutila, S. High-resolution deletion mapping of chromosome 14 in stromal tumors of the gastrointestinal tract suggests two distinct tumor suppressor loci. *Genes Chromosomes Cancer*, *27*: 387–391, 2000.
- Bown, N., Cotterill, S., Lastowska, M., O'Neill, S., Pearson, A. D., Plantaz, D., Meddeb, M., Danglot, G., Brinkschmidt, C., Christiansen, H., Laureys, G., and Speleman, F. Gain of chromosome arm 17q and adverse outcome in patients with neuroblastoma. *N. Engl. J. Med.*, *340*: 1954–1961, 1999.
- Tarkkanen, M., Elomaa, I., Blomqvist, C., Kivioja, A. H., Kellokumpu-Lehtinen, P., Bohling, T., Valle, J., and Knuutila, S. DNA sequence copy number increase at 8q: a potential new prognostic marker in high-grade osteosarcoma. *Int. J. Cancer*, *84*: 114–121, 1999.
- Arnold, N., Hagele, L., Walz, L., Schempp, W., Pfisterer, J., Bauknecht, T., and Kiechle, M. Overrepresentation of 3q and 8q material and loss of 18q material are recurrent findings in advanced human ovarian cancer. *Genes Chromosomes Cancer*, *16*: 46–54, 1996.
- Bärlund, M., Tirkkonen, M., Forozan, F., Tanner, M. M., Kallioniemi, O., and Kallioniemi, A. Increased copy number at 17q22–q24 by CGH in breast cancer is due to high-level amplification of two separate regions. *Genes Chromosomes Cancer*, *20*: 372–376, 1997.
- Tanner, M. M., Tirkkonen, M., Kallioniemi, A., Holli, K., Collins, C., Kowbel, D., Gray, J. W., Kallioniemi, O. P., and Isola, J. Amplification of chromosomal region 20q13 in invasive breast cancer: prognostic implications. *Clin. Cancer Res.*, *1*: 1455–1461, 1995.
- Isola, J. J., Kallioniemi, O. P., Chu, L. W., Fuqua, S. A., Hilsenbeck, S. G., Osborne, C. K., and Waldman, F. M. Genetic aberrations detected by comparative genomic hybridization predict outcome in node-negative breast cancer. *Am. J. Pathol.*, *147*: 905–911, 1995.
- Tanner, M. M., Tirkkonen, M., Kallioniemi, A., Isola, J., Kuukasjärvi, T., Collins, C., Kowbel, D., Guan, X. Y., Trent, J., Gray, J. W., Meltzer, P., and Kallioniemi, O. P. Independent amplification and frequent co-amplification of three nonsyntenic regions on the long arm of chromosome 20 in human breast cancer. *Cancer Res.*, *56*: 3441–3445, 1996.
- Brinkmann, U., Gallo, M., Polymeropoulos, M. H., and Pastan, I. The human *CAS* (*cellular apoptosis susceptibility*) gene mapping on chromosome 20q13 is amplified in BT474 breast cancer cells and part of aberrant chromosomes in breast and colon cancer cell lines. *Genome Res.*, *6*: 187–194, 1996.
- Lydiatt, W. M., Davidson, B. J., Schantz, S. P., Caruana, S., and Chaganti, R. S. 9p21 deletion correlates with recurrence in head and neck cancer. *Head Neck*, *20*: 113–118, 1998.
- Maruno, M., Yoshimine, T., Muhammad, A. K., Tokiyoshi, K., and Hayakawa, T. Loss of heterozygosity of microsatellite loci on chromosome 9p in astrocytic tumors and its prognostic implications. *J. Neuro-Oncol.*, *30*: 19–24, 1996.
- Eiriksdottir, G., Johannesdottir, G., Ingvarsson, S., Bjornsdottir, I. B., Jonasson, J. G., Agnarsson, B. A., Hallgrímsson, J., Gudmundsson, J., Egilsson, V., Sigurdsson, H., and Barkardottir, R. B. Mapping loss of heterozygosity at chromosome 13q: loss at 13q12–q13 is associated with breast tumour progression and poor prognosis. *Eur. J. Cancer*, *34*: 2076–2081, 1998.
- Stanton, S. E., Shin, S. W., Johnson, B. E., and Meyerson, M. Recurrent allelic deletions of chromosome arms 15q and 16q in human small cell lung carcinomas. *Genes Chromosomes Cancer*, *27*: 323–331, 2000.

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

DNA Sequence Copy Number Changes in Gastrointestinal Stromal Tumors: Tumor Progression and Prognostic Significance

Wa'el El-Rifai, Maarit Sarlomo-Rikala, Leif C. Andersson, et al.

Cancer Res 2000;60:3899-3903.

Updated version Access the most recent version of this article at:
<http://cancerres.aacrjournals.org/content/60/14/3899>

Cited articles This article cites 20 articles, 4 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/60/14/3899.full#ref-list-1>

Citing articles This article has been cited by 27 HighWire-hosted articles. Access the articles at:
<http://cancerres.aacrjournals.org/content/60/14/3899.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerres.aacrjournals.org/content/60/14/3899>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.