

Celecoxib Prevents Tumor Growth *in Vivo* without Toxicity to Normal Gut: Lack of Correlation between *in Vitro* and *in Vivo* Models¹

Christopher S. Williams, Alastair J. M. Watson, Hongmiao Sheng, Rania Helou, Jinyi Shao, and Raymond N. DuBois²

Departments of Medicine [C. S. W., H. S., R. H., J. S., R. N. D.], and Cell Biology, [C. S. W., R. N. D.], The Vanderbilt Cancer Center, Vanderbilt University Medical Center, Nashville, Tennessee 37232-2279; Veterans Administration Medical Center [R. N. D.], Nashville, Tennessee 37232-2279; and Department of Medicine, University of Liverpool, Liverpool, United Kingdom [A. J. M. W.]

ABSTRACT

Nonsteroidal anti-inflammatory drugs have potential for use in the prevention and/or treatment of colorectal cancer. We have studied the cytotoxic effect of a specific COX-2 inhibitor, celecoxib, against LLC, HCA-7, and HCT-15 cells grown in cell culture and have compared these results with its effect on HCA-7 cells grown as xenografts in nude mice. "High-dose" celecoxib (>20 μM) reduced the viability of all three cell lines *in vitro* as measured by the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide assay. Flow cytometric analysis demonstrated that this loss of viability was attributable to the induction of apoptosis. Significantly, concentrations of the drug <10 μM had no effect on cell viability *in vitro*. The cytotoxic effects of high-dose celecoxib were independent of COX-2 inhibition because similar effects were observed in *cox-2* (+/+), *cox-2* (+/-) and *cox-2* (-/-) fibroblasts. A plasma concentration of $2.3 \pm 0.7 \mu\text{M}$ was achieved when celecoxib (1250 mg/kg of chow) was fed to animals *ad libitum*. Despite a lack of toxicity at 2–3 μM celecoxib *in vitro*, there was attenuation of HCA-7 xenograft growth *in vivo*. Celecoxib had no effect on apoptosis, cell division, or the epithelial architecture of the normal gut in treated mice. These results support the need for additional clinical evaluation of celecoxib for treatment and/or prevention of colorectal cancer in humans.

INTRODUCTION

Colorectal cancer remains a significant health concern for much of the industrialized world. Diagnosis often occurs at a late stage in the progression of this disease, which reduces the likelihood of effective treatment. Current treatment strategies often involve a combination of surgical resection and adjuvant chemotherapy. Because of the unsatisfactory outcome of present treatment methods, especially with advanced disease, much emphasis has been placed on developing better treatment and prevention measures.

Numerous epidemiological studies indicate that chronic use of NSAIDs³ lowers the mortality rate from colorectal cancer (1, 2). NSAIDs are effective at inducing regression of existing polyps in familial adenomatous polyposis patients (3) and in reducing the tumor burden in three animal models of colorectal cancer: the multiple intestinal neoplasia mouse (4, 5), the azoxymethane-treated rat model (6), and the nude mouse xenograft model (7–9).

Received 5/31/00; accepted 9/1/00.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ This work was supported in part by USPHS Grants RO1DK-47297 (to R. N. D.), P30CA-68485 (to R. N. D.) and PO1CA-77839 (to R. N. D.). A. J. M. W. is supported by Grant 97/21 from the Association of International Cancer Research. R. N. D. is a recipient of a Veterans Affairs Research Merit Grant and the Mina C. Wallace Professor of Medicine.

² To whom requests for reprints should be addressed, at Department of Medicine/GI; MCN C-2104, Vanderbilt University Medical Center, 1161 21st Avenue South, Nashville, TN 37232-2279. Phone: (615) 343-4747; Fax: (615) 343-6229; E-mail: raymond.dubois@mcm.vanderbilt.edu.

³ The abbreviations used are: NSAID, nonsteroidal anti-inflammatory drug; LLC, Lewis lung carcinoma; MEF, mouse embryo fibroblast; TUNEL, terminal deoxynucleotidyl transferase-mediated nick end labeling; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; FBS, fetal bovine serum; PGE₂, prostaglandin E₂.

NSAIDs inhibit the activity of the cyclooxygenases, which are key enzymes in the conversion of arachidonate to PGH₂, the immediate substrate for a number of specific prostaglandin synthases. Unfortunately, prolonged use of NSAIDs can result in gastrointestinal ulceration and bleeding. It is widely believed that this ulcerogenic activity of NSAIDs is attributable to the chronic inhibition of prostaglandin production in the gastric mucosa. Accordingly, researchers have tried to identify NSAID derivatives that retain anti-neoplastic activity but do not affect prostaglandin production in gastric mucosa.

There are two isoforms of cyclooxygenase, COX-1 and COX-2, which differ in their expression pattern and function within the organism. COX-1 is constitutively expressed in many tissues and is thought to be responsible for maintaining gastric mucosal integrity. COX-2 is induced by a variety of stimuli and plays an important role in wound healing, ovulation, fertilization, and in mediating inflammation (for review, see Ref. 10). COX-2 expression levels are increased in colorectal cancer tissues (11–14). Overexpression of COX-2 has been shown to mediate cell cycle progression and to contribute to such diverse processes such as apoptosis, angiogenesis (15), and tissue invasion (16). On the basis of these effects, we have investigated the role of specific COX-2 inhibitors in the treatment of advanced colorectal cancer.

We have studied previously the effects of a selective COX-2 inhibitor, SC-58125, on the growth and viability of colorectal carcinoma cells grown *in vitro* and *in vivo* (7). The precise mechanism for growth inhibition of tumors is under evaluation, but does not seem to involve the induction of apoptosis *in vivo* (17). It has been reported that nonselective COX inhibitors induce a G₂-M cell cycle arrest resulting in a corresponding reduction of p34^{cdc2} levels and activity (18). We have observed a similar effect after SC-58125 treatment of colorectal carcinoma cells.⁴ These results suggest that SC-58125 may have therapeutic potential for treatment of colorectal cancer. However, SC-58125 will not be developed for clinical use in humans. Furthermore, although SC-58125 reduces tumor growth, it does not cause tumor regression (7, 9). Therefore, we evaluated other COX-2 inhibitors for their effect on the growth of colorectal carcinoma cells *in vitro* and *in vivo*.

Here we report that celecoxib (Celebrex), at concentrations >10 μM , potently induces apoptosis and inhibits cell cycle progression in colorectal carcinoma cells grown in culture by mechanisms independent of COX-2 inhibition. Lower concentrations of the drug have no discernible effect on cells grown *in vitro*. We also found that when celecoxib was administered to mice by inclusion in the diet (1250 mg/kg of chow), serum concentrations of $\sim 2.3 \mu\text{M}$ were achieved. Nevertheless, in contrast to the results from cell culture experiments, celecoxib significantly reduced the growth rate of colorectal carcinoma cells grown as xenografts, without toxicity to the normal intestine. These data highlight the point that the biological effects of celecoxib against cells grown in culture do not predict its effects *in*

⁴ Unpublished data.

vivo. However, these findings do support additional investigation for the use of celecoxib as a therapeutic agent against colorectal cancer.

MATERIALS AND METHODS

Reagents. Celecoxib (SC-58635), SC-58125, and SC-560 were provided as kind gifts from G. D. Searle and Co. (St. Louis, MO). Merck and Co. (Rathway, NJ) provided sulindac sulfide. DMSO (EM Science, Gibbstown, NJ) was used as the solvent. Concentrated drug stocks were diluted in DMEM (Life Technologies, Inc., Grand Island, NY) media before addition to cell cultures. The DMSO concentration in cultures was kept at 0.1%.

Cell Culture. HCA-7 cells were a generous gift from Susan Kirkland. LLC and HCT-15 cells and HCT-116 cells were purchased from the American Type Tissue Collection (Manassas, VA).

MEF Derivation. Primary MEFs were derived by passing day 13.5 C57BL/6J mouse embryos through an 18-gauge needle. The cells were expanded over several days in DMEM supplemented with 10% FBS, 1% P/S/1% L-glutamine. The *cox-2* genotype of the MEF cell lines was verified as described previously (19).

Cell Counts. LLC cells were seeded into six-well plates at 2.5×10^4 cells/well. Cells were treated in triplicate with DMSO or 25, 50, or 100 μM celecoxib for 12 h and then harvested and counted using a Coulter counter model Z1 (Coulter, Fullerton, CA).

In Vitro Viability Assay. The MTT assay was used to determine cell viability/proliferation. This assay measures mitochondrial activity. MTT is a yellow-colored tetrazolium salt that is taken up and cleaved only by metabolically active cells, reducing it to a colored, water-insoluble formazan salt. The solubilized formazan product can be quantified via absorbance at 570 nm measured using a 96-well-format spectrophotometer, and the absorbance correlates directly with cell number. Cells were plated at 1.5×10^4 cells/well in a 100- μl volume in 96-well plates and grown for 24 h in DMEM supplemented with 10% FBS. The indicated amount of test drug or DMSO in 1% FBS containing OptiMEM media was then added to the wells. At the indicated times, 10 μl of MTT (5 mg/ml) was added, and the cells were incubated at 37°C for 4 h. The tetrazolium crystals were solubilized by the addition of 10% SDS in 0.01 N HCl. After overnight incubation at 37°C, the absorbance was measured at 570 nm using a 96-well spectrophotometric plate reader (Packard Instruments, Meriden, CT). Results are expressed as the mean \pm SD of six wells.

Prostaglandin Measurement. Subconfluent cell cultures were treated with either celecoxib or SC-58125 for the indicated time. Thirty min before harvesting media, arachidonate was added to the media to a final concentration of 10 μM . PGE₂ was quantified as described previously (20).

Apoptosis Determination. The APO-Direct kit (PharMingen, San Diego, CA) was used for quantitative evaluation of apoptosis in response to celecoxib treatment. This assay relies on the characteristic fragmentation of DNA during the apoptotic process. Terminal nucleotidyl transferase enzyme is used to end-label the free 3'-OH of fragmented DNA using fluorescein-conjugated dUTP is used as the nucleotide for the exchange reaction. Flow cytometric detection of fluorescein-labeled fragmented DNA was conducted to quantitatively evaluate apoptosis in response to celecoxib treatment. Cells were seeded in 100-mm plates, and when 80% confluent cells were treated with DMSO, or 12.5, 25, or 50 μM celecoxib in 1% FBS-supplemented OptiMEM media. Negative control cells were not treated; vehicle-treated cells had an amount of DMSO equivalent to the celecoxib-treated cells added. Cells were harvested after 12 h of treatment, washed in PBS twice then fixed in 1% paraformaldehyde for 15 min and permeabilized by the addition of ice-cold 70% ethanol. The labeling reaction was performed according to the manufacturer's recommendations, with the exception that 2×10^6 cells/condition were used. After labeling, cells were washed three times in PBS, then resuspended in 1 ml of propidium iodide staining solution (5 $\mu\text{g/ml}$ propidium iodide, 40 $\mu\text{g/ml}$ RNASE A, in $1 \times$ PBS). Cells were then filtered through 50- μm mesh immediately before analysis on a Becton Dickinson FACScan flow cytometer. Gating on FL2-width was used to exclude aggregates, and 10^4 gated events were collected and analyzed.

Xenograft Model of Tumor Biology. HCA-7 cells were grown on plastic culture dishes according to standard cell culture techniques (7). The cells were trypsinized and resuspended in sterile PBS, then pelleted by brief centrifuga-

tion at 1500 rpm. The supernatant was aspirated, and cells were resuspended in PBS and counted using a hemocytometer. A final concentration of 5×10^7 cells/ml was made, and 100 μl of cell suspension was injected s.c. using a tuberculin syringe and a 27-gauge needle. Celecoxib was administered in the diet (1250 mg/kg of chow) and the animals were fed *ad libitum*. The size of the tumor was determined by direct measurement of tumor dimensions as described previously (7).

Scoring Apoptosis and Mitosis in Normal Intestinal Epithelium. In brief, mice were fed celecoxib (1250 mg/kg of chow) and the small and large intestines were removed separately and prepared by methods described in detail elsewhere (21, 22). Apoptotic bodies were identified by their characteristic morphological appearance in sections stained with H&E. To score the sections, each cell along the long axis of the crypt was numbered sequentially from the base of the crypt; the cell at the crypt base was designated as number 1. Each cell position was scored as containing either a normal, mitotic, or apoptotic cell. For each cell position, apoptotic and mitotic percentages were calculated. We have already shown this methodology to be more accurate when examining tissues than when identifying apoptotic bodies by TUNEL staining (23).

Gut Morphometry. Villus height and crypt depth was measured on an Axiohome microscope on 20 crypts and 20 villi from at least six animals/experimental group.

RESULTS

Treatment with Celecoxib Induces Concentration-dependent Apoptosis in Vitro. We have observed previously that a compound related to celecoxib, SC-58125, effectively inhibits the growth of LLC and colorectal cancer cells by 40% after 3 days of treatment *in vitro*.⁴ When celecoxib became available, we sought to test its effect on the growth of LLC cells. SC-58125-treated LLC cells remain morphologically unchanged; however, when LLC cells were treated with 50 μM celecoxib for 12 h, they became rounded and detached from the culture dish (Fig. 1A, left panel). As with SC-58125, we found that celecoxib inhibited the growth of LLC cells but was much more potent. After only 12 h of treatment, the IC₅₀ for celecoxib inhibition of cell number was 40 μM (Fig. 1A, right panel). The cytotoxicity of celecoxib against LLC cells was also quantified using the MTT assay. When celecoxib concentrations of $<10 \mu\text{M}$ were used, no effect on cell viability or growth was observed.⁵ However, treatment for 12 h with $>20 \mu\text{M}$ celecoxib resulted in a dramatic decrease in cell viability (Fig. 1B). We next determined whether celecoxib also had potent cytotoxicity against two human colorectal carcinoma cell lines (HCT-15 and HCA-7). Celecoxib had similar cytotoxic effects against these cells with significant loss of viability at concentrations of $>20 \mu\text{M}$ (Fig. 1B). Once again, treatment of these cells with celecoxib for 3 days at concentrations of $<10 \mu\text{M}$ had no effect on cell viability.

To determine whether the cytotoxic effects of celecoxib were the result of induction of apoptosis, LLC, HCA-7, and HCT-15 cells were treated with 25 or 50 μM celecoxib. Apoptotic cells were identified by TUNEL staining and the results analyzed by dual parameter flow cytometry using DNA content to identify the cell cycle position of cells undergoing apoptosis. Treatment with 50 μM celecoxib caused significant apoptosis in all three cell lines with 88%, 37%, and 15% of LLC, HCA-7, and HCT-15 cells, respectively, undergoing apoptosis (Fig. 2). We found that LLC and HCA-7 cells undergoing apoptosis contained 4 N DNA content, suggesting that arrest at the G₂-M stage of the cell cycle was occurring. In contrast, apoptotic HCT-15 cells contained a subdiploid DNA content after celecoxib treatment, suggesting that apoptosis was occurring when the cells contained 2 N DNA. The induction of apoptosis was confirmed by the detection of oligonucleosomal cleavage of DNA in LLC and HCA-7 cells after treatment with 50 μM celecoxib for 6 h.⁵

⁵ Data not shown.




Fig. 1. Celecoxib inhibits the growth of carcinoma cells. *A*, left panel, phase-contrast photomicrographs of cells treated for 12 h with 50 μM of celecoxib (SC-58635) or DMSO (control); right panel, LLC cells treated with the indicated amount of celecoxib for 12 h. Attached cells were suspended and counted using a Coulter counter. Data represents averages and SDs from triplicate wells. *A*, LLC (diamond), HCA-7 (square), and HCT-15 (triangle) cell viability after treatment with vehicle (DMSO) or with the indicated amount of celecoxib for 12 h. Cell viability was determined using the MT assay. Value represents OD_{570} -treated/ OD_{570} control (DMSO), and are expressed as averages \pm SD of six replicate wells/treatment condition.

Induction of Apoptosis *in Vitro* Is Independent of COX-2 Inhibition. Next we sought to test whether cyclooxygenase inhibition was required for the cytotoxic effect of celecoxib treatment *in vitro*. If COX-2 is involved in inhibition of cell growth under these experimental conditions, then one would expect that other COX-2 selective inhibitors would have similar effects. We directly tested this hypothesis by treating LLC cells for 12 h with the COX-2 selective inhibitors, celecoxib and SC-58125, the COX-1 selective inhibitor, SC-560, and the nonselective COX inhibitor, sulindac sulfide. SC-58125 did not affect cell viability over the time course of the experiment and sulindac sulfide and SC-560 decreased MTT activity by 15–25%, whereas celecoxib resulted in virtually complete loss of cell viability (Fig. 3A). The lack of correlation between COX-2 inhibitory activity and cytotoxicity suggests that the effect of celecoxib *in vitro* is independent of its ability to inhibit COX-2. One possible explanation for the different effects of celecoxib and SC-58125 on carcinoma cells might be attributable to variation in inhibition profiles or half-lives of these compounds. We next compared inhibition of PGE_2 production by celecoxib and SC-58125. Celecoxib inhibits PGE_2 production in LLC cells with an IC_{50} of 18 nM. Both compounds had similar cyclooxygenase inhibition profiles, decreasing PGE_2 levels by 85–95% for at least 12 h after treatment (Fig. 3B.2, right panel).

Finally, although LLC cells express both COX-1 and COX-2, HCA-7 cells only express COX-2, and HCT-15 cells express neither isoform. Comparison of the cytotoxic effects of celecoxib between these cell lines is not strictly valid because of differences at many

other genetic loci. Therefore, we tested the effect of celecoxib on MEFs derived from *cox-2* (+/+), (+/-), and (-/-) C57BL/6J mice. The cytotoxicity profile and IC_{50} of celecoxib was similar in all three cell lines regardless of COX-2 status (Fig. 3C). Furthermore, the concentration dependence of cytotoxicity for the cell lines of all three MEFs and LLC cells were similar (Fig. 3C). As with the colon carcinoma cells tested (Fig. 1B), no cell death was observed at celecoxib concentrations of $<10 \mu\text{M}$. These results strongly indicate that the cytotoxicity of celecoxib *in vitro* is independent of inhibition of COX-2 and brings up the question of whether this drug is cytotoxic *in vivo*.

Celecoxib Does Not Cause Apoptosis or Inhibit Cell Division in Normal Gut Epithelium. The COX-2 independent toxicity of celecoxib to MEFs raises the possibility that celecoxib might cause significant apoptosis and/or cell cycle arrest in normal gut epithelium. Therefore, we determined apoptotic and mitotic rates in the small and large intestinal epithelium of nude mice fed celecoxib (1250 mg/kg chow) for 45 days. As described previously, spontaneous rates of apoptosis in the mouse intestine were found to be low (23). Treatment with celecoxib for 45 days did not cause any significant change in apoptotic or mitotic rates in either the small or large intestine (Fig. 4A). Evaluation of intestinal epithelium 4 h after administration of a single dose of celecoxib did not reveal increased rates of apoptosis.⁵ Furthermore, no significant differences in gut morphology could be discerned. In the small intestine, villus height was 303 ± 95 versus $289 \pm 86 \mu\text{m}$ in control and celecoxib-treated animals, and crypt




Fig. 2. Celecoxib induces programmed cell death. Dual parameter flow cytometric analysis measuring both FITC-labeled fragmented DNA (*ordinate*) and cellular DNA content (*abscissa*) in LLC (6 h), HCA-7 (12 h), or HCT-15 (12 h) cells after treatment with celecoxib. *Negative*, TUNEL enzyme was not added to labeling reaction, vehicle (DMSO), or 25 or 50 μM of celecoxib.

depth was $100.4 \pm 17 \mu\text{m}$ in control animals and $100 \pm 38 \mu\text{m}$ treated mice. Similarly, the colon crypt length was unaffected by treatment with celecoxib $150 \pm 25 \mu\text{m}$ in control colon *versus* $167 \pm 31 \mu\text{m}$ in treated colon.

To relate the *in vivo* observations to the cell culture studies above, the serum concentration of celecoxib in six mice was measured and found to range from 1.5–3.5 μM ($2.3 \pm 0.7 \mu\text{M}$, Fig. 4B). Thus the peak serum concentration of celecoxib achieved in mice using this dosing regime was 10-fold less than that which led to cytotoxicity *in vitro*.

In Vivo Tumor Growth Is Attenuated by Celecoxib Treatment.

We have reported previously that celecoxib attenuates the growth of LLC tumors *in vivo* (17). Given the recent interest in the use of COX-2 inhibitors for chemoprevention of colorectal cancer, we wished to confirm and extend these observations in a colorectal carcinoma cell line. HCA-7 cells were implanted *s.c.* into athymic mice. Celecoxib was mixed with mouse chow at a concentration of 1250 mg/kg. (The xenograft and normal intestinal studies reported above were performed in the same mice). Whereas data from our *in vitro* studies may predict that such a low dose of celecoxib (2.3 μM) would be ineffective at inhibiting tumor growth, we found that the

growth of HCA-7 xenograft growth was significantly reduced when compared with animals on the control diet (Fig. 5). Delaying treatment for 10, 20, or 30 days postimplantation, by which point the tumors were well established, still resulted in inhibition of tumor growth, indicating that the compound was not affecting tumor implantation (Fig. 5). These data demonstrate that celecoxib prevents the growth of colorectal carcinoma cells *in vivo* at concentrations that do not cause apoptosis in cells grown *in vitro*.

DISCUSSION

We have shown previously that the specific COX-2 inhibitor, SC-58125, can reduce the growth of COX-2 positive colorectal cancer xenografts (7). We have also observed that SC-58125 primarily affected cell cycle progression via a delay in $G_2\text{-M}$.⁴ Preliminary studies on celecoxib, another specific COX-2 inhibitor, suggested that it might be even more potent than SC-58125. We studied its effects on the growth of LLC cells and two colon carcinoma cell lines grown *in vitro* and compared those results with its effect on tumor growth *in vivo*, as well as evaluating its effect on the normal intestine of treated mice.




Fig. 3. Celecoxib inhibits cell growth independent of COX-2 inhibition. *A*, MTT assay of LLC cells treated for 12 h with 50 μM of SC-58125, NS-398, sulindac sulfide, SC-560, or celecoxib. *B*, prostaglandin-inhibition profile. *B.1*, LLC PGE₂ levels were determined after 3 h of celecoxib treatment at the indicated dose. *B.2*, LLC PGE₂ levels after 12 h of treatment with either celecoxib (black squares) or SC-58125 (black circles) at 1, 5, or 10 μM . Data are represented as the fold-reduction in PGE₂ compared with vehicle-treated cells. *C*, MEFs derived from *cox-2* (+/+), *cox-2* (+/-), and *cox-2* (-/-) C57/BL6 mice were treated with the indicated amount of celecoxib for 12 h. Values represent a fraction of control (DMSO)-treated cells. Viability was determined by the MTT assay and cellular morphology. Each data point represents the mean of six replicates, error bars represent \pm SD.

At concentrations >20 μM , celecoxib induces cell death as measured by cell counts of LLC cells and the assay of LLC, HCA-7, and HCT-15 cells (Fig. 1). Analysis of TUNEL staining by flow cytometry (Fig. 2) and the demonstration of oligonucleosomal DNA cleavage confirms that the loss of viability is attributable to the induction of apoptosis. In addition, a G₂-M block is also induced in LLC and HCA-7 cells by treatment with 50 μM celecoxib. Interestingly, in HCT-15 cells we observed a subdiploid accumulation of fragmented DNA suggesting that apoptosis was occurring during the G₁ to early S transition of the cell cycle. These results show that celecoxib, at concentrations >20 μM , strongly induces apoptosis in cells grown *in vitro*.


Three lines of evidence indicated that induction of apoptosis *in vitro* by high-dose celecoxib occurs by mechanisms independent of COX-2 inhibition: (a) the cytotoxicity of a variety of NSAIDs did not correlate with their ability to inhibit COX-2 (Fig. 3A); (b) although SC-58125 and celecoxib can both inhibit PGE₂ production by $>90\%$ at a concentration of 10 μM , SC-58125 at 50 μM has no effect on cell viability, whereas celecoxib reduces viability by 70% when given at that concentration; and (c) MEFs derived from wild-type, *cox-2* heterozygous or *cox-2* null C57/BL6 mice have similar sensitivities to the cytotoxic effects of celecoxib. These results provide compelling evidence that the cytotoxicity of these agents *in vitro* is independent of

COX-2 inhibition, and indicates that high-dose celecoxib is cytotoxic not only to carcinoma cells *in vitro*, but also to nontransformed mammalian cells grown in culture.

A crucial observation from these studies is that treatment with 10 μM or less of celecoxib for 3 days has no detectable effect on cell death *in vitro*. Despite this lack of effect in cultured cells, we found that celecoxib strongly attenuated the growth of xenografted HCA-7 tumors *in vivo*, although the plasma concentration of celecoxib was ~ 2.3 μM . This discrepancy highlights the fact that tumor growth *in vivo* is determined by the interaction between factors intrinsic to tumor cells, the extracellular matrix, stromal cells, and other host factors. These factors are not always present *in vitro* when cells are grown on plastic culture dishes. Cell culture models are often used to evaluate the therapeutic potential of NSAIDs against cancer, but great caution needs to be taken when extrapolating *in-vitro* results to the whole organism, particularly with respect to the relative dose of agent used.

These studies do not address the issue of the mechanism by which celecoxib attenuates tumor growth *in vivo* or evaluate its COX-2 dependency. *In vivo*, interference with the implantation of carcinoma cells does not seem to play a significant role because growth is still affected even when treatment is started 30 days after inoculation of the host. It has not been well established that COX-2 expression in the

A.



B.




Fig. 4. Celecoxib does not induce apoptosis or mitosis in gut epithelium. *A*, nude mice were treated for 45 days with celecoxib. Both small intestine and colon were harvested, fixed, and H&E-stained. Apoptotic bodies and mitotic figures were counted per high-power field, and indices standardized to saline-treated control epithelium were generated. *B*, serum celecoxib levels. Six animals were treated with 1250 mg/kg celecoxib for 45 days. Serum was harvested and celecoxib levels were measured by mass spectroscopy. Data are presented as the mean (bar) and spread.

carcinoma cells *in vivo* contributes to tumor growth. Previous studies using tumor models lacked the ability to distinguish between tumor-derived COX-2 and extrinsic COX-2. We recently have reported that stromally derived COX-2 is important for tumor growth (17). It is possible that whereas low doses of celecoxib do not affect tumor cells *in vitro*, stromal or host-derived processes dependent on COX-2 may be sensitive to low-dose NSAID treatment. Angiogenesis, a necessary component of neoplastic growth, is also promoted by COX-2 activity, and its inhibition could play a crucial role in the antineoplastic action of celecoxib (15, 17, 24, 25).

In striking contrast to the antitumor effects of celecoxib, no toxic effects were observed in the normal gut by assessing induction of

apoptosis, inhibition of cell division, or reduction of crypt and villus dimensions. The explanation for the differential effect of celecoxib on normal and transformed intestinal epithelial tissue is not known, but it is interesting to note that COX-2 expression and activity is very low to undetectable in normal gut mucosa. Furthermore, gut epithelium is not as dependent on new-vessel growth as tumor xenografts.

In summary, celecoxib can significantly attenuate the growth of colorectal carcinoma xenografts without adverse effects on the normal gut. Although celecoxib causes apoptosis of colorectal cancer cells grown *in vitro* via a COX-2-independent mechanism, this only occurs at concentrations >10-fold higher than what can be achieved *in vivo*. Great caution should be taken in interpreting the clinical significance




Fig. 5. Celecoxib inhibits the *in-vivo* growth of HCA-7 cells. 5×10^6 HCA-7 cells were injected s.c. into the dorsal flank of athymic mice. Mice received either control chow or chow containing 1250 mg/kg of celecoxib. Treatment was initiated either at the time of tumor implantation (red square), or 10 (yellow triangle), 20 (green square), or 30 (purple square) days postimplantation. Tumor dimensions were measured at the indicated intervals, and tumor volume was calculated according to the equation $V = [L \times W^2] \times 0.5$, where V = volume, L = length, and W = width (26).

of cell culture data, particularly with respect to drug levels that could be safely achieved in humans. Overall, celecoxib seems to have significant therapeutic potential against colorectal cancer, and additional clinical evaluation is warranted.

ACKNOWLEDGMENTS

We thank Penny Ottewell for her technical assistance.

REFERENCES

- Thun, M. J., Nambodiri, M. M., and Heath, C. W. J. Aspirin use and reduced risk of fatal colon cancer. *N. Engl. J. Med.*, *325*: 1593–1596, 1991.
- Giovannucci, E., Rimm, E. B., Stampfer, M. J., Colditz, G. A., Ascherio, A., and Willett, W. C. Aspirin use and the risk for colorectal cancer and adenoma in male health professionals. *Ann. Intern. Med.*, *121*: 241–246, 1994.
- Giardiello, F. M., Hamilton, S. R., Krush, A. J., Piantadosi, S., Hyland, L. M., Celano, P., Booker, S. V., Robinson, C. R., and Offerhaus, G. J. Treatment of colonic and rectal adenomas with sulindac in familial adenomatous polyposis. *N. Engl. J. Med.*, *328*: 1313–1316, 1993.
- Mahmoud, N. N., Boolbol, S. K., Dannenberg, A. J., Mestre, J. R., Bilinski, R. T., Martucci, C., Newmark, H. L., Chadburn, A., and Bertagnolli, M. M. The sulfide metabolite of sulindac prevents tumors and restores enterocyte apoptosis in a murine model of familial adenomatous polyposis. *Carcinogenesis (Lond.)*, *19*: 87–91, 1998.
- Beazer-Barclay, Y., Levy, D. B., Moser, A. R., Dove, W. F., Hamilton, S. R., Vogelstein, B., and Kinzler, K. W. Sulindac suppresses tumorigenesis in the min mouse. *Carcinogenesis (Lond.)*, *17*: 1757–1760, 1996.
- Samaha, H. S., Kelloff, G. J., Steele, V., Rao, C. V., and Reddy, B. S. Modulation of apoptosis by sulindac, curcumin, phenylethyl-3-methylcaffeate, and 6-phenylhexyl isothiocyanate: apoptotic index as a biomarker in colon cancer chemoprevention and promotion. *Cancer Res.*, *57*: 1301–1305, 1997.
- Sheng, H., Shao, J., Kirkland, S. C., Isakson, P., Coffey, R., Morrow, J., Beauchamp, R. D., and DuBois, R. N. Inhibition of human colon cancer cell growth by selective inhibition of cyclooxygenase-2. *J. Clin. Investig.*, *99*: 2254–2259, 1997.
- Goldman, A. P., Williams, C. S., Sheng, H., Lamps, L. W., Williams, V. P., Pairet, M., Morrow, J. D., and DuBois, R. N. Meloxicam inhibits the growth of colorectal cancer cells. *Carcinogenesis (Lond.)*, *19*: 2195–2199, 1998.
- Chinery, R., Beauchamp, R. D., Shyr, Y., Kirkland, S. C., Coffey, R. J., and Morrow, J. D. Antioxidants reduce cyclooxygenase-2 expression, prostaglandin production, and proliferation in colorectal cancer cells. *Cancer Res.*, *58*: 2323–2327, 1998.
- Williams, C., Shattuck-Brandt, R. L., and DuBois, R. N. The role of COX-2 in intestinal cancer. *Ann. NY Acad. Sci.*, *889*: 72–83, 1999.
- Eberhart, C. E., Coffey, R. J., Radhika, A., Giardiello, F. M., Ferrenbach, S., and DuBois, R. N. Up-regulation of cyclooxygenase 2 gene expression in human colorectal adenomas and adenocarcinomas. *Gastroenterology*, *107*: 1183–1188, 1994.
- Kutcher, W., Jones, D. A., Matsunami, N., Groden, J., McIntyre, T. M., Zimmerman, G. A., White, R. L., and Prescott, S. M. Prostaglandin H synthase-2 is expressed

abnormally in human colon cancer: evidence for a transcriptional effect. *Proc. Natl. Acad. Sci. USA*, *93*: 4816–4820, 1996.

- Sano, H., Kawahito, Y., Wilder, R. L., Hashiramoto, A., Mukai, S., Asai, K., Kimura, S., Kato, H., Kondo, M., and Hla, T. Expression of cyclooxygenase-1 and -2 in human colorectal cancer. *Cancer Res.*, *55*: 3785–3789, 1995.
- Kargman, S., O'Neill, G., Vickers, P., Evans, J., Mancini, J., and Jothy, S. Expression of prostaglandin G/H synthase-1 and -2 protein in human colon cancer. *Cancer Res.*, *55*: 2556–2559, 1995.
- Tsuji, M., Kawano, S., Tsuji, S., Sawaoka, H., Hori, M., and DuBois, R. N. Cyclooxygenase regulates angiogenesis induced by colon cancer cells. *Cell*, *93*: 705–716, 1998.
- Tsuji, M., Kawano, S., and DuBois, R. N. Cyclooxygenase-2 expression in human colon cancer cells increases metastatic potential. *Proc. Natl. Acad. Sci. USA*, *94*: 3336–3340, 1997.
- Williams, C. S., Tsuji, M., Reese, J., Dey, S. K., and DuBois, R. N. Host cyclooxygenase-2 modulates carcinoma growth. *J. Clin. Investig.*, *105*: 1589–1594, 2000.
- Goldberg, Y., Nassif, I. I., Pittas, A., Tsai, L. L., Dynlacht, B. D., Rigas, B., and Shiff, S. J. The anti-proliferative effect of sulindac and sulindac sulfide on HT-29 colon cancer cells: alterations in tumor suppressor and cell cycle-regulatory proteins. *Oncogene*, *12*: 893–901, 1996.
- Dinchuk, J. E., Car, B. D., Focht, R. J., Johnston, J. J., Jaffee, B. D., Covington, M. B., Contel, N. R., Eng, V. M., Collins, R. J., Czerniak, P. M., Gorry, S. A., and Trzaskos, J. M. Renal abnormalities and an altered inflammatory response in mice lacking cyclooxygenase II. *Nature (Lond.)*, *378*: 406–409, 1995.
- DuBois, R. N., Tsuji, M., Bishop, P., Awad, J. A., Makita, K., and Lanahan, A. Cloning and characterization of a growth factor-inducible cyclooxygenase gene from rat intestinal epithelial cells. *Am. J. Physiol.*, *266*: G822–G827, 1994.
- Merritt, A. J., Allen, T. D., Potten, C. S., and Hickman, J. A. Apoptosis in small intestinal epithelial from p53-null mice: evidence for a delayed, p53-independent G2/M-associated cell death after γ -irradiation. *Oncogene*, *14*: 2759–2766, 1997.
- Pritchard, D. M., Watson, A. J., Potten, C. S., Jackman, A. L., and Hickman, J. A. Inhibition by uridine but not thymidine of p53-dependent intestinal apoptosis initiated by 5-fluorouracil: evidence for the involvement of RNA perturbation. *Proc. Natl. Acad. Sci. USA*, *94*: 1795–1799, 1997.
- Bach, S. P., Chinery, R., O'Dwyer, S. T., Potten, C. S., Coffey, R. J., and Watson, A. J. Pyrrolidinedithiocarbamate increases the therapeutic index of 5-fluorouracil in a mouse model. *Gastroenterology*, *118*: 81–89, 2000.
- Uefuji, K., Ichikura, T., and Mochizuki, H. Cyclooxygenase-2 expression is related to prostaglandin biosynthesis and angiogenesis in human gastric cancer. *Clin. Cancer Res.*, *6*: 135–138, 2000.
- Masferrer, J. L., Leahy, K. M., Koki, A. T., Zweifel, B. S., Settle, S. L., Woerner, B. M., Edwards, D. A., Flickinger, A. G., Moore, R. J., and Seibert, K. Antiangiogenic and antitumor activities of cyclooxygenase-2 inhibitors. *Cancer Res.*, *60*: 1306–1311, 2000.
- Wang, J., Sun, L., Myeroff, L., Wang, X., Gentry, L. E., Yang, J., Liang, J., Zborowska, E., Markowitz, S., Willson, J. K., and Brattain, M. G. Demonstration that mutation of the type II transforming growth factor β receptor inactivates its tumor suppressor activity in replication error-positive colon carcinoma cells. *J. Biol. Chem.*, *270*: 22044–22049, 1995.

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

Celecoxib Prevents Tumor Growth *in Vivo* without Toxicity to Normal Gut: Lack of Correlation between *in Vitro* and *in Vivo* Models

Christopher S. Williams, Alastair J. M. Watson, Hongmiao Sheng, et al.

Cancer Res 2000;60:6045-6051.

Updated version Access the most recent version of this article at:
<http://cancerres.aacrjournals.org/content/60/21/6045>

Cited articles This article cites 26 articles, 10 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/60/21/6045.full#ref-list-1>

Citing articles This article has been cited by 57 HighWire-hosted articles. Access the articles at:
<http://cancerres.aacrjournals.org/content/60/21/6045.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerres.aacrjournals.org/content/60/21/6045>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.