

## NY-ESO-1 Expression and Immunogenicity Associated with Transitional Cell Carcinoma: Correlation with Tumor Grade<sup>1</sup>

Takushi Kurashige, Yuji Noguchi,<sup>2</sup> Takashi Saika, Toshiro Ono, Yasuhiro Nagata, Achim Jungbluth, Gerd Ritter, Yao-Tseng Chen, Elisabeth Stockert, Tomoyasu Tsushima, Hiromi Kumon, Lloyd J. Old, and Eiichi Nakayama

Departments of Urology [T. K., T. S., T. T., H. K.] and Immunology [Y. No., T. O., E. N.], Okayama University Medical School, Okayama 700-8558, Japan; Ludwig Institute for Cancer Research, New York Branch at Memorial Sloan-Kettering Cancer Center, New York, New York 10021 [Y. Na., A. J., G. R., Y-T. C., E. S., L. J. O.]; and Department of Pathology, Cornell University Medical College, New York, New York 10021 [Y-T. C.]

### Abstract

*NY-ESO-1* mRNA expression in transitional cell carcinoma was investigated by reverse transcription-PCR and immunohistochemistry. *NY-ESO-1* mRNA was detected in 20 of 62 (32%) tumor specimens. There was a correlation between *NY-ESO-1* expression and tumor grade: 0 of 4 (0%) grade 1 (G1), 6 of 26 (23%) grade 2 (G2), and 14 of 32 (44%) grade 3 (G3) tumors were *NY-ESO-1* mRNA positive. Immunohistochemical analysis using *NY-ESO-1*-specific monoclonal antibody ES121 showed that 2 of 14 *NY-ESO-1* mRNA-expressing G3 tumors were positive for *NY-ESO-1*. No *NY-ESO-1* staining was observed in the panel of 30 G1 or G2 tumor specimens, including 6 *NY-ESO-1* mRNA-positive cases. Sera from an expanded panel of 124 patients with transitional cell carcinoma were tested for the presence of *NY-ESO-1* antibody. Seropositivity was observed in 9 of 72 (12.5%) patients with G3 tumors, whereas none of 52 patients with G1 or G2 tumors produced antibody against *NY-ESO-1*. In the 9 positive patients with *NY-ESO-1* antibody, 4 had muscular invasive tumors, and 5 had carcinoma *in situ*.

### Introduction

The list of human tumor antigens recognized by human CD8 T cells or antibodies has grown rapidly over the past several years (1). Most of these antigens can be classified according to expression pattern or structural features into one of the following categories: (a) CT<sup>3</sup> antigens, e.g., members of the *MAGE* gene family (2) and *NY-ESO-1* (3); (b) differentiation antigens, e.g., tyrosinase (4), gp 100 (5), and Melan A/MART-1 (6, 7); (c) mutated gene products, e.g.,  $\beta$ -catenin (8) and caspase-8 (9); and (d) overexpressed self-antigens, e.g., *HER2/neu* (10). Because of their broad representation in cancer and restricted expression in normal tissues, CT antigens have received much attention as potential targets for human cancer vaccine (11). Of the ten gene or gene families coding for CT antigens (12), the *NY-ESO-1* product appears to be the most immunogenic, inducing a humoral immune response and specific CD8 T-cell reactivity in ~50% of patients with advanced *NY-ESO-1*-expressing tumors (13, 14). Immunity to *NY-ESO-1* is clearly antigen driven, disappearing with tumor removal or tumor regression (15).

TCC comprises nearly 90% of primary malignant tumors of the bladder, displaying a broad biological spectrum ranging from superficial to invasive tumors (16). Tumor grade (G) is based on cellular dysplasia and architectural abnormalities in tumor tissue and is com-

monly used to classify TCC in terms of malignant potential (17). High-grade (G3) TCC progresses to muscle invasion more frequently than low-grade (G1 and G2) tumors. Although G1 and G2 tumors often recur, they are less likely to become invasive (18). Although CIS is a flat superficial tumor, it is classified as a high-grade lesion, which frequently progresses into invasive tumors (17, 18).

In the initial description for *NY-ESO-1*, we found a high frequency of *NY-ESO-1* mRNA expression in a small number of TCCs (3). In this study, we have evaluated *NY-ESO-1* expression in a large number of TCCs and have found a correlation between *NY-ESO-1* expression and tumor grade. In addition, the sera of patients with TCCs have been screened for antibody against *NY-ESO-1*, and the antibody response has also been found to be restricted to patients with G3 tumors.

### Materials and Methods

**Patients.** TCC specimens were obtained from 62 patients undergoing surgery. The specimens consisted of 54 bladder, 4 ureteral, and 4 renal pelvic TCCs from 49 males and 13 females with a mean age at diagnosis of 65.7 years (range, 25–86 years). Tumor grade was determined according to standard criteria (17), and TNM classification of TCC was assessed according to the consensus report (19). Sera from 124 patients (94 males and 30 females) with a mean age at diagnosis of 66.4 years (range, 25–87 years) were tested for serum antibody to *NY-ESO-1*. All patients, with the exception of patient TCC44, had tumors at the time of serum collection.

**RT-PCR Analysis.** mRNA was isolated from frozen tumor specimens using the QuickPrep Micro mRNA Purification kit (Pharmacia, Uppsala, Sweden). Isolated mRNA was subjected to cDNA synthesis using the First-Strand cDNA Synthesis kit (Pharmacia). Primers for RT-PCR were: ESO1-1, 5'-AGTTCACCTCGCCATGCCT-3'; and ESO1-2, 5'-TCCTCCTCCAGC-GACAAACAA-3'. The amplification program for *NY-ESO-1* was 1 min at 94°C, 1 min at 60°C, and 1.5 min at 72°C for 35 cycles after denaturing at 94°C for 1 min. These cycles were followed by a 10-min elongation step at 72°C. The PCR products (385 bp) were analyzed on 0.8% agarose gel.

**ELISA.** Recombinant *NY-ESO-1* protein solution (100  $\mu$ l/well) at a concentration of 1  $\mu$ g/ml in coating buffer [15 mM Na<sub>2</sub>CO<sub>3</sub>, 35 mM NaHCO<sub>3</sub> in distilled water (pH 9.6)] was added to 96-well plates (Nunc, Roskilde, Denmark) and incubated overnight at 4°C. Plates were washed with 0.05% Tween 20/PBS and blocked with 100  $\mu$ l/well of 5% FCS/PBS for 1 h at room temperature. After washing, patients' sera (100  $\mu$ l/well) serially diluted with 5% FCS/PBS were added to the plate and incubated for 2 h at room temperature. After washing, diluted goat antihuman IgG (100  $\mu$ l/well) labeled with peroxidase (MBL, Nagoya, Japan) was added and incubated for 1 h at room temperature. After washing, substrate solution [50 mM citric acid, 100 mM Na<sub>2</sub>HPO<sub>4</sub>, 0.03% *ortho*-phenylenediamine, 0.1% H<sub>2</sub>O<sub>2</sub> in distilled water (pH 5.0)] was added in each well and incubated for 15 min at room temperature. After adding 3 M H<sub>2</sub>SO<sub>4</sub>, plates were read by U-2001 spectrophotometer (TOSOH, Tokyo, Japan).

**Immunohistostaining.** Tumor specimens were fixed with buffered formalin and embedded in paraffin. Sections (5  $\mu$ m) were placed on glass slides, heated at 60°C overnight, and then deparaffinized with xylene and ethanol. For antigen retrieval, tumor specimens mounted on glass slides were immersed into preheated target retrieval solution (DAKO, Carpinteria, CA) for 15 min and

Received 2/12/01; accepted 5/1/01.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

<sup>1</sup> Supported in part by a Grant-in-Aid for Scientific Research on Priority Area (C) from the Ministry of Education, Science, Sports and Culture of Japan.

<sup>2</sup> To whom requests for reprints should be addressed, at Department of Immunology, Okayama University Medical School, 2-5-1 Shikata-cho, Okayama 700-8558, Japan. Phone: 81-86-235-7188; Fax: 81-86-235-7193; E-mail: noguchi@med.okayama-u.ac.jp.

<sup>3</sup> The abbreviations used are: CT, cancer/testis; TCC, transitional cell carcinoma; RT-PCR, reverse transcription PCR; CIS, carcinoma *in situ*; BCG, *Bacillus Calmette-Guérin*; TNM, tumor-node-metastasis; mAb, monoclonal antibody.

allowed to cool for 20 min at room temperature. After the inactivation of endogenous peroxidase, specimens were incubated with 1.5% horse serum in PBS for 30 min at room temperature. mAb specific for NY-ESO-1 (clone ES121) was then added at a concentration of 2.5 μg/ml and incubated overnight at room temperature. After washing, diluted biotinylated antimouse IgG (Vector Laboratories, Burlingame, CA) was applied and incubated for 30 min at room temperature. Avidin labeled with peroxidase (Vector Laboratories) was added after washing and incubated for 30 min at room temperature. Diaminobenzidine tetrahydrochloride was then added for development, followed by counterstaining with hematoxylin solution.

**Results**

**Expression of NY-ESO-1 mRNA in TCC.** Expression of NY-ESO-1 mRNA in TCC specimens was investigated by RT-PCR. As shown in Table 1, NY-ESO-1 mRNA was detected in 20 of 62 (32%) tumor specimens. The size of PCR product in tumor was the same as in the T24 human bladder cancer cell line and in testis (Fig. 1A). The relationship between NY-ESO-1 mRNA expression and pathological and clinical features is shown in Table 1. NY-ESO-1 mRNA expression was correlated with tumor grade. Higher frequency of NY-ESO-1 mRNA expression was seen in higher grade TCC: 0 of 4 (0%) in G1; 6 of 26 (23%) in G2; and 14 of 32 (44%) in G3. No correlation could be established between NY-ESO-1 expression and age and sex of patients or with presence or absence of muscle invasion. TCC associated with lymph node or systemic metastases appeared to have a higher frequency of NY-ESO-1 expression than nonmetastatic tumors, but the numbers are too small to make a definitive statement.

**Immunohistostaining of NY-ESO-1.** The panel of 62 TCC specimens was analyzed for NY-ESO-1 protein expression by immunohistochemistry using NY-ESO-1-specific mAb ES121. Positive staining was observed in 2 of 14 NY-ESO-1 mRNA-positive G3 tumors. The pattern of staining was heterogeneous rather than homogeneous (Fig. 1B). At higher magnification, staining could be seen to be predominantly cytoplasmic. One of 2 NY-ESO-1 protein positive specimens was from a primary tumor of a patient with systemic metastases; the other one was from a patient with localized and superficial tumors but without metastasis. No mAb ES121 staining was observed in 30 G1 and G2 specimens, including 6 NY-ESO-1 mRNA-positive tumors. The lower frequency of NY-ESO-1 detection by immunohistochemical staining as compared with RT-PCR typing is consistent with our previous experience with ES121 staining of other tumor types.

**Antibody Response to NY-ESO-1 in TCC Patients.** Sera from 124 TCC patients were analyzed for NY-ESO-1 antibody by ELISA




Fig. 1. A, RT-PCR analysis for NY-ESO-1 mRNA expression. mRNA from G1 TCC (Lane 1), G2 TCC (Lanes 2–5), G3 TCC (Lanes 6–8), a human bladder cancer cell line, T24 (Lane 9), and testis (Lane 10) was analyzed for NY-ESO-1 mRNA by RT-PCR. B, H&E staining (left) and immunohistostaining for NY-ESO-1 using mAb ES121 (right) in serial sections of G1, G2, or G3 TCC.

Table 1 Correlation between NY-ESO-1 mRNA expression and pathological and clinical features in transitional cell carcinoma

Pathological and clinical features	NY-ESO-1 positive/tumors examined
All tumors	20/62 (32%)
Tumor grade <sup>a</sup>	
G1	0/4 (0%)
G2	6/26 (23%)
G3	14/32 (44%)
Tumor invasion	
Superficial	10/36 <sup>b</sup> (28%)
Muscle invasion	10/26 (38%)
Regional lymph node metastasis	
Negative	16/54 (30%)
Positive	4/8 (50%)
Systemic metastasis	
Negative	16/57 (28%)
Positive	4/5 (80%)

<sup>a</sup> The difference in frequency of NY-ESO-1 mRNA expression between high-grade (G3; 14 of 32) and low-grade (G1 and G2; 6 of 30) tumors was statistically significant (P < 0.05). Statistical analysis was performed by χ<sup>2</sup> test.

<sup>b</sup> In this series of 36 superficial tumors, 3 were CIS, and 2 of the 3 CIS were NY-ESO-1 positive.

using recombinant NY-ESO-1 protein. Fig. 2 shows titration curves of NY-ESO-1 antibody-positive and -negative sera, and Table 2 summarizes the results. NY-ESO-1 antibody was found in sera from 9 of 72 (12.5%) patients with G3 tumors. No antibody was detected in sera from 52 patients with G1 or G2 tumors or from 23 healthy volunteers. Table 3 lists TNM classification of TCC in patients with NY-ESO-1-positive antibody. Five of 9 seropositive patients had low-stage diseases, including the patients with CIS.

**Discussion**

This study documents that TCC is a tumor type with a high frequency of NY-ESO-1 mRNA expression. NY-ESO-1 expression is correlated with the tumor grade, with 44% of G3 TCCs, in contrast to 20% of G1 and G2 TCCs, expressing NY-ESO-1. Although this frequency of NY-ESO-1 expression in TCCs is higher than reported in several other tumor types, there needs to be a more extensive study of

these other tumor types relating *NY-ESO-1* expression to pathological and prognostic features. In addition, a survey of the *NY-ESO-1* phenotype of primary *versus* metastatic disease and different metastatic deposits in the same patients needs to be determined for TCC as well as other types of tumors. Although there is a tendency for *NY-ESO-1*-positive primary TCCs to be associated with a higher frequency of both local and systemic metastases, the number of samples needs to be increased before any significance can be attached to this observation.

Similar to the association of *NY-ESO-1* mRNA expression and high tumor grade, NY-ESO-1 antibody responses appear to be restricted to patients with G3 TCC. Of the 9 patients with NY-ESO-1 antibody, 5 of the patients had CIS, a flat superficial tumor considered to be a progenitor for invasive tumors (17). BCG has been found to be an effective therapy for patients with CIS and superficial bladder cancer, with prolonged protection from tumor recurrence occurring in a significant proportion of BCG-treated patients (20). In fact, 3 of 5 NY-ESO-1 antibody-positive CIS patients in our series received BCG therapy and have remained tumor free for 5 months to 5 years. One patient, TCC44, underwent transurethral resection of tumor followed by intravesical BCG instillation in 1995 and has been healthy without tumor recurrence since that time. A serum sample from this patient was first obtained in October 1997 and found to be NY-ESO-1 antibody positive, despite the long tumor-free interval. This observation of persistent NY-ESO-1 antibody in the absence of tumor stands in contrast to our experience with melanoma and other cancer types, where the presence of NY-ESO-1 antibody is clearly antigen driven and disappears with tumor removal or therapy-inducing tumor regression (15). The basis for: (a) the strong immunogenicity for NY-ESO-1 presented by CIS; (b) the persistence of NY-ESO-1 antibody in CIS

Table 3 *TNM classification of TCC in patients with NY-ESO-1 antibody*

Patient	Primary tumor	Metastasis	
		Regional lymph node	Systemic
TCC 11	CIS	-	-
TCC 18 <sup>a</sup>	T1 + CIS	-	-
TCC 21	T3 <sup>b</sup>	-	-
TCC 22 <sup>a</sup>	T4	+	+
TCC 25	CIS	-	-
TCC 38	T2	-	-
TCC 44 <sup>b</sup>	CIS	-	-
TCC 84 <sup>a</sup>	T1 + CIS	-	-
TCC 91	T3 <sup>b</sup>	-	-

<sup>a</sup> Tumor specimens with NY-ESO-1 expression confirmed by immunohistochemical staining. Frozen specimens from TCC18 for RT-PCR were not available.

<sup>b</sup> Patient TCC44 has been tumor free since September 1995 after transurethral resection of bladder tumor followed by BCG immunotherapy. Serum for NY-ESO-1 antibody test was obtained in October 1997. Sera from the 8 other patients were obtained at the time of surgical resection of their tumors.

patients after successful therapy; and (c) the possibility that NY-ESO-1 immunity might be involved in delaying tumor progression and mediate the therapeutic activity of BCG in superficial bladder cancer is an important topic requiring careful study.

In a recent study by Jäger *et al.* (14) involving melanoma patients, a humoral immune response to NY-ESO-1 was predictive of a strong CD8 T cell response to NY-ESO-1-derived peptides, as measured by tetramer, enzyme-linked immunospot, and cytotoxicity. We are now conducting a comparable analysis of patients with TCC to determine whether there is a link between the humoral and cellular immune response to NY-ESO-1 in this patient population. Because of the strong immunogenicity of NY-ESO-1, there is considerable interest in vaccine strategies targeting this antigen, and a variety of NY-ESO-1 constructs, including NY-ESO-1 peptides, protein, DNA, and viral and bacterial vectors, are being prepared for clinical evaluation. A recent clinical vaccine trial with HLA-A2-restricted NY-ESO-1 peptides has shown that these peptides can elicit a strong CD8 T-cell response in NY-ESO-1-immunized patients (21). Because of the high frequency of *NY-ESO-1* expression in TCC and the increasing awareness of the limitation of current chemotherapy against TCC, TCC represents a challenging tumor type to test the effectiveness of NY-ESO-1 vaccines.

**Acknowledgments**

We thank M. Isobe for excellent technical assistance and J. Mizuuchi for preparation of the manuscript.

**References**

- Boon, T., and Old, L. J. Tumor antigens. *Curr. Opin. Immunol.*, 9: 681-683, 1997.
- Van der Bruggen, P., Traversari, C., Chomez, P., Lurquin, C., De Plaen, E., Van Den Eynde, B., Knuth, A., and Boon, T. A gene encoding an antigen recognized by cytolytic T lymphocytes on a human melanoma. *Science (Wash. DC)*, 254: 1643-1647, 1991.
- Chen, Y.-T., Scanlan, M. J., Sahin, U., Türeci, Ö., Güre, A. O., Tsang, S., Williamson, B., Stockert, E., Pfreundschuh, M., and Old, L. J. A testicular antigen aberrantly expressed in human cancers detected by autologous antibody screening. *Proc. Natl. Acad. Sci. USA*, 94: 1914-1918, 1997.
- Brichard, V., Van Pel, A., Wölfel, T., Wölfel, C., De Plaen, E., Lethé, B., Coulie, P. G., and Boon, T. The tyrosinase gene codes for an antigen recognized by autologous cytolytic T lymphocytes on HLA-A2 melanomas. *J. Exp. Med.*, 178: 489-495, 1993.
- Kawakami, Y., Eliyahu, S., Delgado, C. H., Robbins, P. F., Sakaguchi, K., Appella, E., Yannelli, J. R., Adema, G. J., Miki, T., and Rosenberg, S. A. Identification of a human melanoma antigen recognized by tumor-infiltrating lymphocytes associated with *in vivo* tumor rejection. *Proc. Natl. Acad. Sci. USA*, 91: 6458-6462, 1994.
- Kawakami, Y., Eliyahu, S., Delgado, C. H., Robbins, P. F., Rivoltini, L., Topalian, S. L., Miki, T., and Rosenberg, S. A. Cloning of the gene coding for a shared human melanoma antigen recognized by autologous T cells infiltrating into tumor. *Proc. Natl. Acad. Sci. USA*, 91: 3515-3519, 1994.
- Coulie, P. G., Brichard, V., Van Pel, A., Wölfel, T., Schneider, J., Traversari, C., Mattei, S., De Plaen, E., Lurquin, C., Szikora, J.-P., Renauld, J.-C., and Boon, T. A




Fig. 2. NY-ESO-1 antibody in patients with G3 TCC. Serially diluted sera from 5 patients with G3 TCC (●), serum from an NY-ESO-1-positive patient (○), and sera from 5 healthy volunteers (□) are shown. ELISA tests with recombinant NY-ESO-1 protein are also shown.

Table 2 *NY-ESO-1 antibody response in 124 patients with transitional cell carcinoma<sup>a</sup>*

Tumor grade	Antibody positive/sera tested
G1	0/5 (0%)
G2	0/47 (0%)
G3	9/72 <sup>b</sup> (12.5%)

<sup>a</sup> In this series of 124 patients with TCC, both frozen tumors and sera were available from 45 patients for NY-ESO-1 RT-PCR and antibody typing. Thirteen of 45 tumors were positive for *NY-ESO-1* mRNA, and 2 of the 13 patients with *NY-ESO-1*-positive tumors had NY-ESO-1 antibody. No NY-ESO-1 antibody was found in the 32 patients with *NY-ESO-1*-negative tumors.

<sup>b</sup> The difference in NY-ESO-1 antibody frequency in patients with high-grade (G3; 9 of 72) and low-grade (G1 and G2; 0 of 52) tumors was statistically significant ( $P < 0.05$ ). Statistical analyses were performed by  $\chi^2$  test.

- new gene coding for a differentiation antigen recognized by autologous cytolytic T lymphocytes on HLA-A2 melanomas. *J. Exp. Med.*, 180: 35–42, 1994.
8. Robbins, P. F., El-Gamil, M., Li, Y. F., Kawakami, Y., Loftus, D., Appella, E., and Rosenberg, S. A. A mutated  $\beta$ -catenin gene encodes a melanoma-specific antigen recognized by tumor infiltrating lymphocytes. *J. Exp. Med.*, 183: 1185–1192, 1996.
  9. Mandruzzato, S., Brasseur, F., Andry, G., Boon, T., and van der Bruggen, P. A CASP-8 mutation recognized by cytolytic T lymphocytes on a human head and neck carcinoma. *J. Exp. Med.*, 186: 785–793, 1997.
  10. Yoshino, I., Goedegebuure, P. S., Peoples, G. E., Parikh, A. S., DiMaio, J. M., Lyerly, H. K., Gazdar, A. F., and Eberlein, T. J. HER2/neu-derived peptides are shared antigens among human non-small cell lung cancer and ovarian cancer. *Cancer Res.*, 54: 3387–3390, 1994.
  11. Old, L. J., and Chen, Y-T. New paths in human cancer serology. *J. Exp. Med.*, 187: 1163–1167, 1998.
  12. Chen, Y-T., Scanlan, M. J., Obata, Y., and Old, L. J. Identification of human tumor antigens by serological expression cloning (SEREX). *In: V. T. Devita, S. Hellman, and S. A. Rosenberg (eds.), Cancer Vaccine: Cancer Antigens*, pp. 557–570. Philadelphia: J. B. Lippincott Co., 1999.
  13. Stockert, E., Jäger, E., Chen, Y-T., Scanlan, M. J., Gout, I., Karbach, J., Arand, M., Knuth, A., and Old, L. J. A survey of the humoral immune response of cancer patients to a panel of human tumor antigens. *J. Exp. Med.*, 187: 1349–1354, 1998.
  14. Jäger, E., Nagata, Y., Gnjatic, S., Wada, H., Stockert, E., Karbach, J., Dunbar, P. R., Lee, S. Y., Jungbluth, A., Jäger, D., Arand, M., Ritter, G., Cerundolo, V., Dupont, B., Chen, Y-T., Old, L. J., and Knuth, A. Monitoring CD8 T cell responses to NY-ESO-1: correlation of humoral and cellular immune responses. *Proc. Natl. Acad. Sci. USA*, 97: 4760–4765, 2000.
  15. Jäger, E., Stockert, E., Zidianakis, Z., Chen, Y-T., Karbach, J., Jäger, D., Arand, M., Ritter, G., Old, L. J., and Knuth, A. Humoral immune responses of cancer patients against “cancer-testis” antigen NY-ESO-1: correlation with clinical events. *Int. J. Cancer*, 84: 506–510, 1999.
  16. Bane, B. L., Rao, J. Y., and Hemstreet, G. P. Pathology and staging of bladder cancer. *Semin. Oncol.*, 23: 546–570, 1996.
  17. Epstein, J. I., Amin, M. B., Reuter, V. R., Mostofi, F. K., and the Bladder Consensus Conference Committee. The World Health Organization/international society of urological pathology consensus classification of urothelial (transitional cell) neoplasms of the urinary bladder. *Am. J. Surg. Pathol.*, 22: 1435–1448, 1998.
  18. Spruck, C. H., III, Ohneseit, P. F., Gonzalez-Zulueta, M., Esrig, D., Miyao, N., Tsai, Y. C., Lerner, S. P., Schmütte, C., Yang, A. S., Cote, R., Dubeau, L., Nichols, P. W., Hermann, G. G., Steven, K., Horn, T., Skinner, D. G., and Jones, P. A. Two molecular pathways to transitional cell carcinoma of the bladder. *Cancer Res.*, 54: 784–788, 1994.
  19. Urogenital tumours. *In: L. H. Sobin and C. Wittekind (eds.), TNM Classification of Malignant Tumours*, Ed. 5, pp. 183–190. New York: Wiley-Liss, Inc., 1997.
  20. Lamm, D. L. Long-term results of intravesical therapy for superficial bladder cancer. *Urol. Clin. North Am.*, 19: 573–580, 1992.
  21. Jäger, E., Gnjatic, S., Nagata, Y., Stockert, E., Jäger, D., Karbach, J., Neumann, A., Rieckenberg, J., Chen, Y-T., Ritter, G., Hoffman, E., Arand, M., Old, L. J., and Knuth, A. Induction of primary NY-ESO-1 immunity: CD8<sup>+</sup> T lymphocyte and antibody responses in peptide-vaccinated patients with NY-ESO-1<sup>+</sup> cancers. *Proc. Natl. Acad. Sci. USA*, 97: 12198–12203, 2000.

# Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

## NY-ESO-1 Expression and Immunogenicity Associated with Transitional Cell Carcinoma: Correlation with Tumor Grade

Takushi Kurashige, Yuji Noguchi, Takashi Saika, et al.

*Cancer Res* 2001;61:4671-4674.

**Updated version** Access the most recent version of this article at:  
<http://cancerres.aacrjournals.org/content/61/12/4671>

**Cited articles** This article cites 19 articles, 14 of which you can access for free at:  
<http://cancerres.aacrjournals.org/content/61/12/4671.full#ref-list-1>

**Citing articles** This article has been cited by 24 HighWire-hosted articles. Access the articles at:  
<http://cancerres.aacrjournals.org/content/61/12/4671.full#related-urls>

**E-mail alerts** [Sign up to receive free email-alerts](#) related to this article or journal.

**Reprints and Subscriptions** To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at [pubs@aacr.org](mailto:pubs@aacr.org).

**Permissions** To request permission to re-use all or part of this article, use this link  
<http://cancerres.aacrjournals.org/content/61/12/4671>.  
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.