

Activated Granulocytes and Granulocyte-derived Hydrogen Peroxide Are the Underlying Mechanism of Suppression of T-Cell Function in Advanced Cancer Patients¹

Jan Schmielau² and Olivera J. Finn

Department of Molecular Genetics and Biochemistry, University of Pittsburgh School of Medicine, Pittsburgh, Pennsylvania 15261

ABSTRACT

Impaired T-cell function in patients with advanced cancer has been a widely acknowledged finding, but mechanisms reported thus far are those primarily operating in the tumor microenvironment. Very few mechanisms have been put forth to explain several well-described defects in peripheral blood T cells, such as reduction in expression of signaling molecules, decreased production of cytokines, or increased apoptosis. We have closely examined the peripheral blood mononuclear cell (PBMC) samples derived from patients and healthy individuals, and we have observed an important difference that may underlie the majority of reported defects. We observed that in samples from patients only, an unusually large number of granulocytes copurify with low density PBMCs on a density gradient rather than sediment, as expected, to the bottom of the gradient. We also show that activating granulocytes from a healthy donor with *N*-formyl-L-methionyl-L-leucyl-L-phenylalanine could also cause them to sediment aberrantly and copurify with PBMCs, suggesting that density change is a marker of their activation. To confirm this, we looked for other evidence of *in vivo* granulocyte activation and found it in drastically elevated plasma levels of 8-isoprostane, a product of lipid peroxidation and a marker of oxidative stress. Reduced T-cell receptor ζ chain expression and decreased cytokine production by patients' T cells correlated with the presence of activated granulocytes in their PBMCs. We showed that freshly obtained granulocytes from healthy donors, if activated, can also inhibit cytokine production by T cells. This action is abrogated by the addition of the hydrogen peroxide (H₂O₂) scavenger, catalase, implicating H₂O₂ as the effector molecule. Indeed, when added alone, H₂O₂ could suppress cytokine production of normal T cells. These findings indicate that granulocytes are activated in advanced cancer patients and that granulocyte-derived H₂O₂ is the major cause of severe systemic T-cell suppression.

INTRODUCTION

Despite increasing efforts, studies into tumor-induced immunosuppression have not provided plausible mechanisms that would explain the severe functional impairment in all circulating T cells seen in patients with advanced cancer (1). Although a growing number of tumor-specific antigens are being identified as potential targets that could be recognized by the immune system, cancer patients largely fail to mount an effective immune response against them. A failed T-cell response early in tumor development appears to be predominantly antigen specific (2). In some instances, tumor antigens are not sufficiently processed or presented to T cells because of the nature of the particular antigen (3) or defects in antigen-presenting cells (4). It has also been reported that proliferation of tumor-specific T cells could be hindered or the tumor-specific response could be switched to

an unfavorable Th²³ that does not support development of cytotoxic effector T cells from precursors stimulated at the tumor site (5). Even if competent T cells are made and capable of infiltrating the tumor, it has been reported that they can undergo apoptosis induced by Fas ligand expressed on the tumor cells (6). Although these mechanisms might explain tumor-specific immunosuppression, they are often restricted to the immediate tumor environment and fail to explain the more generalized immunosuppression that develops with tumor progression when patients stop responding even to common recall antigens (7). The first molecular mechanism described that potentially underlies these defects is the reduced expression of the TCR ζ chain and TCR-associated protein tyrosine kinases in splenocytes from tumor-bearing mice (8). Subsequent studies with cancer patients indeed showed a reduced TCR ζ chain expression that correlated with the tumor state of the patient (9–11). This defect was later correlated with antitumor responses in melanoma patients who underwent immunotherapy with IL-2 (12). Recently, one possible reason for these defects in TCR expression has been proposed to be the absence of L-arginine, known to be depleted in certain disease states (13) and accompanied by a decrease in T-cell proliferation (14, 15). A defect in the translocation of transcription factors in T cells, such as nuclear factor- κ B, has been demonstrated and implicated in tumor-induced suppression (9, 16). As a likely consequence of both defects, an impaired cytokine production by T cells has been shown in patients with a variety of tumors (5, 17–21). Circulating suppressor cells or suppressor factors have been postulated as the cause for some of these defects in T-cell function. One type of candidate suppressor cells investigated have been the macrophages that can undergo activation in the vicinity of the tumor and produce factors, such as H₂O₂, that suppress various other immune effector cells (22). Macrophages, however, are not circulating cells, and their suppressive function is still expected to be limited to the effector cells in the tumor microenvironment.

Here we propose a mechanism that likely underlies all of the others described thus far and that we consider the common cause for most of the observed defects in immune function in advanced cancer patients. This mechanism involves activation of granulocytes and oxidative stress mediated by H₂O₂ that they produce. Unlike macrophages that are primarily found in tissues, granulocytes are abundantly present in circulation. Phagocytosis by granulocytes is supported by an oxidative burst that releases reactive oxygen species from membrane-bound enzymes and facilitates killing of microorganisms. We offer evidence of *in vivo* granulocyte activation in cancer patients that correlates with the inhibition of TCR ζ chain expression and cytokine production by their T cells. We show that cocubation of T cells with activated granulocytes *in vitro* leads to the inhibition of cytokine production and that the mediator of that inhibition is H₂O₂.

Received 12/15/00; accepted 4/11/01.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported by a grant from the "Deutsche Krebshilfe" (to J. S.) and NIH Grant RO1 CA56103 and a grant from The Nathan Aronson Fund for Pancreatic Cancer Research (to O. J. F.).

² To whom requests for reprints should be addressed, at Medizinische Universitätsklinik, Knappschaftskrankenhaus, In der Schornau 23–25, 44892 Bochum, Germany. E-mail: jan.schmielau@ruhr-uni-bochum.de.

³ The abbreviations used are: Th1 and Th2, T helper types 1 and 2; TCR, T-cell receptor; IL, interleukin; PBMC, peripheral blood mononuclear cell; FMLP, *N*-formyl-L-methionyl-L-leucyl-L-phenylalanine; PMA, phorbol myristate acetate; TNF, tumor necrosis factor; PFA, paraformaldehyde.

MATERIALS AND METHODS

Blood Samples. Blood samples were obtained under an Institutional Review Board-approved protocol from patients with metastatic adenocarcinomas of the pancreas, colon, and breast, seen at the University of Pittsburgh Cancer Institute. All radiation or chemotherapy had been concluded at least 4 weeks prior to the blood drawing. PBMCs were purified on a density gradient using a lymphocyte separation medium (ICN, Aurora, OH) and stored at -80°C in fetal bovine serum containing medium with 10% DMSO.

Granulocyte and T-Cell Assays. Granulocytes were purified from peripheral blood with dextran sedimentation as described (23). PBMCs were collected by centrifugation in lymphocyte separation medium (ICN), and residual RBCs were lysed in a hypotonic saline solution. To obtain activated granulocytes, whole blood was incubated for 1 h with FMLP (Sigma Chemical Co., St. Louis, MO) prior to density gradient centrifugation. Cells were stained with anti-CD15 antibody (Becton Dickinson, Mountain View, CA) for flow cytometry or used for cytopspins stained with Diff-Quick (Baxter, McGraw Park, IL). T cells were activated for 4 h with $20\ \mu\text{g/ml}$ PMA and $1\ \mu\text{M}$ ionomycin (Sigma Chemical Co.).

PBMC samples were analyzed for TCR ζ expression and cytokine production by flow cytometry as described (24). For intracellular cytokine staining, 2 mM monensin (Sigma Chemical Co.) was added to T cells for 4 h. Cells were fixed with 2% PFA³ and permeabilized with FACS buffer (PBS supplemented with 5% FBS and 0.1% sodium azide) containing 0.1% saponin (all from Sigma Chemical Co.). An anti-TCR ζ chain antibody, TCR- ζ , TIA-2 (Coulter, Fullerton, CA) was used for indirect staining prior to a secondary goat antimouse R-phycoerythrin-conjugated antibody (Biosource, Camarillo, CA). Antibodies against the cytokines IFN- γ , TNF- α , IL-2, or IL-4 were obtained from PharMingen (San Diego, CA). Live cells were stained with antibodies against CD11b, CD14, and CD15 on granulocytes and monocytes and antibodies against CD3 and CD8 (Becton Dickinson) on T cells. Analysis was done on a FACS-Calibur (Becton Dickinson).

For granulocyte/T-cell coculture assays, both cell types were purified from the same blood sample. Granulocytes (4×10^6) were seeded into the lower compartment of a six-well transwell system (Costar, Cambridge, MA), and 2×10^6 PBMCs were either placed into the upper compartment or directly mixed with the granulocytes in a final volume of 5 ml. The experiment, involving freezing, fixation, or addition of bovine catalase (Sigma Chemical Co.), was performed with 7.5×10^5 T cells and granulocytes in a 24-well plate (Costar). Granulocytes were either frozen overnight in FBS with 10% DMSO at -80°C or fixed in 2% PFA/PBS for 20 min prior to the assay. For experiments involving H₂O₂, PBMCs were treated for 5 min, followed by an additional 5-min incubation with catalase at a final concentration of 1000 units/ml. Dose-response experiments with H₂O₂ were done in the absence of FBS.

Detection of F_{2 α} -Isoprostane Plasma samples that had been stored at -80°C were used in an enzyme immunoassay that detects F_{2 α} -isoprostane (Cayman Chemical, Ann Arbor, MI). To measure total isoprostane, we hydrolyzed plasma samples with potassium hydroxide and performed detection after solid phase extraction with C₁₈ columns (Millipore, Milford, MA) as recommended by the manufacturer.

RESULTS

Evidence of *in Vivo* Granulocyte Activation and Oxidative Stress Damage in Cancer Patients. Density gradient centrifugation has been traditionally used as a method to separate PBMCs, such as lymphocytes, from polymorphonuclear cells, such as granulocytes. Granulocytes are expected to sediment in the cell pellet because of their high buoyant density. We noticed that in the process of purification of blood from cancer patients (Fig. 1A) but not healthy controls (Fig. 1B), unusually large numbers of granulocytes sedimented with the same density as PBMCs. This was detected by flow cytometry analysis of the composition of purified PBMCs by cell size and granularity and confirmed by staining with anti-CD15 antibody specific for granulocytes (Fig. 2). We hypothesized that aberrant sedimentation of granulocytes could be a result of a change in their density attributable to activation. To reproduce these events *in vitro*, we treated whole blood from a healthy donor with FMLP, a peptide




Fig. 1. In cancer patients but not in normal controls, granulocytes copurify with PBMCs on a density gradient. A, PBMC sample from a cancer patient. B, PBMC sample from a healthy control. Blue, CD3⁺ cells; purple, CD14⁺ cells; red, CD15⁺ cells; gray, other cells; SSC, side scatter; FSC, forward scatter.

that induces receptor-mediated granulocyte activation (25). An hour of treatment indeed led to aberrant granulocyte sedimentation. Granulocytes copurified with PBMCs in a FMLP dose-dependent manner (Fig. 2A) and represented up to 12.5% of the population, as confirmed by CD15 expression (Fig. 2B) and by specific morphology observed in the cytopspins (Fig. 2, C and D).

Analysis of additional samples from healthy individuals and cancer patients confirmed that granulocytes are virtually excluded from purified PBMCs of healthy donors but appear in high percentages in PBMCs from cancer patients (Fig. 4A). The same PBMC samples that had large numbers of activated granulocytes had mainly normal neutrophil counts (Fig. 3B). The median lymphocyte count was decreased, however, and some patients were lymphopenic (Fig. 3C). Because we hypothesized that aberrant sedimentation of granulocytes was attributable to their *in vivo* activation, we sought another *in vivo* parameter to measure to further support this hypothesis. Because activation of granulocytes results in oxidative stress and production of short-lived free radicals, we looked for biomarkers for detection of damage caused by free radicals. Isoprostanes are products of lipid peroxidation (26), and we used an enzyme immunoassay to detect F_{2 α} -isoprostane in patient plasma samples. We found exceedingly high plasma levels of isoprostane (Fig. 3D), confirming that patients had been exposed to oxidative stress.

Reduced TCR ζ Expression and Cytokine Production in Peripheral Blood T Cells. We extended our earlier studies into tumor-induced T-cell suppression in cancer patients and tested T-cell function in 19




Fig. 2. Granulocytes that copurify with PBMCs from a healthy donor on a density gradient are activated. *A*, FMLP dose-dependent increase in percentage of granulocytes (CD15+) copurifying with PBMCs. *B*, peripheral blood incubated without (----) and with 1 μ M FMLP (—) prior to density gradient centrifugation and tested for the presence of granulocytes (CD15). Cytospins of PBMCs purified before treatment (*C*) or after treatment (*D*) with FMLP.

pancreatic, 15 colon, and 1 breast cancer patient. TCR ζ chain expression was significantly reduced in all these patients compared with healthy controls ($P < 0.0001$; Fig. 3A). We activated the same T cells with PMA and ionomycin, reagents that do not depend on the TCR ζ chain for their function, and tested their ability to produce cytokines. All patients showed reduced numbers of IFN- γ -producing T cells compared with controls ($P < 0.0001$; Fig. 4B). To address the possibility that this defect may be only in the type 2 T cells, resulting in altered ratios of Th1 *versus* Th2 T cells, we also measured the type 2 cytokine IL-4. There was no compensatory increase in IL-4, the production of which was also reduced (data not shown). A similar reduction was observed in the production of TNF- α and IL-2 (data not shown).

Granulocyte-derived Hydrogen Peroxide Inhibits Cytokine Production. The observation that suppression of the TCR ζ chain expression and T-cell cytokine production coincided with signs of granulocyte activation and elevated isoprostane plasma levels in cancer patients encouraged us to seek a direct proof that activated granulocytes can be responsible for inhibition of T-cell function. We purified granulocytes from healthy donor blood by density sedimentation and obtained a >90% pure population as confirmed by CD15 expression (data not shown). We mixed these granulocytes with lymphocytes from the same donor and coincubated them under different conditions (Fig. 5A). In the absence of granulocytes, >30% of T cells produced IFN- γ when activated with PMA and ionomycin. Coactivation of lymphocytes and granulocytes had a strong inhibitory effect on IFN- γ production by T cells. Coactivation in a transwell system, where granulocytes and lymphocytes were separated by a membrane, showed that inhibition may be caused by soluble factors. The integrity of the granulocytes was important for this inhibition inasmuch as either fixation with paraformaldehyde or freezing greatly reduced their inhibitory effect (Fig. 5B).




Fig. 3. Copurification of activated granulocytes with PBMCs, white blood count, and signs of oxidative stress. In *A*, CD15⁺ cells copurified with PBMCs in patients compared with healthy controls. *B*, absolute neutrophil count in peripheral blood of patients. *C*, absolute lymphocytes count in peripheral blood of patients. *D*, plasma levels of isoprostane in patients. Normal ranges are marked with a *rectangle*.




Fig. 4. TCR ζ chain expression and IFN- γ production by T cells from cancer patients compared with healthy controls. *A*, mean fluorescence intensity of staining of all T cells from TCR ζ chain expression shown as multiples of antibody isotype control. *B*, percentage of IFN- γ -producing T cells among all T cells. *Horizontal bars*, median values.

Confirming that activated granulocytes can suppress *in vitro* IFN- γ production by T cells, we went back to the patients' samples to determine the correlation of these two parameters *in vivo*. We show that low numbers of granulocytes, predominantly found in PBMC samples from

healthy individuals, correlate with high percentages of IFN- γ -producing T cells in those individuals. Similarly, high numbers of granulocytes, predominantly found in PBMC samples from cancer patients, correlate with low percentages of IFN- γ -producing T cells (Fig. 5C).

Granulocytes limit tissue destruction by releasing protecting enzymes such as superoxide dismutase and catalase. Superoxide dismutase converts superoxide anion to H₂O₂, which in turn is degraded to water by catalase. When we added increasing concentrations of catalase to the cocultures of granulocytes and lymphocytes, most of the inhibition of IFN- γ production was abrogated (Fig. 5D). This identified H₂O₂ as the effector molecule mediating inhibition of this T-cell function. Adding H₂O₂ directly to T cells resulted in a reduced number of cytokine-producing cells, not only for IFN- γ but also TNF- α , IL-2, and IL-4, in a dose-dependent manner (Fig. 5, E and F).

DISCUSSION

We have provided evidence here for one mechanism that underlies several well-described defects in T-cell function in patients with advanced cancer. We show that cancer patients respond to the presence of the tumor with activation of granulocytes. This is likely an attempt on the part of the innate immune system to protect the organism. The chronic




Fig. 5. Granulocyte-mediated inhibition of cytokine production *in vitro*. *A*, IFN- γ -producing T cells in the absence or presence of granulocytes in a transwell system (*membrane*) or in direct contact (*direct*). *B*, IFN- γ -producing T cells in the absence or presence of untreated (\emptyset), PFA-fixed (*fixed*), or frozen and thawed (*frozen*) granulocytes. *C*, percentage of IFN- γ -producing T cells after activation, in PBMCs from patients (\bullet) and healthy controls (\circ), plotted against the percentage of contaminating granulocytes. The regression line for patient samples is shown with the 95% confidence interval ($P = 0.006$; slope, -29.6 ± 9.9). *D*, increase in the percentage of IFN- γ -producing T cells in the presence of catalase. Dose-dependent inhibition by hydrogen peroxide of IFN- γ (\bullet), TNF- α (\blacksquare), and IL-2 (\circ) (*E*) and IL-4 (\square) producing T cells (*F*) is shown.

presence of the tumor and thus chronic activation of granulocytes with the accompanying production of H_2O_2 results, ironically, in the suppression of the adaptive immune functions. Here we show specifically that the release of reactive oxygen species during the oxidative burst of granulocytes, *i.e.*, H_2O_2 , is the major contributor to a systemic T-cell dysfunction. This is supported by the observation that cancer patients showed signs of extensive granulocyte activation with massive elevation of plasma levels of 8-isoprostane, a product of lipid oxidation and a marker for oxidative stress, that exceeded measurements found in other diseases with documented oxidative stress (26–28).

Our *in vitro* experiments demonstrated directly that cocubation of granulocytes and lymphocytes could inhibit cytokine production but required the functional integrity and activation of granulocytes. The observation that granulocyte-mediated inhibition of cytokine production could be abrogated by the addition of catalase to inhibit H_2O_2 identified this product as the major soluble mediator of activated granulocyte effects. We confirmed this by adding H_2O_2 to T cells and inducing potent inhibition of cytokine production in a dose-dependent manner. Analyzing *ex vivo* cytokine production by patients' T cells, we found that the inhibition was most pronounced for IFN- γ and, although not significant, was also detectable for IL-4. We did not detect a switch from Th1 cytokines, which promote cell-mediated cytotoxicity, toward an antibody-mediated Th2-dominated responses. A decrease in cytokine production of T cells significantly correlated with the number of circulating activated granulocytes.

The activation of granulocytes could be induced by an inflammatory response or cytokines produced by the tumor. Tumor cells are known to secrete cytokines, among them are cytokines that affect granulocytes (29, 30). Possible candidates are TNF- α and IL-8, which could be elevated in the serum of cancer patients (30, 31) and stimulate the oxidative burst (32, 33). Although we have concentrated on documenting immunosuppressive effects on T cells, it is highly probable that other cells of the immune system could be equally affected by the same mediators of oxidative stress. A decrease in the function of NK cells mediated by H_2O_2 has been documented (22), whereas causes of defects in the function of dendritic cells are unknown (4). Little is known about the functional competence of B cells in cancer patients, but B cells might be affected in a similar manner to the T cells. Our data suggest that inhibition of the oxidative burst of granulocytes might be a useful treatment of cancer patients prior to attempting any form of immunotherapy.

REFERENCES

- Finke, J., Ferrone, S., Frey, A., Mufson, A., and Ochoa, A. Where have all the T cells gone? Mechanisms of immune evasion by tumors. *Immunol. Today*, 20: 158–160, 1999.
- Staveley-O'Carroll, K., Sotomayor, E., Montgomery, J., Borrello, I., Hwang, L., Fein, S., Pardoll, D., and Levitsky, H. Induction of antigen-specific T cell allergy: an early event in the course of tumor progression. *Proc. Natl. Acad. Sci. USA*, 95: 1178–1183, 1998.
- Hiltbold, E. M., Vlad, A. M., Ciborowski, P., Watkins, S. C., and Finn, O. J. The mechanism of unresponsiveness to circulating tumor antigen MUC1 is a block in intracellular sorting and processing by dendritic cells. *J. Immunol.*, 165: 3730–3741, 2000.
- Esche, C., Lokshin, A., Shurin, G. V., Gastman, B. R., Rabinowich, H., Watkins, S. C., Lotze, M. T., and Shurin, M. R. Tumor's other immune targets: dendritic cells. *J. Leukocyte Biol.*, 66: 336–344, 1999.
- Goto, S., Sato, M., Kaneko, R., Itoh, M., Sato, S., and Takeuchi, S. Analysis of Th1 and Th2 cytokine production by peripheral blood mononuclear cells as a parameter of immunological dysfunction in advanced cancer patients. *Cancer Immunol. Immunother.*, 48: 435–442, 1999.
- Strand, S., Hofmann, W. J., Hug, H., Muller, M., Otto, G., Strand, D., Mariani, S. M., Stremmel, W., Krammer, P. H., and Galle, P. R. Lymphocyte apoptosis induced by CD95 (APO-1/Fas) ligand-expressing tumor cells—a mechanism of immune evasion? *Nat. Med.*, 2: 1361–1366, 1996.
- Young, R. C., Corder, M. P., Haynes, H. A., and DeVita, V. T. Delayed hypersensitivity in Hodgkin's disease. *Am. J. Med.*, 52: 63–72, 1972.
- Mizoguchi, H., O'Shea, J. J., Longo, D. L., Loeffler, C. M., McVicar, D. W., and Ochoa, A. C. Alterations in signal transduction molecules in T lymphocytes from tumor-bearing mice. *Science (Wash. DC)*, 258: 1795–1798, 1992.
- Finke, J. H., Zea, A. H., Stanley, J., Longo, D. L., Mizoguchi, H., Tubbs, R. R., Wiltrout, R. H., O'Shea, J. J., Kudoh, S., Klein, E., Bukowski, R. M., and Ochoa, A. C. Loss of T-cell receptor ζ chain and p56^{lck} in T-cells infiltrating human renal cell carcinoma. *Cancer Res.*, 53: 5613–5616, 1993.
- Nakagomi, H., Peterson, M., Magnusson, I., Matsuda, M., Mellstedt, H., Taupin, J.-L., Vivier, E., Anderson, P., and Kiessling, R. Decreased expression of the signal-transducing ζ chains in tumor-infiltrating T-cells and NK cells of patients with colorectal carcinoma. *Cancer Res.*, 53: 5610–5612, 1993.
- Gunji, Y., Hori, S., Aoe, T., Asano, T., Ochiai, T., Isona, K., and Saito, T. High frequency of cancer patients with abnormal assembly of the T cell receptor-CD3 complex in peripheral blood T lymphocytes. *Jpn. J. Cancer Res.*, 85: 1189–1192, 1994.
- Rabinowich, H., Banks, M., Reichert, T. E., Logan, T. F., Kirkwood, J. M., and Whiteside, T. L. Expression and activity of signaling molecules in T lymphocytes obtained from patients with metastatic melanoma before and after interleukin therapy. *Clin. Cancer Res.*, 2: 1263–1274, 1996.
- Taheri, F., Ochoa, J. B., Faghiri, Z., Culotta, K., Park, H.-J., Lan, M. S., Zea, A. H., and Ochoa, A. C. L-Arginine regulates the expression of the T cell receptor ζ chain (CD3 ζ) in Jurkat cells. *Clin. Cancer Res.*, 7: 958s–965s, 2001.
- Roth, E., Steininger, R., Winkler, S., Langle, F., Grunberger, T., Fugger, R., and Muhlbacher, F. L-Arginine deficiency after liver transplantation as an effect of arginase efflux from the graft. Influence on nitric oxide metabolism. *Transplantation*, 57: 665–669, 1994.
- Ochoa, J. B., Udekwi, A. O., Billiar, T. R., Curran, R. D., Cerra, F. B., Simmons, R. L., and Peitzman, A. B. Nitrogen oxide levels in patients after trauma and during sepsis. *Ann. Surg.*, 214: 621–626, 1991.
- Ghosh, P., Sica, A., Young, H. A., Ye, J., Franco, J. L., Wiltrout, R. H., Longo, D. L., Rice, N. R., and Komschlies, K. L. Alteration in NF κ B/Rel family proteins in splenic T-cells from tumor-bearing mice and reversal following therapy. *Cancer Res.*, 54: 2969–2972, 1994.
- Elliott, L. H., Brooks, W. H., and Roszman, T. L. Cytokinetic basis for the impaired activation of lymphocytes from patients with primary intracranial tumors. *J. Immunol.*, 132: 1208–1215, 1984.
- Elsasser-Beile, U., Kolble, N., Grussenmeyer, T., Schultze-Seemann, W., Wetterauer, U., Gallati, H., Schulte Monting, J., and von Kleist, S. Th1 and Th2 cytokine response patterns in leukocyte cultures of patients with urinary bladder, renal cell and prostate carcinomas. *Tumour Biol.*, 19: 470–476, 1998.
- Guilloux, Y., Viret, C., Gervois, N., Le Dréan, E., Pandolfino, M.-C., Diez, E., and Jotereau, F. Defective lymphokine production by most CD8⁺ and CD4⁺ tumor-specific T cell clones derived from human melanoma-infiltrating lymphocytes in response to autologous tumor cells *in vitro*. *Eur. J. Immunol.*, 24: 1966–1973, 1994.
- Nieland, J. D., Loviscek, K., Kono, K., Albain, K. S., McCall, A. R., Potkul, R. K., Fisher, S. G., Velders, M. P., Petersson, M., Kiessling, R., and Kast, W. M. PBLs of early breast carcinoma patients with a high nuclear grade tumor unlike PBLs of cervical carcinoma patients do not show a decreased TCR ζ expression but are functionally impaired. *J. Immunother. Emphasis Tumor Immunol.*, 21: 317–322, 1998.
- Zea, A. H., Curti, B. D., Longo, D. L., Alvord, W. G., Strobl, S. L., Mizoguchi, H., Creekmore, S. P., O'Shea, J. J., Powers, G. C., Urba, W. J., and Ochoa, A. C. Alterations in T cell receptor and signal transduction molecules in melanoma patients. *Clin. Cancer Res.*, 1: 1327–1335, 1995.
- Kono, K., Salazar-Onfray, F., Petersson, M., Hansson, J., Masucci, G., Wasserman, K., Nakazawa, T., Anderson, P., and Kiessling, R. Hydrogen peroxide secreted by tumor-derived macrophages down-modulates signal-transducing ζ molecules and inhibits tumor-specific T cell- and natural killer cell-mediated cytotoxicity. *Eur. J. Immunol.*, 26: 1308–1313, 1996.
- Clark, R. A., and Nauseef, W. M. Isolation and functional analysis of neutrophils. *In: J. E. Coligan, A. M. Kruisbeek, D. H. Margulies, E. M. Shevach, and W. Stober (eds.), Current Protocols in Immunology*, pp. 7.23.1–7.17. New York: John Wiley & Sons, 1998.
- Schmielau, J., Nalesnik, M. A., and Finn, O. J. Suppressed TCR ζ chain expression and cytokine production in pancreatic cancer patients. *Clin. Cancer Res.*, 7: 933s–939s, 2001.
- Panaro, M. A., and Mitolo, V. Cellular responses to FMLP challenging: a mini-review. *Immunopharmacol. Immunotoxicol.*, 21: 397–419, 1999.
- Morrow, J. D., and Roberts, L. J. The isoprostanes: unique bioactive products of lipid peroxidation. *Prog. Lipid Res.*, 36: 1–21, 1997.
- Kluck, R. M., Bossy-Wetzel, E., Green, D. R., and Newmeyer, D. D. The release of cytochrome c from mitochondria: a primary site for Bcl-2 regulation of apoptosis. *Science (Wash. DC)*, 275: 1132–1136, 1997.
- Collins, C. E., Quaggiotto, P., Wood, L., O'Loughlin, E. V., Henry, R. L., and Garg, M. L. Elevated plasma levels of F2 α -isoprostane in cystic fibrosis. *Lipids*, 34: 551–556, 1999.
- Schmielau, J., Kalthoff, H., Roeder, C., and Schmiegel, W. Cytokines and the biology of pancreatic cancer. *In: J. Neoptolemos and N. Lemoine (eds.), Pancreatic Cancer: Molecular and Clinical Advances*, pp. 36–50. London: Blackwell Science, 1995.
- Chen, Z., Malhotra, P. S., Thomas, G. R., Ondrey, F. G., Duffey, D. C., Smith, C. W., Enamorado, I., Yeh, N. T., Kroog, G. S., Rudey, S., McCullagh, L., Mousa, S., Quezado, M., Herscher, L. L., and Van Waes, C. Expression of proinflammatory and angiogenic cytokines in patients with head and neck cancer. *Clin. Cancer Res.*, 5: 1369–1379, 1999.
- Wu, C. W., Chi, C. W., Hsieh, M. C., Chao, M. F., Lui, W. Y., and P'Eng, F. K. Serum tumor necrosis factor in patients with gastric cancer. *Anticancer Res.*, 18: 1597–1599, 1998.
- Menegazzi, R., Cramer, R., Patriarca, P., Scheurich, P., and Dri, P. Evidence that tumor necrosis factor α (TNF)-induced activation of neutrophil respiratory burst on biologic surfaces is mediated by the p55 TNF receptor. *Blood*, 84: 287–293, 1994.
- Jones, S. A., Wolf, M., Qin, S., Mackay, C. R., and Baggiolini, M. Different functions for the interleukin 8 receptors (IL-8R) of human neutrophil leukocytes: NADPH oxidase and phospholipase D are activated through IL-8R1 but not IL-8R2. *Proc. Natl. Acad. Sci. USA*, 93: 6682–6686, 1996.

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

Activated Granulocytes and Granulocyte-derived Hydrogen Peroxide Are the Underlying Mechanism of Suppression of T-Cell Function in Advanced Cancer Patients

Jan Schmielau and Olivera J. Finn

Cancer Res 2001;61:4756-4760.

Updated version Access the most recent version of this article at:
<http://cancerres.aacrjournals.org/content/61/12/4756>

Cited articles This article cites 29 articles, 13 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/61/12/4756.full#ref-list-1>

Citing articles This article has been cited by 97 HighWire-hosted articles. Access the articles at:
<http://cancerres.aacrjournals.org/content/61/12/4756.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerres.aacrjournals.org/content/61/12/4756>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.