

Wild-Type Breast Cancer Resistance Protein (BCRP/*ABCG2*) is a Methotrexate Polyglutamate Transporter¹

Erin L. Volk and Erasmus Schneider²

Wadsworth Center, New York State Department of Health, Albany, New York 12201, and the Department of Biomedical Sciences, School of Public Health, State University of New York at Albany, Albany, New York 12201

ABSTRACT

The existence of an ATP-dependent methotrexate (MTX) efflux mechanism has long been postulated; however, until recently, the molecular components were largely unknown. We have previously demonstrated a role for the ATP-binding cassette transporter breast cancer resistance protein (BCRP) in MTX resistance (Volk *et al.*, *Cancer Res.*, 62: 5035–5040, 2002). Resistance to this antifolate directly correlated with BCRP expression, and was reversible by the BCRP inhibitors fumitremorgin C and GF120918. Here, we provide evidence for BCRP as a MTX-transporter using an *in vitro* membrane vesicle system. Inside-out membrane vesicles were generated from both drug-selected and stably transfected cell lines expressing either wild-type (Arg482) or mutant (Gly482) variants of BCRP. In the presence of the wild-type variant of BCRP, transport of MTX into vesicles was ATP-dependent, osmotically sensitive, and inhibited by fumitremorgin C. In contrast, no transport was observed in vesicles containing the mutant form of BCRP. Wild-type BCRP appeared to have low affinity, but high capacity, for the transport of MTX, with an estimated K_m of 680 μ M and a V_{max} of 2400 pmol/mg/min. MTX accumulation was greatly decreased by mitoxantrone, a known BCRP substrate, suggesting competition for transport. Furthermore, and in contrast to the multidrug resistance-associated proteins, BCRP also transported significant amounts of polyglutamylated MTX. Although transport gradually decreased as the polyglutamate chain length increased, both MTX-Glu₂ and MTX-Glu₃ were substrates for BCRP. Together, these data demonstrate that BCRP is a MTX and MTX-polyglutamate transporter and reveal a possible mechanism by which it confers resistance.

INTRODUCTION

MTX³ was one of the first antimetabolite drugs developed and is used successfully for the treatment of various malignancies. Unfortunately, the efficacy of this chemotherapeutic agent is often compromised by the development of resistance in the cancer cells. Typically, MTX resistance is the result of alterations in its target enzyme, dihydrofolate reductase (1–4), reduced drug import by the reduced folate carrier (5–8), or altered polyglutamylation (9, 10). Reduced polyglutamylation results in lower drug retention, which leads to resistance.

Evidence for the existence of an energy-dependent MTX transport or efflux mechanism has accumulated over several decades. Early reports described a rapid, energy-dependent, MTX efflux from Sarcoma-180 cells (11), and a similar, azide-inhibitable, unidirectional export of MTX was also shown in leukemic L1210 cells (12). Furthermore, efflux of both mono- and diglutamates of MTX was demonstrated in a cervical carcinoma cell line (13). A series of reports throughout the last decade have further classified the cellular export of

MTX as an ATP-dependent, outwardly directed mechanism that transports both MTX and its metabolites (14–18).

Despite clear biochemical evidence, the mediator(s) of MTX efflux have only recently been described at the molecular level. Several members of the MRP or ABCC subfamily of the ABC transporters have been implicated in MTX export and resistance (19–21). These ATP-dependent efflux pumps have been shown to transport MTX *in vitro* and to cause low levels of MTX resistance in both transfected and drug-selected cells, but only in short-term (1–3-h) exposure assays. This limited-resistance phenotype has been attributed to the inability of the MRPs to transport MTX in its polyglutamylated form; the addition of even a single extra glutamate residue to MTX caused an almost complete abrogation of drug transport (22, 23).

In contrast, we have described high-level MTX cross-resistance to continuous (7 days) drug exposure in the mitoxantrone-selected MCF7/MX breast carcinoma cell line, which expresses large amounts of the ABC transporter BCRP (24), also known as mitoxantrone resistance protein (25), placental-specific ABC transporter (26), or *ABCG2* (27). Further characterization of these cells suggested that ATP-dependent, enhanced drug efflux caused the MTX resistance. MTX resistance in MCF7/MX cells was reversible by the BCRP inhibitors GF120918 and FTC, and levels of MTX resistance directly correlated with BCRP expression in this and multiple other cell lines of varied tissue origin (28). Together, these data suggested that BCRP can function as a MTX transporter. However, resistance to MTX occurred only in the presence of the wild-type transporter, which contains an arginine at position 482, whereas cells that overexpress a mutated BCRP (R482T and R482G) remained sensitive to MTX. Here, we expand on the association between BCRP and MTX resistance by demonstrating that wild-type, but not mutated, BCRP is able to transport MTX and its polyglutamates in an *in vitro* membrane vesicle system.

MATERIALS AND METHODS

Cell Culture

The following cell lines were cultured in improved MEM containing 10% fetal bovine serum and 1 μ g/ml ciprofloxacin at 37°C with 5% CO₂: the human breast carcinoma cell line MCF7/WT (29); its mitoxantrone-selected derivative MCF7/MX (30); transfected sublines MCF7/pcDNA3, MCF7/BCRP-Arg clone 67, MCF7/BCRP-Gly clone 52 (28); and the human colon cancer cell line S1 and its mitoxantrone-selected derivative, S1-M1–80 (25).

Vesicle Preparation

Vesicle preparation was adapted from Loe *et al.* (31). Cells were seeded onto 15-cm plates and were grown to 70% confluency, at which time they were washed twice with PBS and scraped into the residual PBS. Cells from 20 plates were combined and centrifuged at 500 \times g at 4°C for 15 min; the pellet was frozen overnight at –80°C. Inside-out membrane vesicles were prepared by thawing the cell pellet at 4°C and resuspending the cells in 7 ml of homogenization buffer containing 50 mM Tris-HCl (pH 7.5), 250 mM sucrose, 0.25 mM CaCl₂, and one EDTA-free protease inhibitor mixture tablet (Roche, Indianapolis, IN). The cell suspension was then transferred to a 50-ml round-bottomed tube and placed on ice into the nitrogen cavitation chamber (Parr Instruments, Moline, IL). Under constant stirring, cells were saturated with

Received 12/31/02; revised 5/6/03; accepted 6/30/03.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

¹ Supported by NIH Grant CA72455 (to E. S.).

² To whom requests for reprints should be addressed, at Wadsworth Center, Empire State Plaza, Albany, NY 12201. Phone: (518) 474-2088; Fax: (518) 474-1850; E-mail: schneid@wadsworth.org.

³ The abbreviations used are: MTX, methotrexate; ABC, ATP-binding cassette; BCRP, breast cancer resistance protein; FTC, fumitremorgin C; MRP, multidrug resistance-associated protein; PVDF, polyvinylidene difluoride.

nitrogen gas at a pressure of 1,250 p.s.i. for 20 min. After release of the pressure, membrane suspensions were centrifuged at $1,000 \times g$ at 4°C for 15 min. The supernatant was overlaid on a sucrose cushion containing 35% sucrose, 10 mM Tris-HCl (pH 7.5), 1 mM EDTA, and was centrifuged at $209,000 \times g$ at 4°C for 120 min. The opaque interface was collected and diluted $\sim 1:30$ into TS buffer containing 10 mM Tris-HCl (pH 7.5) and 250 mM sucrose and was centrifuged at $123,000 \times g$ at 4°C for 40 min. The vesicle-containing pellet was resuspended in 50 mM Tris-HCl (pH 7.5) and 250 mM sucrose, and dispersed 20 times through a 27-gauge needle. Vesicle preparations were assayed for protein concentration using the Bradford method (Bio-Rad, Hercules, CA), and membrane orientation was determined by measuring the activity of the endo-enzyme glyceraldehyde-3-phosphate dehydrogenase (32) in the absence and presence of 0.1% Triton X-100.

Western Blot Analysis

Immunodetection of BCRP in the membrane vesicles was performed as described previously (28). Briefly, 50 μg of the protein of membrane vesicles were electrophoresed on a 10% polyacrylamide gel and were transferred to a PVDF membrane (Millipore, Bedford, MA). BCRP-specific signals were detected using a polyclonal rabbit anti-BCRP primary antibody (kindly provided by Dr. Susan Bates, National Cancer Institute, Bethesda, MD), horseradish peroxidase-conjugated goat antirabbit secondary antibody (Bio-Rad), and enhanced chemiluminescence (Pierce, Rockford, IL).

Vesicular Drug Accumulation Assays

Vesicular drug accumulation assays were adapted from Saxena and Henderson (33). All of the reaction mixtures were prepared as 20- μl samples containing the following: 0.1 mg/ml creatine kinase, 10 mM phosphocreatine (both from Sigma, St. Louis, MO), 10 mM Tris-HCl (pH 7.4), 10 mM magnesium chloride, 250 mM sucrose, 400 μM [^3H]methotrexate (Moravek Biochemicals, Brea, CA), 4 mM ATP or AMP, and 5 μg of vesicles. Reaction mixtures were incubated at 37°C (0 time points were incubated at 4°C), and transport was terminated by dilution into 4 ml of ice-cold TS buffer. Vesicles were then separated from free drug by filtration of the reaction mixture through a glass microfiber filter (GF-F; Whatman, Fairfield, NJ) and rinsing five times with TS buffer. The amount of drug in the vesicles that remained on the filters was quantitated by liquid scintillation counting after 24 h in liquid-scintillation mixture. Unless otherwise indicated, all of the replicate assays were done with separate vesicle preparations.

Time Course Assays. Reactions were incubated for 1, 2, 5, 10, 15, 30, 45, and 60 min at 37°C . Time 0 was incubated at 4°C .

Osmotic Sensitivity Assays. MTX accumulation was measured in the presence of 0.25, 0.5, 0.67, 1.0, or 1.14 M sucrose and 200 μM [^3H]methotrexate, in the presence or absence of ATP at 60 min.

Inhibitor Assays. MTX accumulation was measured in the presence or absence of 5 μM FTC at 60 min.

Kinetic Studies. Drug accumulation was measured after 2-min exposure to 0, 20, 50, 100, 200, 500, 1000, and 2000 μM [^3H]methotrexate.

Competition Assays. MTX accumulation was measured in the presence of 0, 10, 20, 50, and 100 μM mitoxantrone at 60 min.

Accumulation of MTX polyglutamates. Transport of long-chain polyglutamates was assayed by incubation of vesicles in the presence of 400 μM [^3H]methotrexate,⁴ [^3H]methotrexate-Glu₂, [^3H]methotrexate-Glu₃, or [^3H]methotrexate-Glu₄ (Moravek Biochemicals) for 60 min in either the presence or the absence of 5 μM FTC.

Calculation of Net Accumulation. Unless otherwise indicated, all values shown correspond to net accumulation. Net MTX accumulation was determined by measurement of drug uptake in the presence of ATP or AMP at 4°C , and subtracting these background values (T_0) from each respective measurement taken in the presence of ATP or AMP at 37°C , at time points ranging from 1 to 60 min. For example:

$$\text{Net accumulation at 60 min} = (T_{60\text{ATP}} - T_0\text{ATP}) - (T_{60\text{AMP}} - T_0\text{AMP})$$

⁴ MTX has an integral glutamate and, therefore, corresponds to MTX-Glu₁.

RESULTS

BCRP Expression in Membrane Vesicles. Nitrogen cavitation was used to generate inside-out membrane vesicles from the following cells: parental MCF7/WT and S1 and their respective mitoxantrone-selected derivatives MCF7/MX and S1-M1-80 and stable transfectants MCF7/pcDNA3, MCF7/BCRP-Arg, and MCF7/BCRP-Gly. The inverted orientation of the membrane in the vesicles was determined by assaying for the activity of the endo-enzyme glyceraldehyde-3-phosphate dehydrogenase in the presence and absence of detergent. All of the preparations were found to be, on average, 70% (range, 50–96%) inside-out (data not shown).

To confirm the presence of BCRP in the inside-out vesicles, a Western blot analysis was performed. As expected, high levels of BCRP were detected in the vesicles from drug-selected MCF7/MX and S1-M1-80 cells, whereas moderate levels were observed in the vesicles derived from the BCRP-transfected cells MCF7/BCRP-Arg and MCF7/BCRP-Gly (Fig. 1). The apparent slight overexpression of BCRP in the MCF7/BCRP-Gly clone compared with the MCF7/BCRP-Arg clone is also seen in glyceraldehyde-3-phosphate dehydrogenase used as a loading control. It most likely can be attributed to slight differences in the amount of protein loaded and, therefore, does not represent a true difference in the level of BCRP in these vesicles. Using a serial dilution of membranes from MCF7/MX cells, we estimated that these cells expressed approximately 300–400-fold more BCRP than the transfectants (data not shown); this value corresponded well to that obtained by quantitative reverse transcription-PCR (28). Little or no BCRP was detected in vesicles from the parental cell lines MCF7/WT and S1, or from the control transfectants MCF7/pcDNA3. Together, these data indicated that the *in vitro* system was suitable for the intended studies; membrane vesicles were inverted and BCRP levels in vesicles were consistent with those observed in whole cells.

MTX Accumulation into Vesicles. To establish BCRP as an energy-dependent transporter of MTX, steady-state drug accumulation into vesicles from each cell line was measured in the presence of 4 mM either ATP or AMP. As shown in Fig. 2A, high levels of MTX accumulation were found in MCF7/MX and MCF7/BCRP-Arg vesicles, both of which express the arginine variant of BCRP, in the presence of ATP, but not in the presence of AMP. In contrast, vesicles generated from S1-M1-80 or MCF7/BCRP-Gly cells, which express the glycine variant of BCRP, displayed no significant increase in MTX accumulation in the presence of ATP compared with the increase in the presence of AMP. As expected, vesicles derived from MCF7/WT- and control-transfected cells exhibited little or no MTX transport in the presence of either ATP or AMP. Thus, it appeared that vesicles derived from cells containing the arginine variant of BCRP were able to accumulate MTX, and that accumulation was ATP




Fig. 1. Western blot analysis of BCRP expression. Membrane vesicles (50 μg) derived from MCF7/WT, MCF7/pcDNA3, MCF7/MX, MCF7/BCRP-Arg, MCF7/BCRP-Gly, S1, and S1-M1-80 cells were electrophoresed on a 10% polyacrylamide gel and transferred onto a PVDF membrane. BCRP and the loading control glyceraldehyde-3-phosphate-dehydrogenase-specific signals were detected using enhanced chemiluminescence. *kDa*, M_r (MW) in thousands.




Fig. 2. ATP-dependent MTX accumulation into BCRP-expressing vesicles. A, MTX accumulation into vesicles from MCF7/WT, MCF7/MX, MCF7/pcDNA3, MCF7/BCRP-Arg, S1-M1-80, and MCF7/BCRP-Gly was measured after 60-min exposure to 400 μM [^3H]methotrexate at 37°C in the presence of ATP (black columns) or AMP (dotted columns). B and C, time dependence of [^3H]methotrexate accumulation was measured in the presence of ATP or AMP at 0, 1, 2, 5, 10, 15, 30, 45, and 60 min. The data represent net accumulation into vesicles derived from drug-selected and parental cell lines (B), MCF7/MX (●), S1-M1-80 (▲), and MCF7/WT (■), and stably transfected cell lines (C), MCF7/BCRP-Arg (◆), MCF7/BCRP-Gly (▼) and MCF7/pcDNA3 (*). The data represent the mean \pm SE of three independent experiments performed using three separate vesicle preparations from MCF7/MX, MCF7/BCRP-Arg, and MCF7/BCRP-Gly cells, and the mean of two experiments for MCF7/WT and S1-M1-80 vesicles. (The scale of the Y axis is different between panels B and C.)

dependent. In contrast, no accumulation was observed in vesicles containing BCRP with a glycine at position 482. These results were consistent with the previously observed energy-dependence of MTX efflux and the position-482 substrate specificity of BCRP in whole cells (24, 28, 34, 35).

There was a significant difference in the absolute amount of MTX accumulation in the drug-selected MCF7/MX vesicles relative to the BCRP-transfected MCF7/BCRP-Arg vesicles. Steady-state levels of MTX were \sim 8-fold lower in vesicles derived from the transfected cells, which is commensurate with the reduced levels of BCRP expression in the MCF7/BCRP-Arg cells relative to the MCF7/MX cells. In contrast, despite very high levels of BCRP in vesicles from S1-M1-80 cells, no accumulation of MTX was detected.

To further examine MTX transport via BCRP, we measured time-dependent drug accumulation in vesicles derived from both drug-selected and stably transfected cell lines. The MCF7/MX vesicles displayed an ATP-dependent accumulation of MTX that increased with time and was linear for at least two min (Fig. 2B). In contrast, no transport was detected in vesicles derived from either S1-M1-80 or parental cells, even after a 60-min drug exposure. The S1-M1-80 vesicles were, however, capable of ATP-dependent mitoxantrone accumulation (data not shown), indicating that the vesicles were functional. This provides further evidence that the amino acid at codon 482 is critical for BCRP substrate specificity. Similarly, MTX accumulation increased over time in the vesicles derived from the MCF7/BCRP-Arg cells, but no drug uptake was observed in either MCF7/BCRP-Gly vesicles or vesicles from control transfectants (Fig. 2C). Together, these data demonstrated that MTX transport was both energy and time dependent and appeared to occur only in the presence of the wild-type (Arg482) variant of BCRP.

Osmotic Sensitivity. To assess whether the ATP-dependent MTX accumulation was caused by drug import into the lumen of the vesicles or was attributable to nonspecific binding to the membranes, we determined its osmotic sensitivity. ATP-dependent uptake of MTX into the MCF7/MX vesicles was measured under various osmotic conditions. Drug accumulation was linear, relative to the reciprocal of the sucrose concentration in the presence of ATP, whereas no drug accumulation was observed in the absence of ATP under otherwise identical reaction conditions (Fig. 3). These data demonstrated that the apparent intravesicular drug accumulation observed was, indeed, attributable to transport into the lumen of the membrane vesicles, rather than to nonspecific binding.

Inhibition of MTX Transport. FTC has been shown to inhibit BCRP (36–38). Therefore, to confirm that the MTX transport into




Fig. 3. Osmotic sensitivity. MTX uptake into MCF7/MX vesicles was measured after 45-min exposure to 200 μM [^3H]methotrexate in the presence of increasing sucrose concentrations and in the presence (●) or absence (○) of 4 mM ATP.

vesicles was mediated by BCRP, MTX accumulation into vesicles derived from MCF7/MX and MCF7/BCRP-Arg cells was measured in the presence of 5 μM FTC. Essentially complete inhibition of MTX accumulation was observed in both sets of vesicles (Fig. 4), indicating that BCRP is the transporter of MTX in this system.

Kinetics of MTX Transport by BCRP. To determine the kinetic parameters of MTX transport via BCRP, concentration-dependent uptake of MTX was measured in the MCF7/MX-derived vesicles. Accumulation was measured after a 2-min exposure to MTX in concentration that ranged from 0 to 2000 μM , in the presence of either ATP or AMP. As shown in Fig. 5, the velocity of substrate transport increased according to Michaelis-Menten kinetics in the presence of increasing levels of drug. MTX transport by BCRP was found to have a K_m of $681 \pm 280 \mu\text{M}$ and a V_{max} of $2384 \pm 445 \text{ pmol/mg/min}$. Thus, BCRP appeared to be a low-affinity, high-capacity transporter of MTX.

Substrate Competition. Mitoxantrone is the primary substrate for BCRP; almost all BCRP-overexpressing cell lines display high levels of mitoxantrone efflux and resistance (39, 40). Therefore, to further investigate the role of BCRP in MTX transport, we assayed MTX




Fig. 4. MTX accumulation in the presence of a BCRP inhibitor. Net [^3H]methotrexate accumulation into vesicles from the MCF7/MX (black columns) and MCF7/BCRP-Arg (dotted columns) cells was assayed in the presence or absence of 5 μM FTC. Experiments were performed as described in "Materials and Methods." Each column represents the mean \pm SE of three experiments performed with three separate vesicle preparations.




Fig. 5. Kinetics of MTX transport. MCF7/MX vesicles were exposed for 2 min to various concentrations of [^3H]methotrexate, ranging from 0 to 2000 μM , in the presence of either ATP or AMP. Each data point is the mean \pm SE of four independent experiments performed with three separate vesicle preparations. The graph represents a best fit Michaelis-Menten plot of the net initial velocity relative to increasing MTX concentrations, determined using the Prism3 software package.




Fig. 6. Inhibition of MTX transport by mitoxantrone. MCF7/MX vesicles were incubated for 60 min in the presence of 400 μM [^3H]methotrexate and 0–100 μM cold mitoxantrone. The data represent the mean \pm SE (when applicable) of one to three individual assays performed with three separate vesicle preparations.




Fig. 7. Transport of MTX polyglutamates. The net accumulation of [^3H]methotrexate-Glu $_{1-4}$ into MCF7/MX vesicles was determined after 60-min exposure to 400 μM of the respective [^3H]methotrexate species in the absence (shaded columns) or presence (black columns) of 5 μM FTC. The data represent the mean \pm SE of three independent experiments performed with three separate vesicle preparations.

accumulation in the presence of mitoxantrone. MTX uptake into MCF7/MX vesicles was found to decrease with increasing concentrations of mitoxantrone (Fig. 6). The inhibition of MTX transport ranged from 12%, in the presence of 10 μM mitoxantrone, to nearly 60%, in the presence of 100 μM mitoxantrone. These data suggested competition between mitoxantrone and MTX and provided further evidence that MTX was indeed a substrate for BCRP.

BCRP-mediated Transport of Long-Chain Polyglutamates of MTX. MTX transport by several members of the MRP family has been shown to be largely restricted to MTX, with little or no transport observed when even a single extra glutamate was added (22, 23). To address whether BCRP-mediated transport of MTX was similarly limited, we measured accumulation of MTX and MTX-Glu $_2$ to MTX-Glu $_4$ into vesicles derived from MCF7/MX (Fig. 7) and MCF7/BCRP-Arg (data not shown) cells. As expected, drug uptake into vesicles derived from both cell lines decreased as the MTX polyglutamate chain length increased; however, in contrast to the MRPs, significant amounts of MTX-Glu $_2$ and MTX-Glu $_3$ were still transported. Furthermore, MTX polyglutamate transport was sensitive to the BCRP inhibitor FTC, indicating that BCRP, indeed, was transporting MTX as well as its polyglutamylated metabolites.

DISCUSSION

A substantial body of evidence exists to indicate the presence of one or more energy-dependent MTX efflux mechanism(s). For example, studies by Saxena and Henderson (18), using inside-out vesicles from L1210 cells, suggested the presence of two distinct, relatively low-efficiency ATP-dependent MTX efflux systems. In contrast, Schlemmer and Sirotiak (16) reported kinetic data for MTX transport in L1210 cells that suggested a single transporter with a somewhat higher affinity and capacity for MTX. However, neither group identified the actual protein involved. Other studies have shown that MTX is also a substrate for some MRPs. Low MTX uptake was observed into inside-out vesicles from MRP3-overexpressing LLC-PK1 cells (19), and from MRP1-overexpressing Sf9 (41) and GLC4/ADR (42) cells, whereas MTX was more efficiently transported by MRP2-containing Sf9 membrane vesicles (41). Furthermore, it was shown that rat *mrp2* effectively excreted L-MTX into the bile (43). Finally, Chen *et al.* and Zeng *et al.* (22, 23) demonstrated *in vitro* that MRP1–4 were all able to transport MTX, albeit with a low affinity, but with considerable capacity. Thus, there is clear evidence for ATP-dependent MTX transport by multiple ABC proteins. However, none of those described thus far seems to be able to also transport the polyglutamylated MTX metabolites.

In contrast, we present evidence that BCRP transports not only MTX but also MTX-polyglutamates. Using an *in vitro* inside-out vesicle system, we have demonstrated ATP-dependent transport of MTX into vesicles the membranes of which contain BCRP. MTX transport was osmotically sensitive and followed Michaelis-Menten kinetics. The K_m and V_{max} values of 680 μM and 2400 pmol/mg/min, respectively, suggest that BCRP is a low-affinity, high-capacity transporter. Furthermore, there was also substantial transport of polyglutamylated forms of MTX, MTX-Glu₂, and MTX-Glu₃. Accumulation of MTX-Glu₂ and MTX-Glu₃ was 60 and 40% of that observed with MTX. Finally, transport of all MTX species was sensitive to inhibition by the BCRP inhibitor FTC. Thus, the data presented clearly establish BCRP as a transporter of MTX and its polyglutamates.

Interestingly, MTX transport was only detected in vesicles with wild-type BCRP, which contains an arginine at amino acid position 482. In contrast, no uptake of MTX was observed into vesicles containing BCRP with an R482G mutation. This finding was consistent with the phenotype displayed by the cells from which the vesicles were derived, MCF7/MX (BCRP-Arg) and S1-M1–80 (BCRP-Gly), respectively. Whereas MCF7/MX cells are highly resistant to MTX, only minimal cross-resistance to MTX is observed in S1-M1–80 cells (28). Thus, it appears that the R482G mutation resulted in a dramatic change in substrate specificity for BCRP. In agreement with this conclusion, Honjo *et al.* (34) have demonstrated that cells expressing the arginine variant were resistant to mitoxantrone and CPT-11 but remained largely sensitive to Adriamycin. In contrast, cells expressing the glycine variant exhibited enhanced levels of mitoxantrone resistance as well as high levels of Adriamycin resistance, both of which were accompanied by significantly reduced drug accumulation. Furthermore, cellular rhodamine 123 accumulation was drastically reduced in cells with BCRP-Gly but unaffected in cells with BCRP-Arg (34). Similar effects were also observed with mouse *bcrp* when the corresponding arginine was mutated to either serine or methionine (35). Thus, it seems clear that the identity of the amino acid at position 482 has a profound effect on the specificity of BCRP-mediated drug transport. However, additional studies are needed to elucidate the exact mechanism behind this observation.

Several members of the MRP family of ABC transporters have been shown to transport MTX, and their kinetic parameters have been

determined. Reported K_m values for MRP 1, 3, and 4 are 2150, 620, and 220 μM , respectively, and V_{max} values are 2050, 2930, and 240 pmol/mg/min, respectively (22, 23). These values are comparable with those determined for BCRP in the present study. However, in contrast to what was observed for BCRP, the addition of even a single extra glutamate to MTX resulted in essentially complete abrogation of transport by the MRPs. Thus, although the affinities and capacities are in the same range, the ability to transport MTX-Glu₂ and MTX-Glu₃ makes BCRP clearly distinct from the MRPs, and unique among the ABC proteins reported to transport MTX.

It is tempting to speculate that the ability of BCRP to transport polyglutamylated MTX is responsible for the distinct MTX-resistance pattern of BCRP-overexpressing cells. Cells that overexpress BCRP are resistant against MTX in continuous exposure assays when polyglutamylation is likely to occur, suggesting that transport of polyglutamylated MTX may be an important requirement for resistance. In contrast, cells that overexpress MRPs are only resistant to short-term (<3 h) MTX exposure, when it is unlikely that MTX has undergone polyglutamylation (20, 21, 23). Thus, despite the low affinity of BCRP for MTX, its capacity to also transport MTX-polyglutamates may account for the high levels of resistance in long-term assays. It further suggests that MTX-polyglutamates can be exported even as they are formed, as opposed to only after degradation, which is consistent with the observed lower steady-state levels of MTX-polyglutamates in MCF7/MX cells (24).

The physiological significance of a K_m as high as that observed for BCRP is unclear because it is doubtful whether the intracellular free MTX concentration will ever reach levels of that order of magnitude *in vivo*. For example, total MTX concentrations in the order of 500–3500 pmol/10⁹ cells or 2–15 μM , have been found in pediatric leukemia cells (44–46), which is two to three orders of magnitude below the K_m for BCRP as well as for the MRPs. However, plasma MTX levels of 100 μM and greater have been measured 42 h after an initial high-dose MTX infusion (44). Furthermore, it is conceivable that the local concentration on the cytoplasmic side of the plasma membrane may be higher than the average intracellular concentration. Nevertheless, despite the low affinity, when the overexpression of BCRP is sufficiently high, it can apparently cause enhanced MTX efflux and concomitant resistance (28).

BCRP was originally identified and described as a mitoxantrone transporter. Therefore, we determined the effect of mitoxantrone on MTX transport. These studies have demonstrated that the presence of mitoxantrone reduced the transport of MTX. Increasing concentrations of mitoxantrone resulted in decreasing MTX accumulation into MCF7/MX vesicles, indicating that the presence of mitoxantrone interferes and possibly competes with MTX transport. These results further support the idea that BCRP is a MTX transporter. However, the data do not allow for further characterization of the type of competition between the two substrates.

Previously, MTX resistance in the MCF7/MX cells had been attributed to an ATP-dependent efflux mechanism, and it was shown that resistance was reversible with BCRP inhibitors (24, 28). Furthermore, MTX cross-resistance was not unique to the MCF7/MX cells; multiple BCRP-overexpressing cell lines exhibited similar cross-resistance, suggesting that it was caused by BCRP. Here, we extended this association and directly demonstrated that BCRP functions as a MTX efflux pump. BCRP appears to be unique among the MTX exporters described thus far in that it is also able to transport polyglutamylated forms of MTX. The efflux of MTX-polyglutamates may be a major contributor to MTX resistance under long-term exposure conditions.

ACKNOWLEDGMENTS

We gratefully acknowledge Jolene Johnson for her technical support and Dr. Susan Bates for FTC.

Note Added in Proof

After this manuscript was accepted for publication, two reports appeared that independently came to essentially the same conclusions. (1) Z. S. Chen, *et al.* Transport of Methotrexate, Methotrexate Polyglutamates, and 17 β -Estradiol 17-(β -D-glucuronide) by ABCG2: Effects of Acquired Mutations at R482 on Methotrexate Transport. *Cancer Res.*, 63: 4048–4054, 2003. (2) H. Mitomo, *et al.* A functional study on polymorphism of the ATP-binding cassette transporter ABCG2: critical role of arginine-482 in methotrexate transport. *Biochem J.*, 373: 767–774, 2003.

REFERENCES

- Alt, F. W., Kellems, R. E., Bertino, J. R., and Schimke, R. T. Selective multiplication of dihydrofolate reductase genes in methotrexate resistant variants of cultured murine cells. *J. Biol. Chem.*, 253: 1357–1370, 1978.
- Dicker, A. P., Volkenandt, M., Schweitzer, B. I., Banerjee, D., and Bertino, J. R. Identification and characterization of a mutation in the dihydrofolate reductase gene from the methotrexate-resistant Chinese hamster ovary cell line Pro-3 MtxRIII. *J. Biol. Chem.*, 265: 8317–8321, 1990.
- Yu, M., and Melera, P. Allelic variation in the dihydrofolate reductase gene at amino acid position 95 contributes to antifolate resistance in Chinese hamster cells. *Cancer Res.*, 53: 6031–6035, 1993.
- Srimatkandada, S., Schweitzer, B. I., Moroson, B. A., Dube, S., and Bertino, J. R. Amplification of a polymorphic dihydrofolate reductase gene expressing an enzyme with decreased binding to methotrexate in a human colon carcinoma cell line, HCT-8R4, resistant to this drug. *J. Biol. Chem.*, 264: 3524–3528, 1989.
- Dixon, K. H., Lanpher, B. C., Chiu, J., Kelley, K., and Cowan, K. H. A novel cDNA restores reduced folate carrier activity and methotrexate sensitivity to transport deficient cells. *J. Biol. Chem.*, 269: 17–20, 1994.
- Jansen, G., Mauritz, R., Drori, S., Sprecher, H., Kathmann, I., Bunni, M., Priest, D. G., Noordhuis, P., Schornagel, J. H., Pinedo, H. M., Peters, G. J., and Assaraf, Y. G. A structurally altered human reduced folate carrier with increased folic acid transport mediates a novel mechanism of antifolate resistance. *J. Biol. Chem.*, 273: 30189–30198, 1998.
- Roy, K., Tolner, B., Chiao, J., and Sirotnak, F. A single amino acid difference within the folate transporter encoded by the murine *RFC-1* gene selectively alters its interaction with folate analogues. *J. Biol. Chem.*, 273: 2526–2531, 1998.
- Zhao, R., Sharina, I. G., and Goldman, I. D. Pattern of mutations that results in loss of reduced folate carrier function under antifolate selective pressure augmented by chemical mutagenesis. *Mol. Pharmacol.*, 56: 68–76, 1999.
- Roy, K., Egan, M., Sirlin, S., and Sirotnak, F. Posttranscriptionally mediated decreases in folypolyglutamate synthetase gene expression in some folate analogue-resistant variants of the L1210 cell. *J. Biol. Chem.*, 272: 6903–6908, 1997.
- Rhee, M. S., Wang, Y., Nair, M. G., and Galivan, J. Acquisition of resistance to antifolates caused by enhanced γ -glutamyl hydrolase activity. *Cancer Res.*, 53: 2227–2230, 1993.
- Hakala, M. T. On the nature of permeability of sarcoma-180 cells to amethopterin *in vitro*. *Biochim. Biophys. Acta*, 102: 210–225, 1965.
- Goldman, I. D. Transport energetics of the folic acid analogue, methotrexate, in L1210 leukemia cells. Enhanced accumulation by metabolic inhibitors. *J. Biol. Chem.*, 244: 3779–3785, 1969.
- Barakat, R. R., Li, W. W., Lovelace, C., and Bertino, J. R. Intrinsic resistance of cervical squamous cell carcinoma cell lines to methotrexate (MTX) as a result of decreased accumulation of intracellular MTX polyglutamates. *Gynecol. Oncol.*, 51: 54–60, 1993.
- Dembo, M., Sirotnak, F. M., and Moccio, D. M. Effects of metabolic deprivation on methotrexate transport in L1210 leukemia cells: further evidence for separate influx and efflux systems with different energetic requirements. *J. Membr. Biol.*, 78: 9–17, 1984.
- Sirotnak, F., and O'Leary, D. The issues of transport multiplicity and energetics pertaining to methotrexate efflux in L1210 cells addressed by an analysis of *cis*- and *trans*- effects of inhibitors. *Cancer Res.*, 51: 1412–1417, 1991.
- Schlemmer, S., and Sirotnak, F. Energy-dependent efflux of methotrexate in L1210 leukemia cells. *J. Biol. Chem.*, 267: 14746–14752, 1992.
- Schlemmer, S., and Sirotnak, F. Retentiveness of methotrexate polyglutamates in cultured L1210 cells. *Biochem. Pharmacol.*, 45: 1261–1266, 1993.
- Saxena, M., and Henderson, G. B. Identification of efflux systems for large anionic conjugates as the mediators of methotrexate efflux in L1210 cells. *Biochem. Pharmacol.*, 51: 975–982, 1996.
- Hirohashi, T., Suzuki, H., and Sugiyama, Y. Characterization of the transport properties of cloned rat multidrug Resistance-associated protein 3 (MRP3). *J. Biol. Chem.*, 274: 15181–15185, 1999.
- Hooijberg, J. H., Broxterman, H. J., Kool, M., Assaraf, Y. G., Peters, G., Noordhuis, P., Scheper, R. J., Borst, P., Pinedo, H. M., and Jansen, G. Antifolate resistance mediated by the multidrug resistance proteins MRP1 and MRP2. *Cancer Res.*, 59: 2532–2535, 1999.
- Lee, K., Klein-Szanto, A. J., and Kruh, G. D. Analysis of the MRP4 drug resistance profile in transfected NIH3T3 cells. *J. Natl. Cancer Inst.* (Bethesda), 92: 1934–1940, 2000.
- Chen, Z. S., Lee, K., Walther, S., Blanchard Raftogianis, R., Kuwano, M., Zeng, H., and Kruh, G. D. Analysis of methotrexate and folate transport by multidrug resistance protein 4 (ABCC4): MRP4 is a component of the methotrexate efflux system. *Cancer Res.*, 62: 3144–3150, 2002.
- Zeng, H., Chen, Z.-S., Belinsky, M. G., Rea, P. A., and Kruh, G. D. Transport of methotrexate and folates by multidrug resistance protein (MRP) 3 and MRP1: effect of polyglutamylamoylation on MTX transport. *Cancer Res.*, 61: 7225–7232, 2001.
- Volk, E. L., Rohde, K., Rhee, M., McGuire, J. J., Doyle, L. A., Ross, D. D., and Schneider, E. Methotrexate cross-resistance in a mitoxantrone-selected multidrug resistant MCF7 breast cancer cell line is attributable to enhanced energy-dependent drug efflux. *Cancer Res.*, 60: 3514–3521, 2000.
- Miyake, K., Mickley, L., Litman, T., Zhan, Z., Robey, R., Cristensen, B., Brangi, M., Greenberger, L., Dean, M., Fojo, T., and Bates, S. E. Molecular cloning of cDNAs which are highly overexpressed in mitoxantrone-resistant cells: demonstration of homology to ABC transport genes. *Cancer Res.*, 59: 8–13, 1999.
- Allikmets, R., Schriml, L., Hutchinson, A., Romano-Spica, V., and Dean, M. A human placenta-specific ATP-binding cassette gene (*ABCP*) on chromosome 4q22 that is involved in multidrug resistance. *Cancer Res.*, 58: 5337–5339, 1998.
- Klein, I., Sarkadi, B., and Varadi, A. An inventory of the human ABC proteins. *Biochim. Biophys. Acta*, 1461: 237–262, 1999.
- Volk, E. L., Farley, K. M., Wu, Y., Li, F., Robey, R. W., and Schneider, E. Overexpression of wild-type breast cancer resistance protein mediates methotrexate resistance. *Cancer Res.*, 62: 5035–5040, 2002.
- Soule, H. D., Vazquez, J., Long, A., Albert, S., and Brennan, M. A human cell line from a pleural effusion derived from a breast carcinoma. *J. Natl. Cancer Inst.* (Bethesda), 51: 1409–1413, 1973.
- Nakagawa, M., Schneider, E., Dixon, K. H., Horton, J., Kelley, K., Morrow, C., and Cowan, K. H. Reduced intracellular drug accumulation in the absence of P-glycoprotein (*mdr1*) overexpression in mitoxantrone-resistant MCF-7 breast cancer cells. *Cancer Res.*, 52: 6175–6181, 1992.
- Loe, D. W., Deeley, R. G., and Cole, S. P. Characterization of vincristine transport by the M_r 190,000 multidrug resistance protein (MRP): evidence for cotransport with reduced glutathione. *Cancer Res.*, 58: 5130–5136, 1998.
- Steck, T. L., and Kant, J. A. Preparation of impermeable ghosts and inside-out vesicles from human erythrocyte membranes. *Methods Enzymol.*, 31: 172–180, 1974.
- Saxena, M., and Henderson, G. B. MOAT4, a novel multispecific organic-anion transporter for glucuronides and mercapturates in mouse L1210 cells and human erythrocytes. *Biochem. J.*, 320: 273–281, 1996.
- Honjo, Y., Hrycyna, C. A., Yan, Q. W., Medina-Perez, W. Y., Robey, R. W., van de Laar, A., Litman, T., Dean, M., and Bates, S. E. Acquired mutations in the *MXR/BCRP/ABCP* gene alter substrate specificity in *MXR/BCRP/ABCP*-overexpressing cells. *Cancer Res.*, 61: 6635–6639, 2001.
- Allen, J. D., Jackson, S. C., and Schinkel, A. H. A mutation hot spot in the *Bcrp1* (*Abcg2*) multidrug transporter in mouse cell lines selected for doxorubicin resistance. *Cancer Res.*, 62: 2294–2299, 2002.
- Rabindran, S. K., He, H., Singh, M., Brown, E., Collins, K. I., Annable, T., and Greenberger, L. M. Reversal of a novel multidrug resistance mechanism in human colon carcinoma cells by fumitremorgin C. *Cancer Res.*, 58: 5850–5858, 1998.
- Rabindran, S. K., Ross, D. D., Doyle, L. A., Yang, W., and Greenberger, L. M. Fumitremorgin C reverses multidrug resistance in cells transfected with the breast cancer resistance protein. *Cancer Res.*, 60: 47–50, 2000.
- Robey, R. W., Medina-Perez, W. Y., Nishiyama, K., Lahusen, T., Miyake, K., Litman, T., Senderowicz, A. M., Ross, D. D., and Bates, S. E. Overexpression of the ATP-binding cassette half-transporter, ABCG2 (*MXR/BCRP/ABCP1*), in flavopiridol-resistant human breast cancer cells. *Clin. Cancer Res.*, 7: 145–152, 2001.
- Ross, D. D., Yang, W., Abruzzo, L. V., Dalton, W. S., Schneider, E., Lage, H., Dietel, M., Greenberger, L., Cole, S. P. C., and Doyle, L. A. Atypical multidrug resistance: breast cancer resistance protein messenger RNA expression in mitoxantrone-selected cell lines. *J. Natl. Cancer Inst.* (Bethesda), 91: 429–433, 1999.
- Doyle, L. A., Yang, W., Abruzzo, L. V., Krogmann, T., Gao, Y., Rishi, A. K., and Ross, D. D. A multidrug resistance transporter from human MCF-7 breast cancer cells. *Proc. Natl. Acad. Sci. USA*, 95: 15665–15670, 1998.
- Bakos, E., Evers, R., Sinko, E., Varadi, A., Borst, P., and Sarkadi, B. Interactions of the human multidrug resistance proteins MRP1 and MRP2 with organic anions. *Mol. Pharmacol.*, 57: 760–768, 2000.
- Heijn, M., Hooijberg, J. H., Scheffer, G. L., Szabo, G., Westerhoff, H. V., and Lankelma, J. Anthracyclines modulate multidrug resistance protein (MRP) mediated organic anion transport. *Biochim. Biophys. Acta*, 1326: 12–22, 1997.
- Masuda, M., Iizuka, Y., Yamazaki, M., Nishigaki, R., Kato, Y., Ni'inuma, K., Suzuki, H., and Sugiyama, Y. Methotrexate is excreted into the bile by canalicular multispecific organic anion transporter in rats. *Cancer Res.*, 57: 3506–3510, 1997.
- Masson, E., Relling, M. V., Synold, T. W., Liu, Q., Schuetz, J. D., Sandlund, J. T., Pui, C.-H., and Evans, W. E. Accumulation of methotrexate polyglutamates in lymphoblasts is a determinant of antileukemic effects *in vivo*. *J. Clin. Invest.*, 97: 73–80, 1996.
- Whitehead, V., Payment, C., Cooley, L., Lauer, S., Mahoney, D., Shuster, J., Vuchich, M.-J., Bernstein, M., Look, A., Pullen, D., and Camitta, B. The association of the TEL-AML1 chromosomal translocation with the accumulation of methotrexate polyglutamates in lymphoblasts and with ploidy in childhood B-progenitor cell acute lymphoblastic leukemia: a Pediatric Oncology Group study. *Leukemia* (Baltimore), 15: 1081–1088, 2001.
- Panetta, J., Yanishevski, Y., Pui, C., Sandlund, J., Rubnitz, J., Rivera, G., Ribiero, R., Evans, W., and Relling, M. A mathematical model of *in vivo* methotrexate accumulation in acute lymphoblastic leukemia. *Cancer Chemother. Pharm.* 50: 419–428, 2002.

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

Wild-Type Breast Cancer Resistance Protein (BCRP/ABCG2) is a Methotrexate Polyglutamate Transporter

Erin L. Volk and Erasmus Schneider

Cancer Res 2003;63:5538-5543.

Updated version Access the most recent version of this article at:
<http://cancerres.aacrjournals.org/content/63/17/5538>

Cited articles This article cites 44 articles, 31 of which you can access for free at:
<http://cancerres.aacrjournals.org/content/63/17/5538.full#ref-list-1>

Citing articles This article has been cited by 28 HighWire-hosted articles. Access the articles at:
<http://cancerres.aacrjournals.org/content/63/17/5538.full#related-urls>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cancerres.aacrjournals.org/content/63/17/5538>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.