

Loss of *Phosphatase and Tensin Homologue Deleted on Chromosome 10* Engages ErbB3 and Insulin-Like Growth Factor-I Receptor Signaling to Promote Antiestrogen Resistance in Breast Cancer

Todd W. Miller,¹ Marianela Pérez-Torres,² Archana Narasanna,¹ Marta Guix,¹ Olle Stål,⁵ Gizeh Pérez-Tenorio,⁵ Ana M. Gonzalez-Angulo,^{6,7} Bryan T. Hennessy,^{7,8} Gordon B. Mills,⁷ J. Phillip Kennedy,³ Craig W. Lindsley,³ and Carlos L. Arteaga^{1,2,4}

Departments of ¹Medicine, ²Cancer Biology, and ³Chemistry, and ⁴Breast Cancer Research Program, Vanderbilt-Ingram Comprehensive Cancer Center, Vanderbilt University, Nashville, Tennessee; ⁵Department of Biomedicine and Surgery, Division of Oncology, Faculty of Health Sciences, Linköping University, Linköping, Sweden; Departments of ⁶Breast Medical Oncology, ⁷Systems Biology, and ⁸Gynecology Medical Oncology, University of Texas, M. D. Anderson Cancer Center, Houston, Texas

Abstract

Knockdown of the tumor suppressor phosphatase *Phosphatase and tensin homologue deleted on chromosome 10* (*PTEN*) with shRNA in three estrogen receptor (ER)-positive breast cancer cell lines resulted in increased phosphatidylinositol-3 kinase (PI3K) and AKT activities, resistance to tamoxifen and fulvestrant, and hormone-independent growth. *PTEN* knockdown induced the up-regulation of ER transcriptional activity in MCF-7 cells but decreased ER protein levels and transcriptional activity in T47D and MDA-361 cells. Tamoxifen and fulvestrant treatment inhibited estradiol-induced ER transcriptional activity in all shPTEN cell lines but did not abrogate the increased cell proliferation induced by *PTEN* knockdown. *PTEN* knockdown increased basal and ligand-induced activation of the insulin-like growth factor-I (IGF-I) and ErbB3 receptor tyrosine kinases, and prolonged the association of the p85 PI3K subunit with the IGF-I receptor (IGF-IR) effector insulin receptor substrate-1 and with ErbB3, implicating *PTEN* in the modulation of signaling upstream of PI3K. Consistent with these data, *PTEN* levels inversely correlated with levels of tyrosine-phosphorylated IGF-IR in tissue lysate arrays of primary breast cancers. Inhibition of IGF-IR and/or ErbB2-mediated activation of ErbB3 with tyrosine kinase inhibitors restored hormone dependence and the growth inhibitory effect of tamoxifen and fulvestrant on shPTEN cells, suggesting that cotargeting both ER and receptor tyrosine kinase pathways holds promise for the treatment of patients with ER+, *PTEN*-deficient breast cancers. [Cancer Res 2009;69(10):4192–201]

Introduction

Loss-of-function mutations of the *Phosphatase and tensin homologue deleted on chromosome 10* (*PTEN*) tumor suppressor gene occur in 5% to 45% of human cancers (1), with reduced *PTEN* protein found in 31% to 48% of breast cancers (2–4). The main tumor suppressive action of *PTEN* is its lipid phosphatase activity

to antagonize phosphatidylinositol-3 kinase (PI3K) by dephosphorylating its product, phosphatidylinositol (3,4,5)-trisphosphate (PIP₃), resulting in inhibition of the serine/threonine kinase AKT and other pleckstrin homology domain-containing proteins that modulate cell growth, survival, and angiogenesis. *PTEN* can also act as a protein phosphatase with targets including focal adhesion kinase (5), platelet-derived growth factor receptor, epidermal growth factor receptor (EGFR; ref. 6), and itself (7), and as a binding partner to increase p53 activity (8).

Two-thirds of breast cancers express estrogen receptor (ER)- α , which drives breast cancer cell growth. Although endocrine therapies designed to block estrogen action (e.g., tamoxifen, aromatase inhibitors) have changed the natural history of hormone-dependent breast cancer, many tumors exhibit *de novo* or acquired therapeutic resistance. Crosstalk between receptor tyrosine kinase (RTK) and ER signaling promotes resistance to endocrine therapy (9). Tumor overexpression of RTKs and RTK ligands, and increased RTK pathway activation, have been linked to antiestrogen resistance (10–12). For example, the RTK effectors AKT and mitogen-activated protein kinase (MAPK) can phosphorylate ER (13, 14), and MAPK can phosphorylate the ER coactivator AIB1, to promote ER transcriptional activity (15). In turn, ER drives transcription of *insulin-like growth factor-I* (*IGF-I*), *IGF-II*, *IGF-I receptor* (*IGF-IR*), and its effector *insulin receptor substrate-1* (*IRS-1*). Activated *IRS-1* induces PI3K activation (16) and is stabilized by ER (17). Furthermore, estrogen induces the expression of genes encoding the EGFR ligands transforming growth factor α and amphiregulin (18, 19) and can activate EGFR, HER2/ErbB2, and downstream signal transducers by transcription-independent mechanisms (20). Although the traditional role of ER as a transcription factor is central to ER+ breast cancer, ER has also been implicated in nongenomic, plasma membrane-initiated signaling with IGF-IR, EGFR, and PI3K (21, 22).

Mutational activation of the PI3K pathway, by *PTEN* loss and/or gain-of-function mutations in *PIK3CA* (which encodes the PI3K p110 α catalytic subunit), occurs in 56% to 62% of ER+ breast cancers (23, 24). Patients with cancers exhibiting a gene expression signature of *PTEN* loss show poor disease outcome (24). Although PI3K mutations and *PTEN* loss are both thought to confer increased PI3K activity, the cellular effects of these mutations may be different, as suggested by the coexistence of these alterations in 5% to 14% of primary breast tumors (2, 23, 24). We therefore investigated the effects of *PTEN* loss in three ER+ human

Note: Supplementary data for this article are available at Cancer Research Online (<http://cancerres.aacrjournals.org/>).

Requests for reprints: Carlos L. Arteaga, Vanderbilt University Medical Center, 2200 Pierce Avenue, 777 PRB, Nashville, TN 37232-6307. Phone: 615-936-3524; Fax: 615-936-1790; E-mail: carlos.arteaga@vanderbilt.edu.

©2009 American Association for Cancer Research.
doi:10.1158/0008-5472.CAN-09-0042

breast cancer cell lines on PI3K activation, hormone-independent growth, and response to antiestrogens.

Materials and Methods

Cell lines. MCF-7, T47D, and MDA-361 cells (American Type Culture Collection) were stably transduced with retrovirus-encoding shRNA targeting *PTEN* or mismatch control (shMM; as in Supplementary Materials and Methods). Experiments were performed using phenol red-free IMEM + dextran-charcoal-treated-fetal bovine serum (DCC-FBS; Hyclone) unless otherwise indicated.

Phospholipid analysis. MCF-7 lines were labeled $\times 16$ h with 100 $\mu\text{Ci}/\text{mL}$ [^{32}P]- P_i (Perkin-Elmer) in phosphate-free DMEM + 10% dialyzed FBS (Hyclone). Radiolabeled lipids were extracted, concentrated, and separated by TLC as described (25). ^{32}P incorporation into phosphatidylinositol species was detected by autoradiography.

Cell proliferations assays. Cells were seeded in triplicate in 12-well plates (2.5×10^4 per well). The next day, medium was changed to IMEM + DCC-FBS \pm 17- β -estradiol (E2), 4-hydroxy-tamoxifen (4-OH-T), fulvestrant (faslodex; ICI182780; gift from AstraZeneca), testosterone, letrozole (gift from Dean Evans, Novartis, Basel, Switzerland), the allosteric AKT1/2 inhibitor 0360263-1 (AKTi; ref. 26), BEZ235 (27), AEW541 (both provided by Carlos Garcia-Echeverria, Novartis, Basel, Switzerland; ref. 28), or lapatinib ditosylate (GW-572016; LC Laboratories). For siRNA of ErbB3/HER3, cells were transfected as in Supplementary Materials and Methods. Media were refreshed every 2 to 3 d, and after 5 to 8 d, cells were trypsinized and counted using a Coulter counter.

ER transcriptional reporter assays. Cells were plated as above and transfected with pGLB-MERE (provided by Dorraya El-Ashry, University of Miami, Miami, FL) and pCMV-Renilla (Promega) luciferase plasmids. Cells were then treated as above, and luciferase activity was measured 16 to 20 h later as described (29).

Immunoprecipitation and immunoblotting. Cells were treated as indicated (IGF-I and heregulin- β 1; R&D Systems). Cells were lysed in NP40 buffer plus protease and phosphatase inhibitors, sonicated for 10 s,

centrifuged at 14k rpm for 10 min., and protein was quantitated using BCA assay (Pierce). Immunoprecipitations were performed using Dynal protein-G beads (Invitrogen) and p85 antibody (Upstate) as described (30). Immunoprecipitates and cell lysates were subjected to SDS-PAGE and transferred to nitrocellulose. Primary antibodies for immunoblotting included PTEN, IGF-IR β , HER3, ER α (Santa Cruz Biotechnology), AKT, P-AKT_{S473}, P-HER2_{Y1248}, P-HER3_{Y1289}, P-IGF-IR β _{Y1131} (Cell Signaling), PR (Dako), P-tyrosine (4G10, Vanderbilt Monoclonal Antibody Core), IRS-1 (Upstate), and actin (Sigma). Phospho-RTK arrays were performed as per manufacturer's protocol (R&D Systems).


Reverse phase protein lysate microarray analysis. Three hundred and eighty-three hormone receptor-positive, primary breast tumor samples were obtained from the Breast Tissue Frozen Tumor Bank at M. D. Anderson Cancer Center. Specimens were collected under Institutional Review Board-approved protocols. Tumor lysates were analyzed as in Supplementary Materials and Methods (31) using antibodies against PTEN, P-IGF-IR β _{Y1135/1136} (may cross-react with P-InsR β _{Y1150/1151}), and IGF-IR β . Relative protein levels were quantified, and PTEN scores were plotted against P-IGF-IR β /IGF-IR β ratio.

Statistical analysis. In cell proliferation assays and ER transcriptional reporter assays, significant differences were determined by two-tailed *t* test. In reverse phase protein lysate microarray analysis, relative units for PTEN, P-IGF-IR β , and IGF-IR β levels were converted to logarithms, and the relationship between PTEN and ratio of P-IGF-IR β /IGF-IR β was analyzed using two-tailed *t* test and Pearson correlation. *P* value of <0.05 was considered significant.

Results

PTEN loss results in hormone-independent growth and antiestrogen resistance. We stably knocked down *PTEN* in MCF-7, T47D, and MDA-361 ER+ breast cancer cells using shMM. In shPTEN cell lines, PTEN protein was undetectable and AKT phosphorylation at Ser₄₇₃ (P-AKT) was up-regulated compared with shMM controls (Fig. 1A). All three of these cell lines harbor

Figure 1. PTEN loss increases PIP₃ and P-AKT, and alters ER and PR levels. **A**, cells were treated with medium containing DCC-FBS (MCF-7, 2%; T47D and MDA-361, 0.5%) $\times 24$ h, and lysates were analyzed by immunoblotting with the indicated antibodies. PR was not detected in MDA-361 cells. **B**, MCF-7 lines were metabolically labeled with ^{32}P -orthophosphate in 10% dialyzed FBS $\times 16$ h. Lipids were extracted, resolved by TLC, and ^{32}P -PIP species were detected by autoradiography (arrows). Origin of spotting is indicated. Cell lysates were also used for immunoblotting to confirm PTEN status. Fold changes in PR isoforms (A) and PIP₃ (B) normalized to actin were determined by densitometry analysis (bar graphs).







Figure 2. PTEN loss increases hormone-independent growth and antiestrogen resistance. **A**, cell proliferation assays. MCF-7 (*i* and *iv*), T47D (*ii*), and MDA-361 (*iii*) lines were treated with medium containing DCC-FBS (MCF-7, 2%; T47D and MDA-361, 0.5%) with the indicated compounds (E2; 4-OH-T; Fulvest, Fulvestrant; Lz, letrozole). Media and drugs were refreshed every 2 to 3 d. Adherent cells were counted after 5 to 8 d. **B**, ER transcriptional reporter assays. MCF-7 (*i* and *iv*), T47D (*ii*), and MDA-361 (*iii*) lines were transfected with luciferase reporter plasmids. Cells were treated as in (**A**), and luciferase activities were measured after 16 to 20 h. *RLU*, relative light units (firefly/*Renilla*). All data are presented as % untreated shMM control; columns, mean of triplicates; bars, SD. *, $P < 0.05$ by *t* test comparing shPTEN to shMM under each condition.

activating mutations in *PIK3CA*, which encodes the PI3K p110 α catalytic subunit (2). In addition, PTEN loss increased levels of the PI3K product PIP₃ (Fig. 1B). ER levels were unchanged by PTEN loss in MCF-7 cells but were decreased in T47D and MDA-361 cells. PTEN loss reduced progesterone receptor (PR) levels in MCF-7 and

T47D cells, which is consistent with the negative correlation between PTEN and PR levels observed in human breast cancers (2).

In cell proliferation assays, shPTEN cells significantly outgrew shMM controls under hormone-depleted and E2-induced conditions ($P < 0.05$; Supplementary Fig. S1; Fig. 2A). In 5 of 5 nude mice,

Figure 3. PTEN loss activates RTKs upstream of PI3K. **A**, lysates from MCF-7 (*i*), T47D (*ii*), and MDA-361 (*iii*) lines treated with medium containing DCC-FBS (MCF-7, 2%; T47D and MDA-361, 0.5%) overnight were used to probe phospho-RTK arrays. Antibodies against 42 RTKs are spotted in duplicate on a membrane. Membranes are incubated with cell lysates followed by probing with a P-tyrosine antibody. A positive signal is indicative of receptor phosphorylation. Blots from each pair of lines are exposure-matched. Tyrosine-phosphorylated RTKs are labeled as follows: 1, EGFR; 2, HER2; 3, HER3; 4, InsR β ; 5, IGF-IR β ; 6, ErbB4; 7, ROR2; 8, EphA1. Positive controls are spotted at corners. **B**, lysates from MCF-7 (*i*), T47D (*ii*), and MDA-361 (*iii*) lines treated as in **A** overnight \pm the indicated kinase inhibitors were used for immunoblotting with the indicated antibodies. Short and long exposures (*exp*) are shown in *iii*. **C**, tumor lysates from 383 hormone receptor-positive breast cancers were analyzed by reverse phase protein lysate microarray analysis to quantify relative levels of PTEN, P-IGF-IR β ^{Y1135/1136}, and IGF-IR β . Shown is a scatterplot of PTEN versus ratio of P-IGF-IR β /IGF-IR β (indicative of the fraction of activated IGF-IR β). This relationship was analyzed using two-tailed *t* test and Pearson correlation.



MCF-7/shPTEN cells formed palpable tumors (≥ 3 mm in diameter) versus only 1 of 5 animals injected with MCF-7/shMM cells (Supplementary Fig. S1). Although all lines were sensitive to the inhibitory effects of the selective ER modulator 4-OH-T and the ER down-regulator fulvestrant, shPTEN cells exhibited significantly increased growth compared with shMM controls ($P < 0.05$; Fig. 2A), indicative of relative antiestrogen resistance. The hormone-independent proliferation of MDA-361/shMM cells, but not MDA-361/shPTEN cells, was inhibited by 4-OH-T and fulvestrant. Furthermore, testosterone induced the growth of MCF-7/shPTEN but not MCF-7/shMM cells ($P < 0.05$; Fig. 2A, *iv*). This response was blocked by the aromatase inhibitor letrozole, suggesting that *PTEN* knockdown up-regulated the cellular response to E2 produced by the aromatization of testosterone. Aromatase mRNA levels were unchanged by *PTEN* knockdown (determined by microarray analysis; data not shown). Aromatase protein was undetectable by immuno-

blot analysis of MCF-7 lines, but low levels of aromatase are expressed in MCF-7 cells (32).

PI3K/AKT pathway activation has been associated with increased ER transcriptional activity and reduced ER expression in MCF-7 cells (14, 33). Here, MCF-7/shPTEN cells showed increased basal, E2-induced, and testosterone-induced ER transcriptional reporter activity compared with shMM cells (Supplementary Fig. S2; Fig. 2B, *i* and *iv*). However, *PTEN* loss decreased basal and E2-stimulated ER transcriptional activity in T47D and MDA-361 cells (Supplementary Fig. S2; Fig. 2B, *ii* and *iii*), reflective of the reduced ER levels in these cells (Fig. 1). We also observed variable effects of *PTEN* loss on the expression of the E2-inducible genes *AREG* and *EGR3* (18). In MCF-7 and T47D cells, *PTEN* knockdown increased the basal and E2-induced mRNA levels of *AREG* and *EGR3* compared with shMM controls (Supplementary Fig. S3). Opposite effects were observed in MDA-361 cells.

Treatment with 4-OH-T or fulvestrant did not consistently suppress the effects of *PTEN* knockdown on *AREG* or *EGR3* expression, suggesting that *PTEN* loss alters their expression by both ER-dependent and ER-independent mechanisms. Although treatment with 4-OH-T, fulvestrant, or letrozole did not abrogate the growth advantage of sh*PTEN* cell lines (Fig. 2A), these inhibitors markedly suppressed E2- and testosterone-induced ER transcriptional activity in all lines (Supplementary Fig. S4; Fig. 2B). Overall, these data suggest that the increased hormone-independent growth and relative resistance to 4-OH-T and fulvestrant conferred by *PTEN* loss cannot be solely explained by increased ER transcriptional activity.

PTEN loss increases IGF-IR- and ErbB3-mediated activation of PI3K. We next examined mechanisms that can activate PI3K in *PTEN*-deficient cells. Using phospho-RTK array analysis, we detected predominant tyrosine phosphorylation of IGF-IR β , ErbB3/HER3, insulin receptor- β (InsR β), and EGFR in MCF-7 cells (Fig. 3A, *i*), and of EGFR, HER2, and HER3 in T47D and MDA-361 cells (Fig. 3A, *ii* and *iii*). *PTEN* loss up-regulated tyrosine phosphorylation of EGFR, InsR β , and IGF-IR β in MCF-7 cells, and of EGFR, HER3, ErbB4/HER4, and ROR2 in T47D cells. We next used a loss-of-function approach to assess which RTKs activated PI3K, measuring P-AKT_{S473} as a surrogate of PI3K activity. The IGF-IR tyrosine kinase inhibitor AEW541 reduced P-AKT in MCF-7 cells (Fig. 3B, *i*), whereas the EGFR/HER2 tyrosine kinase inhibitor lapatinib reduced P-AKT in T47D and MDA-361 cells (Fig. 3B, *ii* and *iii*). Treatment with lapatinib decreased P-HER3 in all three cell

lines (Supplementary Fig. S4), suggesting that HER3 activation is EGFR/HER2-dependent in these cells, and that HER3 is the predominant activator of PI3K in T47D and MDA-361 cells. In all sh*PTEN* lines, the combination of AEW541 plus lapatinib inhibited P-AKT more effectively than either drug alone. AKTi and the PI3K/mammalian target of rapamycin inhibitor BEZ235 decreased P-AKT in all lines.

To ascertain whether *PTEN* loss is linked with IGF-IR activation in primary breast cancers, we analyzed 383 hormone receptor-positive tumors by reverse phase protein lysate microarray for levels of *PTEN*, tyrosine-phosphorylated IGF-IR β (P-IGF-IR β), and total IGF-IR β . We observed a statistically significant inverse correlation between *PTEN* levels and P-IGF-IR β /IGF-IR β ratio ($R^2 = 0.182$; $P < 10^{-15}$; Fig. 3C), consistent with the results observed upon *PTEN* knockdown in MCF-7 cells (Fig. 3A). Because P-IGF-IR β antibodies may cross-react with P-InsR β , we cannot rule out that P-InsR β may also contribute to this correlation.

Growth factor receptors activate PI3K by phosphorylating adaptor proteins such as GAB1, GAB2, IRS-1, IRS-2, and HER3. Tyrosine-phosphorylated adaptors engage the N-SH2 domain of the PI3K regulatory subunit p85, relieving the inhibition of the p110 catalytic subunit by p85, and recruiting the p85-p110 heterodimer to its substrate phosphatidylinositol-4,5-bisphosphate at the plasma membrane (34). p110 phosphorylates phosphatidylinositol-4,5-bisphosphate to produce PIP₃. The interaction between p85 and tyrosine-phosphorylated adaptors permits the identification of PI3K activators by their coprecipitation with p85




Figure 4. *PTEN* loss increases PI3K activation and sensitivity to RTK ligands. **A**, p85 was immunoprecipitated from cell lysates of MCF-7 (*i*), T47D (*ii*), and MDA-361 (*iii*) lines treated overnight \pm 10% DCC-FBS. Short and long exposures are shown in *ii*. **B**, immunoblotting with the indicated antibodies of (*i*) lysates from MCF-7 cells serum-starved overnight, then treated with IGF-I (0–100 ng/mL \times 15 min.); (*ii*) lysates from T47D and MDA-361 lines treated overnight \pm 10% DCC-FBS; (*iii*) lysates from T47D cells serum-starved overnight, then treated with heregulin- β 1 (20 ng/mL \times 0–12 min).




Figure 5. PTEN loss prolongs IGF-IR and HER3 signaling, and increases E2-induced nongenomic signaling through IGF-IR. *A*, p85 was immunoprecipitated (IP) from lysates of MCF-7 cells that had been pretreated overnight with serum-free medium \pm AEW541 (1 μ mol/L), then stimulated \pm 100 ng/mL IGF-1 \pm AEW541 \times 5, 60, or 180 min. *Arrowhead*, P-IGF-IR β . *B*, lysates from MCF-7 cells pretreated as in *A* \pm AEW541 (1 μ mol/L) or lapatinib (1 μ mol/L), then stimulated \pm IGF-1 (100 ng/mL) \pm inhibitors \times 15 min. *C*, p85 was immunoprecipitated from lysates of T47D cells treated with 0.5% DCC-FBS \pm lapatinib [1 μ mol/L] \times 15, 30, 60, 120, or 180 min, or overnight (o/n). Short and long exposures for P-HER3 are shown. *D*, p85 was immunoprecipitated from lysates of MCF-7 cells pretreated overnight with 10% DCC-FBS \pm 1 μ mol/L 4-OH-T, 1 μ mol/L fulvestrant, 1 μ mol/L AEW541, or 1 μ mol/L lapatinib, and then stimulated \pm 1 nmol/L E2 \pm inhibitors \times 20 min. *Arrowhead*, tyrosine-phosphorylated IRS-1 (\sim 150 kDa). All immunoprecipitates and cell lysates were analyzed by immunoblotting with the indicated antibodies.

antibodies. Immunoprecipitation of p85 followed by immunoblot analysis showed increased p85-IRS-1 binding under basal and serum-stimulated conditions in MCF-7/shPTEN cells compared with shMM control (Fig. 4A). Similarly, T47D/shPTEN cells showed increased p85-HER3 association compared with shMM cells. PTEN loss did not significantly alter the high basal p85-HER3 association in MDA-361 cells. Additionally, MCF-7/shPTEN cells showed increased PI3K/AKT pathway sensitivity to IGF-1 ligand compared with shMM control (Fig. 4B, *i*). In contrast, PTEN loss maximally activated PI3K under basal conditions in T47D and MDA-361 cells, whereas shMM cells showed increased P-AKT upon stimulation with serum or the HER3 ligand heregulin- β 1 (Fig. 4B, *ii* and *iii*).

A time course in IGF-1-stimulated MCF-7 cell lines showed that

the association of p85 with IRS-1 upon ligand addition was enhanced by PTEN loss (Fig. 5A). P-IGF-IR β was detectable after 5 min. of IGF-1 stimulation and returned to baseline within 1 hour in MCF-7/shMM cells. In contrast, P-IGF-IR β and the increased p85-IRS-1 association remained detectable for \geq 3 hours after IGF-1 stimulation in MCF-7/shPTEN cells. Therefore, PTEN loss increased and prolonged the activation of IGF-IR and IRS-1. Prior work has shown that PI3K pathway activation suppresses IRS-1 expression (35). Indeed, extended IGF-1 stimulation for 24 hours modestly decreased IRS-1 levels in MCF-7/shPTEN but not MCF-7/shMM cells (Supplementary Fig. S5). Therefore, PTEN loss may promote IRS-1 down-regulation due to increased negative feedback from PI3K signaling.

Additionally, we detected IGF-I-induced tyrosine phosphorylation of HER3 (Fig. 5A), suggestive of crosstalk between IGF-IR and ErbB receptors. This phosphorylation was more robust and sustained in MCF-7/shPTEN compared with MCF-7/shMM cells and was inhibited by AEW541. IGF-I-induced activation of IGF-IR, HER3, and AKT was inhibited by either AEW541 or lapatinib, suggesting that EGFR/HER2 tyrosine kinase activity was required for HER3 activation in response to IGF-I (Supplementary Fig. S7; Fig. 5B).

PTEN loss induced maximal activation of PI3K/AKT in T47D cells (Fig. 4B). Lapatinib inhibited PI3K in these cells, suggesting that PI3K activation was EGFR/HER2/HER3 dependent (Supplementary Fig. S5; Figs. 3A–B and 4A). Therefore, we examined the temporal effect of lapatinib on HER3 inactivation and p85-HER3 interaction. T47D/shPTEN cells exhibited constitutive association between p85 and HER3, and higher levels of P-HER3 and P-AKT compared with shMM control (Supplementary Fig. S7; Fig. 5C). In T47D/shPTEN cells, p85-HER3 association, P-HER3, and P-AKT remained detectable after overnight treatment with lapatinib, compared with near-complete inhibition after 2 to 3 hours in T47D/shMM cells. These data suggest that, like for

IGF-IR, *PTEN* knockdown increases and prolongs the activation of HER3 and PI3K. *PTEN* loss did not alter the ability of lapatinib to suppress P-HER3 and p85-HER3 association in MDA-361 cells (not shown).

PTEN loss increases nongenomic estrogen signaling through IGF-IR. Nongenomic estrogen signaling to activate PI3K and MAPK has been proposed as a mechanism of resistance to hormonal therapy (22). We found that *PTEN* loss enhances an E2-induced increase in P-AKT and p85-IRS-1 interaction in MCF-7 cells (Fig. 5D). This signaling was unaffected by pretreatment with 4-OH-T or fulvestrant, although these compounds respectively increased and decreased ER protein levels. AEW541 but not lapatinib blocked p85-IRS-1 binding and P-AKT, suggesting that IGF-IR permits E2-induced PI3K activation in these cells. E2 did not increase P-AKT in T47D or MDA-361 cells (data not shown).

Combined blockade of IGF-IR and ErbB signaling inhibits *PTEN*-deficient cell growth. Treatment with AKTi, BEZ235, or AEW541 inhibited the growth of MCF-7/shPTEN cells in hormone-depleted medium and in the presence of tamoxifen or fulvestrant (Fig. 6A). In contrast, lapatinib was effective mainly when combined with 4-OH-T or fulvestrant. MCF-7/shPTEN cells showed




Figure 6. Combined inhibition of IGF-IR and EGFR/HER2 synergizes with 4-OH-T, fulvestrant, and hormone deprivation to block cell proliferation. A, MCF-7/shPTEN, (B) T47D/shPTEN, and (C) MDA-361/shPTEN cells were treated with medium containing DCC-FBS (MCF-7, 2%; T47D and MDA-361, 0.5%) as indicated [0.1 nmol/L E2 (MCF-7), 1 nmol/L E2 (T47D, MDA-361), 1 μ mol/L 4-OH-T, 1 μ mol/L fulvestrant, 1 μ mol/L AKTi, 200 nmol/L BEZ235, 1 μ mol/L AEW541, 1 μ mol/L lapatinib]. Media and drugs were refreshed every 2 to 3 d. Adherent cells were counted after 5 to 8 d. Data are presented as % untreated shMM control; columns, mean of triplicates; bars, SD. *, $P < 0.05$ by t test comparing kinase inhibitor-treated cells to control (Ctl) cells within each group. D, MCF-7/shPTEN and shMM cells transfected with siRNA against *HER3* or control (siCtl) were treated and analyzed as in A. *, $P < 0.05$ by t test compared with untreated shMM/siCtl within each group.

increased sensitivity to AEW541 and the PI3K inhibitor LY294002 compared with shMM control (Supplementary Fig. S7), suggesting that PTEN loss increases dependence upon IGF-IR and PI3K. Similarly, AKTi, BEZ235, and lapatinib significantly inhibited T47D/shPTEN (Fig. 6B) and MDA-361/shPTEN (Fig. 6C) cell growth. In these cells, AEW541 was only effective when combined with 4-OH-T or fulvestrant.

In all shPTEN lines, the combination of AEW541 plus lapatinib was significantly more inhibitory than either agent alone (Fig. 6). These results imply that in MCF-7 cells, IGF-IR mainly drives PI3K/AKT (Figs. 3–5), but EGFR/HER2/HER3 may also contribute to hormone-independent growth and antiestrogen resistance (Fig. 6A). Indeed, siRNA-mediated knockdown of *HER3* (verified in Supplementary Fig. S8) inhibited the growth of MCF-7/shPTEN and/shMM cells in hormone-depleted medium, and sensitized them to 4-OH-T and fulvestrant (Fig. 6D). The inhibitory effect of *HER3* knockdown was augmented by AEW541 treatment, implicating both *HER3* and IGF-IR in the modulation of MCF-7/shPTEN cell growth. Similarly, *HER3* drives PI3K in T47D and MDA-361 cells (Figs. 3–5), but the synergistic effect of lapatinib plus AEW541 also implicates IGF-IR in the modulation of growth of T47D/shPTEN and MDA-361/shPTEN cells (Fig. 6B–C). Because these kinase inhibitors synergized with 4-OH-T, fulvestrant, and hormone-deprivation to block cell growth, PTEN-deficient, ER+ breast cancer patients may be effectively treated with drug combinations targeting ER and RTK pathways that activate PI3K.

Discussion

Herein, we show that shRNA-mediated knockdown of *PTEN* in three ER+ breast cancer cell lines resulted in antiestrogen resistance and hormone-independent growth by both genomic and nongenomic mechanisms. PTEN loss up-regulated PI3K/AKT and enhanced IGF-IR/IRS-1 and *HER3* signaling, implicating PTEN in the modulation of RTK signaling upstream of PI3K. Notably, all cell lines used herein harbor activating mutations in *PIK3CA*, the gene encoding p110 α (2). These results suggest that (a) different signaling outputs result from PTEN loss versus *PIK3CA* mutations, and (b) PTEN loss is a more transforming event than *PIK3CA* mutations. This is consistent with reports that *PIK3CA* mutations and PTEN loss coexist in 5% to 14% of primary breast cancers (2, 23, 24). Cross-talk between IGF-IR and EGFR/HER2 was also enhanced by PTEN loss, as IGF-I–induced *HER3* phosphorylation was inhibited by lapatinib (Supplementary Fig. S7; Fig. 5B). Combined inhibition of these RTKs with tyrosine kinase inhibitors and/or RNAi most effectively inhibited PI3K activation (Fig. 3B) and cell growth (Fig. 6).

The lipid phosphatase activity of PTEN mediates its tumor suppressive function through dephosphorylation of PIP₃. However, evidence also suggests a tumor suppressor role for PTEN protein phosphatase activity. First, a *PTEN*_{G129E} mutation was found in two kindreds of Cowden's disease, a cancer predisposition disorder. PTEN_{G129E} lacks lipid phosphatase activity but retains protein phosphatase activity (36), inhibits cell migration (5) and epithelial-to-mesenchymal transition (37), and promotes G₁ cell cycle arrest (38). Second, we found that PTEN loss dysregulates kinase signaling upstream of PI3K. PTEN directly modulates tyrosine phosphorylation of EGFR and platelet-derived growth factor receptor (6), and regulates IRS-1 activation and InsR-IRS-1 binding (39). We showed that PTEN loss increases and prolongs IGF-IR and *HER3* tyrosine-phosphorylation, p85-IRS-1 and p85-*HER3* binding,

and E2-induced, IGF-IR/IRS-1–dependent activation of PI3K/AKT (Supplementary Fig. S7; Figs. 3–5). Furthermore, PTEN levels were inversely correlated with P-IGF-IR β in hormone receptor–positive breast cancers (Fig. 3C).

These findings collectively suggest that PTEN modulates RTKs and adaptors that activate PI3K, implicating PTEN in regulatory processes both upstream and downstream of PI3K. Because PTEN loss increases RTK activation and sensitivity to RTK ligands (Figs. 3–5), we speculate that other RTK-initiated signaling pathways besides PI3K are likely to also be activated in PTEN-deficient cells. Although the mechanism(s) by which PTEN regulates IGF-IR and *HER3* remains unclear, possibilities include genomic effects, feedback signaling to RTKs or their adaptors (39), modulation of RTK ligand production, or PTEN binding and/or dephosphorylation of RTKs or adaptor proteins such as focal adhesion kinase (5). Notably, focal adhesion kinase is a PTEN substrate (5), and focal adhesion kinase phosphorylation has been implicated in IGF-IR (40), *HER2/HER3* (41, 42), EGFR, and platelet-derived growth factor receptor signaling (43). To explore changes in RTK ligand production upon PTEN loss in MCF-7 cells, we used reverse transcription-PCR and ELISA assays for IGF-I and IGF-II, but we found no change in mRNA or protein levels (data not shown).

We show that IGF-IR/ErbB cross-talk is enhanced by PTEN loss. IGF-I stimulation increased P-*HER3* in direct correlation with P-IGF-IR β , and these effects were prolonged in PTEN-deficient cells (Fig. 5A). These responses were blocked by lapatinib or AEW541 (Supplementary Fig. S7; Fig. 5B), suggesting that IGF-IR and EGFR/HER2 kinases were required. IGF-IR has been shown to complex with P-*HER2* upon stimulation with IGF-I or heregulin (44). IGF-IR has also been shown to activate EGFR (45). Interestingly, lapatinib suppressed IGF-I–induced IGF-IR phosphorylation (Supplementary Fig. S7; Fig. 5B), suggesting that EGFR and/or *HER2* kinases are also permissive of IGF-IR activation. This RTK cross-talk supports the synergistic inhibitory effect of AEW541 plus either lapatinib or *HER3* knockdown on shPTEN cell growth (Fig. 6) and PI3K activation (Fig. 3B). We speculate that PTEN-deficient cancer cells use an alternative RTK pathway(s) to drive growth and survival when the primary PI3K-activating RTK is blocked. Dissecting the roles of IGF-IR and ErbB RTKs in heteromeric complex formation, stability, and signaling requires further evaluation, but these data suggest that inhibition of one RTK type (EGFR/HER2) may disrupt signaling of another RTK type (IGF-IR), and that combinations of tyrosine kinase inhibitors should be considered for the treatment of PTEN-deficient cancers.

Breast cancer cells may maintain a balance between ER and RTK pathways. Upon treatment with 4-OH-T or hormone deprivation, breast cancer cells adapt by up-regulating RTK network components (e.g., P-AKT, P-MAPK, RTK ligands; ref. 46). In turn, RTK pathway activation modulates ER function. For example, activation of the Src tyrosine kinase can induce ER degradation (47). Prior findings with MCF-7 cells suggest that PI3K activation suppresses ER expression while increasing transcriptional activity (14, 33). However, we found that PTEN loss decreased both ER protein levels and transcriptional activity in T47D and MDA-361 cells (Supplementary Fig. S2; Fig. 1A and 2B), indicating that the increase in ER transcriptional activity observed upon PTEN loss in MCF-7 cells (Supplementary Fig. S2; Fig. 2B) is not widely applicable to all hormone-dependent breast cancers. The decreased levels of PR seen upon *PTEN* knockdown in MCF-7 and T47D cells (Fig. 1A) may be due to increased PR degradation or decreased

gene expression (determined by microarray analysis; data not shown). PI3K pathway activation has been shown to increase the expression of the E2-inducible gene *Cyclin D1* while suppressing *PgR* expression (48). Furthermore, heregulin- β 1 stimulation of T47D cells increased PR phosphorylation at Ser₂₉₄ (49), which, in turn, promotes PR degradation (50). Therefore, PI3K activation as a result of PTEN loss may up-regulate or down-regulate transcription of ER target genes, as well as modulate their gene products posttranslationally.

In summary, we report that induced PTEN loss in hormone-dependent human breast cancer cells resulted in (a) hormone-independent growth and resistance to 4-OH-T and fulvestrant, (b) variable alterations in ER transcriptional activity, and (c) up-regulation of kinase signaling upstream and downstream of PI3K. Furthermore, we showed that *PTEN* knockdown in three *PIK3CA*-mutant breast cancer cell lines confers gain-of-function effects, indicating different signaling outputs as a result of PTEN loss versus *PIK3CA* mutation. The use of three cell lines allowed us to discover pathways commonly dysregulated by PTEN loss in different systems, and we found that PTEN loss up-regulates intact pathways upstream of PI3K. Inhibition of IGF-IR and ErbB signaling synergized with 4-OH-T, fulvestrant, and hormone-

deprivation to overcome the growth advantage conferred by PTEN loss. These data hold promise for the treatment of PTEN-deficient, ER+ breast cancer patients with combinations of drugs targeting both ER and RTK/PI3K pathways.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

Acknowledgments

Received 1/13/09; revised 3/20/09; accepted 3/23/09.

Grant support: NIH R01CA80195 (C.L. Arteaga), F32CA121900 (T.W. Miller), T32CA78136 (M.P. Torres), K23CA121994, R21CA120248 (A.M. Gonzalez), Breast Cancer Specialized Program of Research Excellence P50CA98131, and Vanderbilt-Ingram Comprehensive Cancer Center Support Grant P30CA68485; the Breast Cancer Research Foundation (C.L. Arteaga); ACS Clinical Research Professorship Grant CRP-07-234 (C.L. Arteaga); Kleberg Center for Molecular Markers, M. D. Anderson Cancer Center; American Society of Clinical Oncology Career Development Award (A.M. Gonzalez); M. D. Anderson [Physician Scientist Award (B.T. Hennessy)]; Susan G. Komen Breast Cancer Foundation [EAS0703849 (B.T. Hennessy, A.M. Gonzalez, and G.B. Mills)]; American Cancer Society [IRG-58-009-49 (C.W. Lindsley)]; and Lee Jeans Translational Breast Cancer Research Program (C.L. Arteaga).

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

References

- Li J, Yen C, Liaw D, et al. PTEN, a putative protein tyrosine phosphatase gene mutated in human brain, breast, and prostate cancer. *Science* 1997;275:1943-7.
- Saal LH, Holm K, Maurer M, et al. PIK3CA mutations correlate with hormone receptors, node metastasis, and ERBB2, and are mutually exclusive with PTEN loss in human breast carcinoma. *Cancer Res* 2005;65:2554-9.
- Depowski PL, Rosenthal SI, Ross JS. Loss of expression of the PTEN gene protein product is associated with poor outcome in breast cancer. *Mod Pathol* 2001;14:672-6.
- Perren A, Weng LP, Boag AH, et al. Immunohistochemical evidence of loss of PTEN expression in primary ductal adenocarcinomas of the breast. *Am J Pathol* 1999;155:1253-60.
- Tamura M, Gu J, Matsumoto K, Aota S, Parsons R, Yamada KM. Inhibition of cell migration, spreading, and focal adhesions by tumor suppressor PTEN. *Science* 1998;280:1614-7.
- Mahimainathan L, Choudhury GG. Inactivation of platelet-derived growth factor receptor by the tumor suppressor PTEN provides a novel mechanism of action of the phosphatase. *J Biol Chem* 2004;279:15258-68.
- Raftopoulos M, Etienne-Manneville S, Self A, Nicholls S, Hall A. Regulation of cell migration by the C2 domain of the tumor suppressor PTEN. *Science* 2004;303:1179-81.
- Freeman DJ, Li AG, Wei G, et al. PTEN tumor suppressor regulates p53 protein levels and activity through phosphatase-dependent and -independent mechanisms. *Cancer Cell* 2003;3:117-30.
- Arpino G, Wiechmann L, Osborne CK, Schiff R. Crosstalk between the estrogen receptor and the HER tyrosine kinase receptor family: molecular mechanism and clinical implications for endocrine therapy resistance. *Endocr Rev* 2008;29:217-33.
- Folgiero V, Avetrani P, Bon G, et al. Induction of ErbB-3 expression by α 6 β 4 integrin contributes to tamoxifen resistance in ER β 1-negative breast carcinomas. *PLoS ONE* 2008;3:e1592.
- Kurokawa H, Arteaga CL. ErbB (HER) receptors can abrogate antiestrogen action in human breast cancer by multiple signaling mechanisms. *Clin Cancer Res* 2003;9:511-55.
- Massarweh S, Schiff R. Unraveling the mechanisms of endocrine resistance in breast cancer: new therapeutic opportunities. *Clin Cancer Res* 2007;13:1950-4.
- Kato S, Endoh H, Masuhiro Y, et al. Activation of the estrogen receptor through phosphorylation by mitogen-activated protein kinase. *Science* 1995;270:1491-4.
- Martin MB, Franke TF, Stoica GE, et al. A role for Akt in mediating the estrogenic functions of epidermal growth factor and insulin-like growth factor I. *Endocrinology* 2000;141:4503-11.
- Font de Mora J, Brown M. AIB1 is a conduit for kinase-mediated growth factor signaling to the estrogen receptor. *Mol Cell Biol* 2000;20:5041-7.
- Hadari YR, Tzahar E, Nativ O, et al. Insulin and insulinomimetic agents induce activation of phosphatidylinositol 3'-kinase upon its association with pp185 (IRS-1) in intact rat livers. *J Biol Chem* 1992;267:17483-6.
- Morelli C, Garofalo C, Bartucci M, Surmacz E. Estrogen receptor- α regulates the degradation of insulin receptor substrates 1 and 2 in breast cancer cells. *Oncogene* 2003;22:4007-16.
- Frasor J, Stossi F, Danes JM, Komm B, Lyttle CR, Katzenellenbogen BS. Selective estrogen receptor modulators: discrimination of agonistic versus antagonistic activities by gene expression profiling in breast cancer cells. *Cancer Res* 2004;64:1522-33.
- Reddy KB, Yee D, Hilsenbeck SG, Coffey RJ, Osborne CK. Inhibition of estrogen-induced breast cancer cell proliferation by reduction in autocrine transforming growth factor α expression. *Cell Growth Differ* 1994;5:1275-82.
- Losel R, Wehling M. Nongenomic actions of steroid hormones. *Nat Rev Mol Cell Biol* 2003;4:46-56.
- Simoncini T, Hafezi-Moghadam A, Brazil DP, Ley K, Chin WW, Liao JK. Interaction of estrogen receptor with the regulatory subunit of phosphatidylinositol-3-OH kinase. *Nature* 2000;407:538-41.
- Song RX, Fan P, Yue W, Chen Y, Santen RJ. Role of receptor complexes in the extranuclear actions of estrogen receptor α in breast cancer. *Endocr Relat Cancer* 2006;13 Suppl 1:53-13.
- Perez-Tenorio G, Alkhorri L, Olsson B, et al. PIK3CA mutations and PTEN loss correlate with similar prognostic factors and are not mutually exclusive in breast cancer. *Clin Cancer Res* 2007;13:3577-84.
- Saal LH, Johansson P, Holm K, et al. Poor prognosis in carcinoma is associated with a gene expression signature of aberrant PTEN tumor suppressor pathway activity. *Proc Natl Acad Sci U S A* 2007;104:7564-9.
- Yi JY, Shin I, Arteaga CL. Type I transforming growth factor β receptor binds to and activates phosphatidylinositol 3-kinase. *J Biol Chem* 2005;280:10870-6.
- Lindsley CW, Zhao Z, Leister WH, et al. Allosteric Akt (PKB) inhibitors: discovery and SAR of isozyme selective inhibitors. *Bioorg Med Chem Lett* 2005;15:761-4.
- Maira SM, Stauffer F, Brueggen J, et al. Identification and characterization of NVP-BEZ235, a new orally available dual phosphatidylinositol 3-kinase/mammalian target of rapamycin inhibitor with potent *in vivo* antitumor activity. *Mol Cancer Ther* 2008;7:1851-63.
- Garcia-Echeverria C, Pearson MA, Marti A, et al. *In vivo* antitumor activity of NVP-AEW541-A novel, potent, and selective inhibitor of the IGF-IR kinase. *Cancer Cell* 2004;5:231-9.
- Kurokawa H, Lenferink AE, Simpson JF, et al. Inhibition of HER2/neu (erbB-2) and mitogen-activated protein kinases enhances tamoxifen action against HER2-overexpressing, tamoxifen-resistant breast cancer cells. *Cancer Res* 2000;60:5887-94.
- Engelman JA, Janne PA, Mermel C, et al. ErbB-3 mediates phosphoinositide 3-kinase activity in gefitinib-sensitive non-small cell lung cancer cell lines. *Proc Natl Acad Sci U S A* 2005;102:3788-93.
- Tibes R, Qiu Y, Lu Y, et al. Reverse phase protein array: validation of a novel proteomic technology and utility for analysis of primary leukemia specimens and hematopoietic stem cells. *Mol Cancer Ther* 2006;5:2512-21.
- MacIndoe JH. Estradiol formation from testosterone by continuously cultured human breast cancer cells. *J Clin Endocrinol Metab* 1979;49:272-7.
- Stoica GE, Franke TF, Moroni M, et al. Effect of estradiol on estrogen receptor- α gene expression and activity can be modulated by the ErbB2/PI 3-K/Akt pathway. *Oncogene* 2003;22:7998-8011.
- Engelman JA, Luo J, Cantley LC. The evolution of phosphatidylinositol 3-kinases as regulators of growth and metabolism. *Nat Rev Genet* 2006;7:606-19.
- Lee AV, Gooch JL, Oesterreich S, Guler RL, Yee D. Insulin-like growth factor I-induced degradation of insulin receptor substrate 1 is mediated by the 26S proteasome and blocked by phosphatidylinositol 3'-kinase inhibition. *Mol Cell Biol* 2000;20:1489-96.
- Myers MP, Stolarov JP, Eng C, et al. P-TEN, the tumor suppressor from human chromosome 10q23, is a dual-specificity phosphatase. *Proc Natl Acad Sci U S A* 1997;94:9052-7.
- Leslie NR, Yang X, Downes CP, Weijer CJ. PtdIns(3,4,5)P(3)-dependent and -independent roles for PTEN in the control of cell migration. *Curr Biol* 2007;17:115-25.

38. Hlobilkova A, Guldberg P, Thullberg M, Zeuthen J, Lukas J, Bartek J. Cell cycle arrest by the PTEN tumor suppressor is target cell specific and may require protein phosphatase activity. *Exp Cell Res* 2000;256:571-7.
39. Vivanco I, Palaskas N, Tran C, et al. Identification of the JNK signaling pathway as a functional target of the tumor suppressor PTEN. *Cancer Cell* 2007;11:555-69.
40. Liu W, Bloom DA, Cance WG, Kurenova EV, Golubovskaya VM, Hochwald SN. Fak and Igf-Ir interact to provide survival signals in human pancreatic adenocarcinoma cells. *Carcinogenesis* 2008;29:1096-107.
41. Vartanian T, Goodearl A, Lefebvre S, Park SK, Fischbach G. Neuregulin induces the rapid association of focal adhesion kinase with the erbB2-3 receptor complex in schwann cells. *Biochem Biophys Res Commun* 2000;271:414-7.
42. Benlimame N, He Q, Jie S, et al. FAK signaling is critical for ErbB-2/ErbB-3 receptor cooperation for oncogenic transformation and invasion. *J Cell Biol* 2005;171:505-16.
43. Sieg DJ, Hauck CR, Ilic D, et al. FAK integrates growth-factor and integrin signals to promote cell migration. *Nat Cell Biol* 2000;2:249-56.
44. Balana ME, Labriola L, Salatino M, et al. Activation of ErbB-2 via a hierarchical interaction between ErbB-2 and type I insulin-like growth factor receptor in mammary tumor cells. *Oncogene* 2001;20:34-47.
45. Ahmad T, Farnie G, Bundred NJ, Anderson NG. The mitogenic action of insulin-like growth factor I in normal human mammary epithelial cells requires the epidermal growth factor receptor tyrosine kinase. *J Biol Chem* 2004;279:1713-9.
46. Johnston SR, Martin LA, Leary A, Head J, Dowsett M. Clinical strategies for rationale combinations of aromatase inhibitors with novel therapies for breast cancer. *J Steroid Biochem Mol Biol* 2007;106:180-6.
47. Chu I, Arnaout A, Loiseau S, et al. Src promotes estrogen-dependent estrogen receptor α proteolysis in human breast cancer. *J Clin Invest* 2007;117:2205-15.
48. Cui X, Zhang P, Deng W, et al. Insulin-like growth factor-I inhibits progesterone receptor expression in breast cancer cells via the phosphatidylinositol 3-kinase/Akt/mammalian target of rapamycin pathway: progesterone receptor as a potential indicator of growth factor activity in breast cancer. *Mol Endocrinol* 2003;17:575-88.
49. Labriola L, Salatino M, Proietti CJ, et al. Heregulin induces transcriptional activation of the progesterone receptor by a mechanism that requires functional ErbB-2 and mitogen-activated protein kinase activation in breast cancer cells. *Mol Cell Biol* 2003;23:1095-111.
50. Lange CA, Shen T, Horwitz KB. Phosphorylation of human progesterone receptors at serine-294 by mitogen-activated protein kinase signals their degradation by the 26S proteasome. *Proc Natl Acad Sci U S A* 2000;97:1032-7.

Cancer Research

The Journal of Cancer Research (1916–1930) | The American Journal of Cancer (1931–1940)

Loss of *Phosphatase and Tensin Homologue Deleted on Chromosome 10* Engages ErbB3 and Insulin-Like Growth Factor-I Receptor Signaling to Promote Antiestrogen Resistance in Breast Cancer

Todd W. Miller, Marianela Pérez-Torres, Archana Narasanna, et al.

Cancer Res 2009;69:4192-4201. Published OnlineFirst May 12, 2009.

Updated version	Access the most recent version of this article at: doi: 10.1158/0008-5472.CAN-09-0042
Supplementary Material	Access the most recent supplemental material at: http://cancerres.aacrjournals.org/content/suppl/2009/05/11/0008-5472.CAN-09-0042.DC1

Cited articles	This article cites 50 articles, 25 of which you can access for free at: http://cancerres.aacrjournals.org/content/69/10/4192.full#ref-list-1
Citing articles	This article has been cited by 18 HighWire-hosted articles. Access the articles at: http://cancerres.aacrjournals.org/content/69/10/4192.full#related-urls

E-mail alerts	Sign up to receive free email-alerts related to this article or journal.
Reprints and Subscriptions	To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org .
Permissions	To request permission to re-use all or part of this article, use this link http://cancerres.aacrjournals.org/content/69/10/4192 . Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.